

BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE- liquid bandage liquid
Chain Drug Consortium, LLC

Premier Value Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

Warnings

For external use only. Flammable: Keep away from fire or flame.

Do not use

- in the eyes
- overlarge areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious burns

When using this product

- avoid contact with eyes

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs
- symptoms persists for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age; consult a doctor

Other information

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

Premier Value

Liquid Bandage

FOR SMALL CUTS & WOUNDS

Topical analgesic and antiseptic to help prevent infection

Antiseptic formula dries quickly to form an invisible protective cover that is flexible


0.3 FL OZ (9mL)

Use Liquid Bandage for:

blisters, chapped and cracked finger tips, hangnails, paper cuts, shaving nicks and help in preventing the formation of calluses.

Great for:

athletes, active lifestyles, outdoor enthusiasts and musicians.



Liquid Bandage

FOR SMALL CUTS & WOUNDS

Topical analgesic and antiseptic
to help prevent infection

Antiseptic formula dries quickly to form an
invisible protective cover that is flexible

NET 0.3 FL OZ (9 mL)



Use Liquid Bandage for:

blisters, chapped and cracked finger tips, hangnails, paper cuts,
shaving nicks and help in preventing the formation of calluses.

Great for:

athletes, active lifestyles, outdoor enthusiasts and musicians.

53-033PV-02



Liquid Bandage

FOR SMALL CUTS & WOUNDS
Topical analgesic and antiseptic
to help prevent infection

NET 0.3 FL OZ (9 mL)



NO VARNISH
AREA



If for any reason you are not satisfied with
this product, please return it to the store
where purchased for a full refund.

Distributed By:
Pharmacy Value Alliance, LLC
407 East Lancaster Avenue
Wayne, PA 19087

Made in USA with U.S. and imported parts



DO NOT USE IF NECKBAND IS BROKEN

Drug Facts

Active Ingredients
Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%
First aid antiseptic
Topical analgesic

Purpose
First aid to prevent bacterial contamination in minor cuts, scrapes and burns

Uses
First aid to prevent bacterial contamination in minor cuts, scrapes and burns

Warnings
For external use only. Flammable: Keep away from fire or flame.
Do not use in the eyes over large areas of the body
longer than 1 week unless directed by a doctor
Ask a doctor before use if you have deep or puncture wounds
animal bites serious burns
When using this product avoid contact with eyes
Stop use and ask a doctor if condition persists or gets worse infection occurs
symptoms persist for more than 7 days or clear up and occur again within a few days
Keep out of reach of children. If swallowed, get medical help or call a Poison
Control Center right away.

Directions
for adults and children 2 years of age and older
clean affected area
apply a small amount on the area 1 to 3 times daily and let dry
a second coating
may be applied for extra protection
children under 2 years of age: consult a doctor
Other information cap bottle tightly store between 20° to 25°C
(68° to 77°F) may temporarily sting upon application to remove, apply more
Liquid Bandage and quickly wipe off

Inactive ingredients acetone, amyl acetate, castor oil, ethyl acetate,
nitrocellulose, SD alcohol 40

Questions? Call 1-866-964-0939

This product is not manufactured or distributed by Moberg Pharma North America, LLC,
owner of the registered trademark New-Skin®.



BENZETHONIUM CHLORIDE PLUS DYCLONINE HYDROCHLORIDE

liquid bandage liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-615
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 9 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 9 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-	9 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a	12/07/2005	

615-00	Combination Product	12/07/2005	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	12/07/2005	

Labeler - Chain Drug Consortium, LLC (101668460)

Revised: 12/2024

Chain Drug Consortium, LLC