

ASPIRIN 325 MG- aspirin tablet
WALGREENS

331R Walgreens 0363-6453 Aspirin 325 mg

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- For the temporary relief of minor aches and pains due to:
 - o headache
 - o toothache
 - o muscle pain
 - o minor arthritis pain
 - o menstrual pain
 - o colds
- or as recommended by a doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDS (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use:

- If you have ever had an allergic reaction to any other pain reliever/fever reducer.

Ask a doctor before use if:

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis.

Stop use and ask a doctor if:

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs

Seek medical help right away if:

- new symptoms occur
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose.
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor.
- children under 12 years: ask a doctor.

Other information

- **each tablet contains:** sodium 1 mg (VERY LOW SODIUM)
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).
- use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD & C yellow #6 aluminum lake, hypromellose, methacrylic acid and ethyl acrylate

Questions or comments? Call 1-877-290-4008

STOP PEELING	<p>Drug Facts (continued)</p> <p>Do not use: ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer.</p> <p>Ask a doctor before use if:</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ taking a prescription drug for diabetes, gout, or arthritis. <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ an allergic reaction occurs <p>Seek medical help right away if:</p> <ul style="list-style-type: none"> ■ new symptoms occur ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days 	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ drink a full glass of water with each dose. ■ adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor. ■ children under 12 years: ask a doctor. <p>Other information</p> <ul style="list-style-type: none"> ■ each tablet contains: sodium 1 mg (VERY LOW SODIUM) ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F). ■ use by expiration date on package <p>Inactive ingredients corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD & C yellow #6 aluminum lake, hypromellose, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate</p>
	<p>Questions or comments? Call 1-877-290-4008</p>	<p>Questions or comments? Call 1-877-290-4008</p>

aspirin tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6453
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MINERAL OIL (UNII: T5L8T28FGP)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)				
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)				
SODIUM HYDROXIDE (UNII: 55X04QC32I)				
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)				
STARCH, CORN (UNII: O8232NY3SJ)				
TALC (UNII: 7SEV7J4R1U)				
Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	T	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6453-53	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		06/15/2022	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-6453)

Revised: 7/2022

WALGREENS