

**DYE-FREE ASPIRIN 81- aspirin tablet
WALGREENS**

480R Walgreens 0363-6455 Dye-Free Aspirin 81 mg

DRUG FACTS

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: •hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problem
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn

- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present

new symptoms occur

ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments?

Call **1-877-290-4008**

Walgreens
free+pure[†]

NDC 0363-6455-10

Heart health[†]

Aspirin 81

81 mg, pain reliever (NSAID)

Compare to Bayer[®] Low Dose Aspirin^{††}

Low dose | Enteric-coated

[†]Free + Pure is:

- No artificial flavors
- No synthetic colors
- Gluten free^{††}
- Non-GMO

^{††}Talk to your doctor or other healthcare provider before using this product for your heart.

1000
enteric-coated
tablets

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)^{**}.....Pain reliever
^{**}nonsteroidal anti-inflammatory drug

Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

Warnings

- **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
- **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
- **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Drug Facts (continued under label)

Walgreens Pharmacist Recommended. 480R 1023 WFFORG112S-F
[†]We invite you to compare to national brands. Walgreens is a registered trademark and/or distributed by Bayer AG, owner of the registered trademark Bayer.
^{††}Contains less than 20ppm of gluten
DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
WALGREENS ©2023 Walgreen Co.



Lot No.:

Exp Date:



NDC 0363-6455-12

Walgreens

Dye-Free Aspirin 81

81 mg / PAIN RELIEVER (NSAID)

Low Dose

Enteric-Coated

120 TABLETS

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)^{**}.....Pain reliever
^{**}nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.** ■ ask your doctor about other uses for enteric-coated 81 mg Aspirin.

Warnings (See carton for complete warnings information)

- **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
- **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
- **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 - are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.** **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other Information

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
480R 0222
Store at room temperature. Avoid excessive heat above 40°C (104°F)
DISTRIBUTED BY: WALGREEN CO. ITEM 285144
W00000-0000-0
100% SATISFACTION GUARANTEED
W3ORG0422-F
WALGREENS.COM ©2022 Walgreen Co.
MADE IN U.S.A. OF U.S. AND IMPORTED INGREDIENTS

Lot No.:

Exp. Date:

Varnish Omit Area

Lot No.:

Exp Date:



Drug Facts (continued)

Ask a doctor before use if

- you have stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have taken a prescription drug for:
 - gout
 - diabetes
 - arthritis

Stop use and ask a doctor if

- you have an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - heartburn
 - vomit blood
 - have bloody or black stools
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.** **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

Store at room temperature. Avoid excessive heat above 40°C (104°F)
Inactive ingredients: anhydrous lactose, canola wax, colloidal silicon dioxide, croscarmellose sodium, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polypropylene 80, stearic acid, sodium hydroxide, sodium lauryl sulfate, stearic acid, talc, triethyl citrate

Questions or comments? Call 1-877-296-4008

ITEM 285144 W00000-0000-0
 3 11917 13645 5

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING. REPRINTED FOL SEAL OVER THE MOUTH OF THE BOTTLE.



Drug Facts

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Active ingredient Purpose (in each tablet)

Aspirin 81 mg (NSAID)*Pain reliever
 *nonsteroidal anti-inflammatory drug

Uses ■ for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

■ ask your doctor about other uses for enteric coated 81 mg Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling
- asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

Drug Facts (continued)

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn ■ you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.

- you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important **not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- Store at room temperature. Avoid excessive heat above 40°C (104°F)

Inactive ingredients anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

NDC 0363-8465-12



Walgreens
 Compare to the active ingredient in Bayer® Low Dose Aspirin®



Dye-Free Aspirin 81
 81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

120 TABLETS

ACTUAL SIZE
 **Talk to your doctor or other healthcare provider before using this product for your heart.

W3ORG0422-F

Drug Facts (continued)
Questions or comments?
 Call 1-877-290-4008

Our pharmacist recommends the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.
 DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED 480R 0222 walgreens.com ©2022 Walgreen Co. MADE IN U.S.A. OR U.S. AND IMPORTED INGREDIENTS

VALUE SIZE NDC 0363-6455-50

Walgreens
Compare to the active ingredient in Bayer® Low Dose Aspirin®

Dye-Free Aspirin 81
81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

500 TABLETS ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN

HEART HEALTH ⁺₊

Drug Facts
Active ingredient (in each tablet)
Aspirin 81 mg (NSAID)*Pain reliever
*nonsteroidal anti-inflammatory drug

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma
■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
*This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.
480F 0222 W3ORG0422-F ITEM 285351 W00000-0000-0
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
Walgreens.com ©2022 Walgreen Co.
100% SATISFACTION GUARANTEED
Walgreens is a U.S. PATENTED
IMPORTED INGREDIENTS
Lot No.: EXP. DATE:
PEEL HERE

Varnish Omit Area

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
● feel faint ● vomit blood ● have bloody or black stools
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
■ children under 12 years: consult a doctor

Other information
■ Store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients
anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysoxate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, triethyl citrate

Questions or comments? Call 1-877-290-4008

STOP PEELING

DYE-FREE ASPIRIN 81

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6455
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TALC (UNII: 7SEV7J4R1U)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6455-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2022	
2	NDC:0363-6455-12	120 in 1 CARTON	06/15/2022	
2		120 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0363-6455-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	01/15/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/15/2022	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(0363-6455)

Revised: 12/2023

WALGREENS