

ULINE IBUPROFEN- ibuprofen tablet
ULINE IBUPROFEN- ibuprofen tablet, coated
Uline

Uline Ibuprofen

Drug Facts

Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

■ headache ■ toothache ■ backache ■ menstrual cramps ■ common cold ■
muscular aches ■ minor arthritis pain

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■
blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer

- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
- you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children: (12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Ask a doctor.

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 40°C (above 104°F)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone K30*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide

*may contain

Questions or comments? 1-800-295-5510

Uline

Tamper evident sealed packets:

Do not use if packet is open or torn

This package is for households without young children

Compare to Motrin IB*

Ibuprofen


Pull to Open

Ibuprofen 200 mg (NSAID*)

- Pain Reliever
- Fevr Reducer

50 Packages

2 Tablets Each



Uline

Tamper evident sealed packets:

Do not use if packet is open or torn

This package is for households without young children

Compare to Motrin IB*

Ibuprofen


Pull to Open

Ibuprofen 200 mg (NSAID*)

- Pain Reliever
- Fever Reducer

50 Packages

2 Tablets Each



ULINE IBUPROFEN

ibuprofen tablet

| Product Information | |
|-------------------------|----------------|
| Product Type | HUMAN OTC DRUG |
| Route of Administration | ORAL |
| Item Code (Source) | NDC:69790-709 |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| POVIDONE K30 (UNII: U725QWY32X) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TALC (UNII: 7SEV7J4R1U) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | brown | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | G;2 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69790-709-33 | 50 in 1 BOX | 04/04/2022 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA079174 | 04/04/2022 | |

ULINE IBUPROFEN

ibuprofen tablet, coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:69790-609 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | brown | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | 44;291 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69790-609-33 | 50 in 1 BOX | 04/04/2022 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA075010 | 04/04/2022 | |

Labeler - Uline (039612668)

Revised: 9/2025

Uline