# BURKHART- sodium fluoride gel Burkhart Dental Supply Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

## **Active Ingredient:**

Fluoride Ion 1.23%

Available from 2.09% Sodium Fluoride and Hydrofluoric Acid

## **Purpose:**

Fluoride Treatment Gel

## Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Use Only. This product is not intended for home or unsupervised use.

## **Warnings:**

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

# **Dosage and Administration:**

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

- 1. After thorough prophylaxsis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
- 2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
- 3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

### Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C).

## **Inactive Ingredients:**

Citric Acid, FD&C Blue #1, FD&C Red #40, FD&C Yellow #5 (tartrazine) as a color additive, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

# **APF GEL**

Acidulated Phosphate Fluoride Treatment Gel with Xylitol

> 1.23% Fluoride Ion Gluten Free

> > Ronly

PIÑA COLADA



Reorder No. 72721424



Active Ingredient:

Purpose:

Fluoride Ion 1.23%......Fluoride Treatment Gel Available from 2.09% Sodium Fluoride and Hydrofluoric Acid.

### **Indications and Usage:**

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Office Use Only. This product is not intended for home or unsupervised consumer use.

#### Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- · Read directions carefully before using.

### Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

- After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
- 2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
- 3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

### Other Information:

 $\bullet$  Store at controlled room temperature 59° to 86°F (15°-30°C). Protect from freezing.

### **Inactive Ingredients:**

Citric Acid, FD&C Blue #1, FD&C Red #40, FD&C Yellow #5 (tartrazine) as a color additive, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

Net wt. 16 oz (454 g)

MADE IN THE USA

Manufactured for Burkhart Dental Supply 2502 S. 78th Street Tacoma, Washington 98409 (800) 562-8176 www.burkhartdental.com

99092604 Rev 9/13

## **BURKHART**

sodium fluoride gel

## **Product Information**

**Product Type** 

HUMAN PRESCRIPTION DRUG

Item Code (Source)

NDC:43498-104

**Route of Administration** 

DENTAL

<b>Active</b>	Ingredient/Active	Moiety
---------------	-------------------	--------

Ingredient Name	Basis of Strength	Strength
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU4080)	FLUORIDE ION	5.6 g in 454 g

Inactive Ingredients		
Ingredient Name	Strength	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)		
PHOSPHORIC ACID (UNII: E4GA8884NN)		
POLYSORBATE 20 (UNII: 7T1F30V5YH)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
WATER (UNII: 059QF0KO0R)		
XANTHAN GUM (UNII: TTV12P4NEE)		
XYLITOL (UNII: VCQ006KQ1E)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C YELLOW NO. 5 (UNII: I753WB2F1M)		

Product Characteristics		
Color	yellow	Score
Shape		Size
Flavor	PINEAPPLE (Pina Colada)	Imprint Code
Contains		

l	P	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:43498- 104-15	454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2013	

Marketing Information		
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	12/14/2012	
	• •	Citation Date

Revised: 2/2022 Burkhart Dental Supply Inc