

MAX RELIEF JUNIOR- children acetaminophen suspension
Akron Pharma Inc.

MAX relief junior
Children's Pain Reliever
Acetaminophen Oral Suspension
160mg per 5mL

Dye-Free

Active Ingredient

| Active ingredient (in each 5mL) | Purpose |
|--|-----------------------------|
| Acetaminophen 160 mg | Pain reliever/fever reducer |

Uses

temporarily: reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:

if sore throat is severe, persists or more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor or pharmacist before use if the child has

liver disease.

Ask doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin. **When using this product do not exceed recommended dose (see overdose warning)**

Stop use and ask a doctor if

- pain gets worse or lasts for more than 5 days
- fever gets worse or last for more than 3 days
- new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Overdose warning:

In case of overdose, get medical help to contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Keep out of Reach of children

Directions

- this product does not contain directions or complete warnings for adult use
- shake well before using
- mL = milliliter
- find right dose on chart below. if possible, use weight to dose, otherwise use age
- use only the enclosed dosing cup designed for use with product.

| Weight (lbs) | Age (yrs.) | Dose (tsp or mL) | Weight (lbs) | Age (yrs.) | Dose (tsp or mL) |
|--------------|------------|---------------------|--------------|------------|----------------------|
| under 24 | under 2 | ask a doctor | 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 25 to 35 | 2 to 3 | 1 tsp or 5 mL | 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL | 72 to 95 | 11 | 2 1/2 tsp or 12.5 mL |

Other Information

- Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Inactive Ingredients

Anhydrous Citric Acid, Cherry Flavor, Glycerin, Microcrystalline cellulose and Sodium Methyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.

Questions or comments?

call toll-free 1-877-225-6999

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

NDC 71399-0038-4

MAXrelief™
junior



Dye-Free
Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER

Acetaminophen
Oral Suspension

160mg per 5mL

Cherry
flavor

Akron Pharma

4 FL.OZ (118 mL)

Drug Facts

Active ingredient (in each 5 mL)
Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold
■ flu ■ headache ■ sore throat ■ toothache

Warning Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists or more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor use if the child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present. These could be signs of a serious condition. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use ■ shake well before using ■ mL = milliliter ■ find ring dose on chart below. If possible, use weight to dose; otherwise, use age. ■ use only the enclosed dosing cup designed for use with product.

| Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) | Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) |
|---------------|------------|---------------------|---------------|------------|----------------------|
| under 24 | under 2 | ask a doctor | 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 24 to 35 | 2 to 3 | 1 tsp or 5 mL | 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL | 72 to 95 | 11 | 2 1/2 tsp or 12.5 mL |

Other information store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients Anhydrous Citric Acid, Cherry Flavor, Glycerin, Microcrystalline cellulose and Sodium Carboxy Methyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.


Questions or comments? call toll-free 1-877-225-6999.

*** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING

DRUGS/AP/12297

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com



MRS-1/23/10

NDC 71399-0038-4

MAXrelief[™]
Dye-Free
junior

Children's
PAIN RELIEVER
and fever Reducer

Acetaminophen
Oral Suspension

160mg per 5mL

PAIN+FEVER



Akron Pharma

4 FL.OZ (118 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor use if the child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use ■ shake well before using ■ mL—milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.

NDC 71399-0038-4

MAXrelief[™]
Dye-Free
junior

Children's
PAIN RELIEVER
and fever Reducer

Acetaminophen
Oral Suspension

160mg per 5mL

PAIN+FEVER



Akron Pharma

4 FL.OZ (118 mL)



Drug Facts (continued)

■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by doctor.

| Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) |
|---------------|------------|----------------------|
| under 24 | under 2 | ask a doctor |
| 24 to 35 | 2 to 3 | 1 tsp or 5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL |
| 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 72 to 95 | 11 | 2 1/2 tsp or 12.5 mL |

Other information
store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive ingredients: Anhydrous Citric Acid, Cherry Flavor, Glycerin, Microcrystalline cellulose and Sodium Carboxymethyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DRUGS/W/12097

Akron Pharma

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com



NDC 71399-0038-8

MAXrelief[™]
Dye-Free
junior

Children's
PAIN RELIEVER
and fever Reducer

Acetaminophen
Oral Suspension

160mg per 5mL

PAIN+FEVER

Akron Pharma

8 FL.OZ (236 mL)



Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL—milliliter ■ find right dose on chart below. If possible, use weight to dose; other wise, use age. ■ use only the enclosed dosing cup designed for use with product.

Drug Facts (continued)

Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours.

| Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) |
|---------------|------------|----------------------|
| under 24 | under 2 | ask a doctor |
| 24 to 35 | 2 to 3 | 1 tsp or 5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL |
| 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 72 to 95 | 11 | 3 tsp or 15 mL |


Other information
store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive ingredients: anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.



Akron Pharma

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com



Drug Facts

Active ingredient (in each 5 mL) *Purpose*
Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to:
■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if
■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
■ use only the enclosed dosing cup designed for

NDC 71399-0038-6

MAXrelief[™] junior

Dye-Free
Children's
PAIN RELIEVER
and fever Reducer

PAIN + FEVER

Acetaminophen Oral Suspension
160mg per 5mL

Cherry Flavor



Akron Pharma

16 FL.OZ (473 mL)

Drug Facts (continued)

use with product. Do not use any other dosing device.
■ if needed, repeat dose every 4 hours while symptoms last
■ do not give more than 5 times in 24 hours
■ do not give more than 5 days unless directed by doctor.

| Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) |
|---------------|------------|----------------------|
| under 24 | under 2 | ask a doctor |
| 24 to 35 | 2 to 3 | 1 tsp or 5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL |
| 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 72 to 95 | 11 | 3 tsp or 15 mL |

*or as directed by a doctor

Other Information
store at room temperature 20°-25°C (68-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethyl cellulose, potassium sorbate, purified water, sorbitol, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.



Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com

N 3 71399 00386 7


NDC 71399-0038-8

MAXrelief[™] junior

Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen Oral Suspension
160mg per 5mL

PAIN + FEVER

Cherry Flavor



Akron Pharma

8 FL.OZ (236 mL)

Size : 58x160x58mm Lock Bottom

Drug Facts

Active ingredient (in each 5 mL) *Purpose*
Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
■ more than 5 doses in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions
■ this product does not contain directions or complete warnings for adult use, shake well before using
■ mL = milliliter
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.
■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.
■ if needed, repeat dose every 4 hours while symptoms last

NDC 71399-0038-8

MAXrelief[™] junior

Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen Oral Suspension
160mg per 5mL

PAIN + FEVER

Cherry Flavor



Akron Pharma

8 FL.OZ (236 mL)

Drug Facts (continued)

■ do not give more than 5 times in 24 hours
■ do not give more than 5 days unless directed by doctor.
■ do not give more than directed (see overdose warning)

| Weight (lbs.) | Age (yrs.) | Dose (tsp or mL) |
|---------------|------------|----------------------|
| under 24 | under 2 | ask a doctor |
| 24 to 35 | 2 to 3 | 1 tsp or 5 mL |
| 36 to 47 | 4 to 5 | 1 1/2 tsp or 7.5 mL |
| 48 to 59 | 6 to 8 | 2 tsp or 10 mL |
| 60 to 71 | 9 to 10 | 2 1/2 tsp or 12.5 mL |
| 72 to 95 | 11 | 3 tsp or 15mL |

*or as directed by a doctor

Other Information
store at room temperature 20°-25°C (68-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxy methyl cellulose, potassium sorbate, purified water, sorbitol, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com

N 3 71399 00388 1

children acetaminophen suspension

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:71399-0038 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|-----------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 160 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| SUCROSE (UNII: C151H8M554) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |

Product Characteristics

| | | | |
|-----------------|--------|---------------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | CHERRY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:71399-0038-4 | 1 in 1 CARTON | 03/18/2024 | |
| 1 | | 118 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:71399-0038-8 | 1 in 1 CARTON | 04/24/2024 | |
| 2 | | 236 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 3 | NDC:71399-0038-6 | 473 mL in 1 BOTTLE; Type 0: Not a Combination Product | 04/24/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug | M013 | 03/18/2024 | |

Labeler - Akron Pharma Inc. (067878881)

Revised: 4/2024

Akron Pharma Inc.