

MAX RELIEF JUNIOR- children acetaminophen suspension
Akron Pharma Inc.

MAX relief junior Cherry flavor
Children's Pain Reliever
Acetaminophen Oral Suspension
160mg per 5mL

Dye-Free

Active Ingredient

Active ingredient (in each 5mL)	Purpose
Acetaminophen 160 mg	Pain reliever/fever reducer

Uses

temporarily: reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:

if sore throat is severe, persists or more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor or pharmacist before use if the child has

liver disease.

Ask doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin. **When using this product do not exceed recommended dose (see overdose warning)**

Stop use and ask a doctor if

- pain gets worse or lasts for more than 5 days
- fever gets worse or last for more than 3 days
- new symptoms occur redness or swelling is present. These could be signs of a serious condition.

Overdose warning:

In case of overdose, get medical help to contact Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Keep out of Reach of children

Directions

- this product does not contain directions or complete warnings for adult use
- shake well before using
- mL = milliliter
- find right dose on chart below. if possible, use weight to dose, otherwise use age
- use only the enclosed dosing cup designed for use with product.

Weight (lbs)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
25 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	2 1/2 tsp or 12.5 mL

Other Information

- Store at room temperature 20°-25°C (68°-77°F). Protect from freezing. Protect from light.

Inactive Ingredients

Anhydrous Citric Acid, Cherry Flavor, Glycerin, Microcrystalline cellulose and Sodium Methyl Cellulose, Potassium Sorbate, Purified Water, Sorbitol, Sucralose, Sucrose, Xanthan gum.

Questions or comments?

call toll-free 1-877-225-6999

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

NDC 71399-0038-4



MAX reliefTM junior
Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen
Oral Suspension
160mg per 5mL

PAIN + FEVER

Cherry
Flavor

Akron Pharma

4 FL.OZ (118 mL)

Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days, ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ do not give more than 5 days unless directed by doctor. ■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.



52/90794
3 71399-00384 3

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com

160mg per 5mL
Acetaminophen
and fever Reducer
Children's
PAIN RELIEVER
 Dye-Free
MAXrelief[™] junior

NDC 71399-0038-4

MAXrelief[™] junior
 Dye-Free
Children's
PAIN RELIEVER
 and fever Reducer
Acetaminophen
 Oral Suspension
160mg per 5mL

PAIN + FEVER



Cherry flavor
Akron Pharma 4 FL.OZ (118 mL)

Drug Facts

Active ingredient (in each 5 mL) Purpose
 Acetaminophen 160 mg Pain reliever / fever reducer

Uses temporarily: reduces fever, relieves minor aches and pains due to: the common cold, flu, headache, sore throat, toothache

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
 more than 5 doses in 24 hours, which is the maximum daily amount
 with other drugs containing acetaminophen.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist, or if your child is allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
When using this product, do not exceed recommended dose (see overdose warning).
Stop use and ask a doctor if pain gets worse or lasts for more than 5 days, fever gets worse or lasts for more than 3 days, new symptoms occur, or redness or swelling is present. These could be signs of a serious condition.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-232-1223). Quick medical attention is critical even if you do not notice any signs or symptoms.
Directions
 This product does not contain directions or complete warnings for adult use. Shake well before using.
 Find right dose on chart below. If possible, use weight to dose; otherwise, use age.
 Use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.
 If needed, repeat dose every 4 hours while symptoms last.

NDC 71399-0038-4

MAXrelief[™] junior
 Dye-Free
Children's
PAIN RELIEVER
 and fever Reducer
Acetaminophen
 Oral Suspension
160mg per 5mL

PAIN + FEVER



Cherry flavor
Akron Pharma 4 FL.OZ (118 mL)

Drug Facts (continued)

- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor
- do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information
 Store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments?
 call toll-free 1-877-225-8999.

**This product is not manufactured or distributed by McKesson Consumer Healthcare, distributor of Infant Tylenol Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Akron Pharma
 Manufactured for:
 Akron Pharma, Inc.
 Fairfield, NJ 07004
 www.akronpharma.com

811399003843

NDC 71399-0038-8

MAXrelief™
Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen
Oral Suspension
160mg per 5mL

PAIN + FEVER

Cherry
flavor



Akron Pharma

8 FL.OZ (236 mL)

Drug Facts

Active ingredient (in each 5 mL) **Purpose**

Acetaminophen 160 mgPain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device. ■ if needed, repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours

Drug Facts (continued)

■ do not give more than 5 days unless directed by doctor.
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Akron Pharma

Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004

www.akronpharma.com



160mg per 5mL
Oral Suspension

Acetaminophen
and fever Reducer
PAIN RELIEVER

Children's
Dye-Free
MAXrelief™ junior

NDC 71399-0038-8

MAXrelief™ junior
Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen
Oral Suspension

160mg per 5mL

PAIN + FEVER



Akron Pharma

8 FL.OZ (236 mL)

Drug Facts

Active ingredient (in each 5 mL) Purpose
Acetaminophen 160 mg _____ Pain reliever / fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
■ more than 5 doses in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ hives ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days.
■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-422-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

■ this product does not contain directions or complete warnings for adult use, shake well before using

■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ use only the enclosed dosing cup designed for use with product. Do not use any other dosing device.

■ if needed, repeat dose every 4 hours while symptoms last
■ do not give more than 5 times in 24 hours

NDC 71399-0038-8

MAXrelief™ junior
Dye-Free
Children's
PAIN RELIEVER
and fever Reducer
Acetaminophen
Oral Suspension

160mg per 5mL

PAIN + FEVER



Akron Pharma

8 FL.OZ (236 mL)

Drug Facts (continued)

■ do not give more than 5 days unless directed by doctor
■ do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethyl cellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments?

call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infant Tylenol Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com



Drug Facts	
Active ingredient (in each 5 mL)	Purpose
Acetaminophen 160 mg	Pain reliever / fever reducer
Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache	
Warning	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
Do not use	
■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.	
Ask a doctor before use if your child has liver disease.	
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.	
When using this product, do not exceed recommended dose (see overdose warning)	
Stop use and ask a doctor if	
■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.	
Keep out of reach of children.	
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.	
Directions	
■ this product does not contain directions or complete warnings for adult use, shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; other wise, use age. ■ if needed, repeat dose every 4 hours while symptoms last	

NDC 71399-0038-6

MAXreliefTM
junior

Dye-Free

Children's

PAIN RELIEVER
and fever Reducer

PAIN+FEVER Acetaminophen
Oral Suspension

160mg per 5mL



Akron Pharma

16 FL.OZ (473 mL)

Drug Facts (continued)

- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor.
- do not give more than directed (see overdose warning)

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

*or as directed by a doctor

Other Information

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

Inactive Ingredients anhydrous citric acid, cherry flavor, glycerin, microcrystalline cellulose and sodium carboxymethylcellulose, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? call toll-free 1-877-225-6999.

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Infants' Tylenol® Oral Suspension.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

Akron Pharma

Manufactured for:

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

Rev.06/25



MAX RELIEF JUNIOR

children acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71399-0038
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3Z D4)	

SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-0038-4	1 in 1 CARTON	03/18/2024	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:71399-0038-8	1 in 1 CARTON	04/24/2024	
2		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:71399-0038-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/24/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/18/2024	

Labeler - Akron Pharma Inc. (067878881)

Revised: 10/2025

Akron Pharma Inc.