

**POLYETHYLENE GLYCOL 3350, NF POWDER FOR SOLUTION, LAXATIVE-
polyethylene glycol 3350, nf powder for solution, laxative powder, for
solution**

LGM PHARMA SOLUTIONS, LLC

PEG 3350

Drug Facts

Active Ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Manufactured by:

LGM Pharma Solutions, LLC

Irvine, CA 92614

7116-0119-99-EC-PD

Rev. 01/2022

Ask a doctor before use if you have:

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if:

- you have rectal bleeding, or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to the top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other Information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-877-288-1495

Polyethylene Glycol 3350, NF increases frequency of bowel movements and

softens the stool.

Principal Display Panel

LGM PHARMA SOLUTIONS, LLC

PRODUCT: Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

Net Wt. 4.1 OZ (119g)

7 ONCE-DAILY DOSES

NDC: 79739-7116-1

Dissolves in Any Beverage

LGM Pharma
PARTNER SMART.

NDC 79739-7116-1

Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

Net Wt. 4.1 OZ (119 g)
7 ONCE-DAILY DOSES

Drug Facts Lift Here

Active Ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line)	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Manufactured by
LGM Pharma Solutions, LLC
Irvine, CA 92614

7116-0119-99-EC-PD
Rev 01/2022


N 3 79739-7116-1 8

Drug Facts (continued)

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 7 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- **do not take more than directed unless advised by your doctor**
- The single bag is a resealable cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap).
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
- fill and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficult swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under, ask a doctor

Drug Facts (continued)

Other information

- store at 20° - 25°C (68° - 77°F)
- longer-evoked, do not use if hot seal under cap, provided with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments? 1-877-288-1498

Polyethylene Glycol 3350, NF increases frequency of bowel movements and softens the stool.

Drug Facts Lift Here

Active Ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line)	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days


Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Manufactured by
LGM Pharma Solutions, LLC
Irvine, CA 92614

7116-0119-99-EC-PD
Rev 01/2022


3 79739-7116-1 8

2.3750"

Principal Display Panel

LGM PHARMA SOLUTIONS, LLC

PRODUCT: Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

NDC: 79739-7116-2

Net Wt. 8.3 OZ (238g)

14 ONCE-DAILY DOSES

Dissolves in Any Beverage

NDC 79739-7116-2

Polyethylene Glycol 3350, NF increases frequency of bowel movements and softens the stool.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- children 16 years of age or older: ask a doctor

Other information

- store at 15° - 25°C (59° - 77°F)
- tamper-evident: do not use if top seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients: none

Questions or comments? 1-877-288-1488

Manufactured by:
LGM Pharma Solutions, LLC 7116-0238-99-BL-PD
Irvine, CA 92614 Rev 01/2022

LGM Pharma
PARTNER PHARMA

Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

Net Wt. 8.3 OZ (238 g)
14 ONCE-DAILY DOSES

Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line)	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

79739-7116-2

Principal Display Panel

LGM PHARMA SOLUTIONS, LLC

PRODUCT: Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relives Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

NDC: 79739-7116-3

Net Wt. 17.9 OZ (510g)

30 ONCE-DAILY DOSES

Dissolves in Any Beverage

NDC 79739-7116-3

Polyethylene Glycol 3350, NF
increases frequency of bowel
movements and softens the stool.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section of cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil seal under cap, printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken

Inactive ingredients none

Questions or comments? 1-877-258-1486



Polyethylene Glycol 3350, NF Powder for Solution, Osmotic Laxative

- Relieves Occasional Constipation/Irregularity
- Softens Stool
- Unflavored Powder
- Sugar Free

Net Wt. 17.9 OZ (510 g)
30 ONCE-DAILY DOSES

Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line)	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have:

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- intable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if:

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.



POLYETHYLENE GLYCOL 3350, NF POWDER FOR SOLUTION, LAXATIVE

polyethylene glycol 3350, nf powder for solution, laxative powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79739-7116
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 1 d

Product Characteristics

Color	white (colorless upon dissolution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79739-7116-1	7 d in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
2	NDC:79739-7116-2	14 d in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
3	NDC:79739-7116-3	30 d in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090812	10/07/2009	

Labeler - LGM PHARMA SOLUTIONS, LLC (117549198)

Establishment

Name	Address	ID/FEI	Business Operations
LGM Pharma Solutions, LLC		117549200	analysis(79739-7116) , label(79739-7116) , manufacture(79739-7116) , pack(79739-7116)

Revised: 1/2024

LGM PHARMA SOLUTIONS, LLC