

**WAL TUSSIN DM MAX NIGHTTIME WAL TUSSIN DM MAX- dextromethorphan hbr,
doxylamine succinate, guaifenesin
Walgreen Company**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Walgreen Co. Wal-Tussin® DM Max Nighttime Wal-Tussin® DM Max Drug Facts

Active ingredients (in each 20mL) - NIGHTTIME

Dextromethorphan HBr, USP 30 mg

Doxylamine succinate, USP 12.5 mg

Purposes

Cough suppressant

Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do Not Use

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 6 hours
children under 12 years	do not use

Other information

- **each 20 mL contains:** sodium 11 mg
- store at 20-25°C (68-77°F)

Inactive Ingredients

anhydrous citric acid, benzoic acid, benzyl alcohol, carboxymethylcellulose sodium, FD&C blue #1, FD&C red #40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Active ingredients (in each 20 mL) - DAYTIME

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 400 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
-----	------

adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Other information

- **each 20 mL contains:** sodium 13 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive Ingredients

acetic acid, anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

DAY & NIGHT PACK

Compare to Robitussin® Maximum Strength Cough + Chest Congestion DM & Robitussin® Maximum Strength Nighttime Cough DM active ingredients

ADULT

NON-DROWSY

Wal-Tussin®

DM Max

COUGH & CHEST CONGESTION DM

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

GUAIFENESIN / EXPECTORANT

MAXIMUM STRENGTH

ALCOHOL FREE

Relieves cough, chest congestion & mucus

For maximum strength mucus relief

12 years & older

SEE NEW DOSING

RASPBERRY MENTHOL FLAVOR

ADULT

Nighttime Wal-Tussin® DM Max

COUGH

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

MAXIMUM STRENGTH

ALCOHOL FREE

Relieves cough, itchy throat & runny nose

12 years & older

SEE NEW DOSING

RASPBERRY, BLACKBERRY & MENTHOL FLAVOR

TOTAL 8 FL OZ (236 mL) – 2 x 4 FL OZ (118 mL)

DAY & NIGHT PACK

NDC 0363-0913-12

Walgreens

ADULT
NON-DROWSY

Wal-Tussin®
DM Max

COUGH & CHEST
CONGESTION DM

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT

Compare to Robitussin® Maximum
Strength Cough + Chest Congestion
DM & Robitussin® Maximum Strength
Nighttime Cough DM active ingredients†

ADULT

Nighttime
Wal-Tussin®
DM Max

COUGH

DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
DOXYLAMINE SUCCINATE /
ANTIHISTAMINE

MAXIMUM STRENGTH

ALCOHOL FREE

- Relieves cough, chest congestion & mucus
- For maximum strength mucus relief
- 12 years & older



RASPBERRY
MENTHOL
FLAVOR

SEE NEW DOSING

MAXIMUM STRENGTH

ALCOHOL FREE

- Relieves cough, itchy throat & runny nose
- 12 years & older



RASPBERRY, BLACKBERRY
& MENTHOL FLAVOR

SEE NEW DOSING

TOTAL 8 FL OZ (236 mL) – 2 x 4 FL OZ (118 mL)



Drug Facts (continued)

Other information

- each 20 mL contains: sodium 13 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients acetic acid, anhydrous citric acid, carboxymethylcellulose sodium, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

PHARMACIST

Walgreens
Trusted since 1901™

Health expertise
you rely on.™

RECOMMENDED

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

ORG0518-F



ADULT
Nighttime
Wal-Tussin[®]
DM Max
COUGH

ADULT
NON-DROWSY
Wal-Tussin[®]
DM Max
COUGH & CHEST
CONGESTION DM

Walgreens

Wal-Tussin[®] DM Max

Nighttime Wal-Tussin[®] DM Max

DO NOT TAKE BOTH PRODUCTS WITHIN 6 HOURS OF EACH OTHER

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg.....Cough suppressant
Guaifenesin, USP 400 mg.....Expectorant

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 4 hours
children under 12 years	do not use

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 30 mg.....Cough suppressant
Doxylamine succinate, USP 12.5 mg.....Antihistamine

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- controls the impulse to cough to help you sleep

Warnings

Do not use

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Drug Facts (continued)

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- do not take more than 4 doses in any 24-hour period
- this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	20 mL every 6 hours
children under 12 years	do not use

Other information

- each 20 mL contains: sodium 11 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, benzoic acid, benzyl alcohol, carboxymethylcellulose sodium, FD&C blue #1, FD&C red #40, flavor, glycerin, menthol, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-800-719-9260

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

Gluten Free

¹Walgreens Pharmacist Survey
¹¹These products are not manufactured or distributed by Pfizer, distributor of Robitussin[®] Maximum Strength Cough + Chest Congestion DM and Robitussin[®] Maximum Strength Nighttime Cough DM.

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015

Walgreens
**100% SATISFACTION
GUARANTEED**

walgreens.com ©2018 Walgreen Co.

ITEM 782141 W10106-0818-L



9Z212 94 C1

WAL TUSSIN DM MAX NIGHTTIME WAL TUSSIN DM MAX

dextromethorphan hbr, doxylamine succinate, guaifenesin kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0913
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0913-12	1 in 1 CARTON; Type 0: Not a Combination Product	10/27/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	1 BOTTLE	118 mL

Part 1 of 2**NIGHTTIME WAL TUSSIN DM MAX**

dextromethorphan hbr, doxylamine succinate solution

Product Information

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 20 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BENZOIC ACID (UNII: 8SKN0B0MIM)	

BENZYL ALCOHOL (UNII: LKG8494WBH)
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)
FD&C RED NO. 40 (UNII: WZB9127XOA)
GLYCERIN (UNII: PDC6A3C0OX)
MENTHOL (UNII: L7T10EP3A)
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
WATER (UNII: 059QF0KO0R)
SODIUM BENZOATE (UNII: OJ245FE5EU)
SORBITOL (UNII: 506T60A25R)
SUCRALOSE (UNII: 96K6UQ3ZD4)
XANTHAN GUM (UNII: TTV12P4NEE)

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

WAL TUSSIN DM MAX
dextromethorphan hbr, guaifenesin solution

Product Information

Item Code (Source)	NDC:0363-1023
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL

GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)

GUAIFENESIN

400 mg
in 20 mL**Inactive Ingredients**

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	RED	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1023-26	118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/09/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/27/2018	

Labeler - Walgreen Company (008965063)