

**VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU- acetaminophen,  
dextromethorphan hydrobromide, and doxylamine succinate liquid  
The Procter & Gamble Manufacturing Company**

-----  
**VICKS® NyQuil™ HIGH BLOOD PRESSURE COLD & FLU**

***Drug Facts***

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:**

This product contains acetaminophen.

Severe liver damage may occur if you take

- more than 4 doses (30 mL each dose) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:**

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL per dose every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

---

## Other information

- **each 30 mL contains:** sodium 28.23 mg
- store at no greater than 25°C and do not refrigerate

## Inactive ingredients

citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

## Questions?

**1-800-362-1683**

**TAMPER EVIDENT:** Do not use if printed shrinkband seal around the neck is broken or missing

**DIST. BY:**

**PROCTER & GAMBLE**

**CINCINNATI, OH 45202**

**PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label**

VICKS®

NyQuil™

HIGH BLOOD PRESSURE

## COLD & FLU

**Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr**  
Headache, Fever, Sore Throat, Minor Aches & Pains  
Sneezing, Runny Nose

Cough

SUGAR & ALCOHOL FREE

*Nighttime Relief*

DECONGESTANT FREE

8 FL OZ (236 mL)



**Drug Facts (continued)**  
**Directions**  
• take only as directed  
• only use the dose cup provided  
• do not exceed 4 doses per 24 hrs  
adults & children 12 yrs & over 30 mL per dose every 9 hrs  
children 4 to under 12 yrs ask a doctor  
children under 4 yrs do not use  
**Other information**  
• each 30 mL contains: sodium 28.23 mg  
• store at no greater than 25°C and do not refrigerate  
**Inactive ingredients:** citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum  
**Questions?** 1-800-362-1083

**Drug Facts (continued)**  
• a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema  
• trouble urinating due to enlarged prostate gland  
**Ask a doctor or pharmacist before use if you are**  
• taking sedatives or tranquilizers  
• taking the blood thinning drug warfarin  
**When using this product**  
• excitability may occur, especially in children  
• marked drowsiness may occur  
• avoid alcoholic drinks  
• be careful when driving a motor vehicle or operating machinery  
• alcohol, sedatives, and tranquilizers may increase drowsiness  
**Stop use and ask a doctor if**  
• pain or cough gets worse or lasts more than 7 days  
• fever gets worse or lasts more than 3 days  
• redness or swelling is present  
• new symptoms occur  
• cough comes back or occurs with rash or headache that lasts  
These could be signs of a serious condition.  
**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.**

**Drug Facts (continued)**  
**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
• more than 4 doses (30 mL each dose) in 24 hours, which is the maximum daily amount for this product  
• with other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product  
**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
• skin redness • hives • rash  
If a skin reaction occurs, stop use and seek medical help right away  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use**  
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
• If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have**  
• liver disease • glaucoma  
• cough that occurs with too much phlegm (mucus)



## VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69423-973
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL

<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
---	----------------------	---------------------

### Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-973-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/12/2022	
2	NDC:69423-973-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/12/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/12/2022	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)