

DR. SCHOLLS BLISTER PAIN RELIEF COMPLETE- lidocaine
Scholl's Wellness Company LLC

Dr. Scholl's Blister Pain Relief Complete Kit

Drug Facts

Active ingredient

Lidocaine 4%

Purpose

External Analgesic

Use

Temporarily relieves pain

Warnings

For external use only

- Avoid contact with eyes
- If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

Do not use

- More than one patch on your body at a time
- If you have had an allergic reaction to lidocaine or other local anesthetics
- Do not use in large quantities, particularly over raw surfaces or blistered areas
- For more than one week without consulting a doctor.

When using this product

- Use only as directed. Read and follow all directions and warnings on this carton

Stop use and ask a doctor if

- condition worsens
- severe burning sensation, redness, rash or irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.

If pregnant or breast-feeding,

ask a health care professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children over 12 years:

Step 1 (Lidocaine patch application) • apply to the affected area not more than 3-4 times daily

- clean and dry affected area
- remove backing from lidocaine patch by firmly grasping both ends and gently pulling away backing
- once exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area
- once applied, press firmly on patch to help adhesive adhere to blister
- do not remove patch and reapply
- once patch is applied, move to Step 2.

Step 2 (Bandage application) • remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over skin to seal lidocaine patch and blister.

- remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when bandage begins to loosen

children 12 years or younger: ask a doctor

Other information

Store between 20-25°C (68-77°F)

Inactive ingredients

Cellulose Gum, Dihydroxyaluminum Aminoacetate, Ethylhexylglycerin, Glycerol, Laurocapram, PEG-40 Hydrogenated Castor Oil, Phenoxyethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopherol Acetate, Vanillyl Butyl Ether, Urea, Water (Aqua)

Questions?

1-866-360-3226

Package Labeling:

BL-CM1-20096-1-24.10.9

Drug Facts

Active ingredient

Lidocaine 4%

Purpose

External Analgesic

Use Temporarily relieves pain

Warnings

For external use only

■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

Do not use

■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.

When using this product

■ Use only as directed. Read and follow all directions and warnings on this carton

Stop use and ask a doctor if ■ condition worsens ■ severe burning sensation, redness, rash or irritation develops ■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.

If pregnant or breast feeding, ask a health care professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions Adults and children over 12 years:

Step 1 (Lidocaine patch application) ■ apply to the affected area not more than 3-4 times daily ■ clean and dry affected area ■ remove backing from lidocaine patch by firmly grasping both ends and gently pulling away backing ■ once exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area ■ once applied, press firmly on patch to help adhesive adhere to blister ■ do not remove patch and reapply ■ once patch is applied, move to Step 2.

Step 2 (Bandage application) ■ remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over skin to seal lidocaine patch and blister. ■ remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when bandage begins to loosen

Children 12 years or younger: ask a doctor

Other information Store between 20-25°C (68-77°F)

Inactive ingredients Cellulose Gum, Dihydroxyaluminum Aminoacetate, Ethylhexylglycerin, Glycerol, Laureocapram, PEG-40 Hydrogenated Castor Oil, Phenoxylethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopherol Acetate, Vanillyl Butyl Ether, Zinc, Water (Aqua/Eau)

Questions? 1-866-360-3228

FSA/HSA ELIGIBLE

For additional information and instructions visit: drscholls.com
Pat: patents.drscholls.com

INSTRUCTIONS FOR USE:

Lidocaine Patch Application
Clean and dry affected area. Remove backing from lidocaine patch and apply to blister. Press firmly on patch to help adhesive adhere to blister. Move to hydrogel bandage application.



Hydrogel Bandage Application

Step 1: Remove backing film from hydrogel bandage marked "Step 1" and place center (adhesive side down) over blister.



Step 2: Remove backing film marked "Step 2" and firmly smooth down adhesive edge to skin to seal lidocaine patch and blister. Replace when strip begins to loosen.



NEW

Dr.Scholl's

Dr.Scholl's

BLISTER PAIN RELIEF COMPLETE KIT

SEAL & HEAL BANDAGES

LIDOCAINE 4%

CLINICALLY TESTED 12-HOUR PAIN RELIEF

- ✓ Lidocaine patch numbs pain
- ✓ Hydrogel bandage seals & cushions blister



12 LIDOCAINE PATCHES

12 HYDROGEL BANDAGES

BLISTER PAIN RELIEF COMPLETE KIT

SEAL & HEAL BANDAGES

- ✓ Maximum strength lidocaine without a prescription
- ✓ Only blister treatment with an active pain reliever

©/© Scholl's Wellness Company 2024
Dist. by: Scholl's Wellness Company LLC, Parsippany, NJ 07054
Made in China.

C1187



彩面

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Drug Facts (continued)
burning sensation, redness, rash or irritation develops
■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied.
■ **pregnant or breast feeding**, ask a health care professional before use. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.
Directions Adults and children over 12 years:
■ apply to the affected area not more than 3-4 times daily. ■ **clean** and dry affected area ■ **remove** backing from lidocaine patch by firmly grasping both ends and gently pulling away backing. ■ **once** exposed portion of patch is positioned, carefully remove remaining backing completely to apply patch to affected area. ■ **once** applied, press firmly on patch to help adhesive adhere to skin. ■ **do not** remove patch and reapply. ■ **once** patch is applied, move to Step 2. **Children 12 years or younger**; ask a doctor.
Other information Store between 20-25°C (68-77°F)
Inactive ingredients Cellulose Gum, Dihydroxyaluminum Ammonioacetate, Ethylhexylglycerin, Glycolol, Lauracopran, PEG-40 Hydrogenated Castor Oil, Phenoxymethanol, Polyacrylic Acid, Propylene Glycol, Silica Silylate, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Tocopheryl Acetate, Vanillin Butyl Ether, Urea, Water (Aqua/Eau)
Questions? 1-866-360-3226

Div. by Scholl's Wellness Company LLC
Paterson, NJ 07654
Made in China
CIB-remnant packaging
LOT: 20241115
EXP: 2026-11-14

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶

Dr. Scholl's
4% LIDOCAINE PATCH
External Analgesic

1 PATCH
1.9" L x 0.813" W (3 cm x 2 cm) Weight: 0.37g

Drug Facts	Active ingredient	Purpose
Lidocaine 4%	External Analgesic	

Use Temporarily relieves pain
Warnings
For external use only
■ Avoid contact with eyes ■ If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.
Do not use
■ More than one patch on your body at a time ■ If you have had an allergic reaction to lidocaine or other local anesthetics ■ Do not use in large quantities, particularly over raw surfaces or blistered areas ■ For more than one week without consulting a doctor.
When using this product
■ Use only as directed, Read and follow all directions and warnings on this carton
Stop use and ask a doctor if ■ condition worsens ■ severe ▶



DR. SCHOLLS BLISTER PAIN RELIEF COMPLETE

lidocaine kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73469-0488
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73469-0488-2	1 in 1 KIT	02/15/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	12 PATCH	4.44 g

Part 1 of 1

DR SCHOLL S 4 LIDOCAINE EXTERNAL ANALGESIC

lidocaine patch

Product Information

Item Code (Source)	NDC:75568-024
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE GUM (UNII: K679OBS311)	
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
GLYCERIN (UNII: PDC6A3C0OX)	
LAUROCAPRAM (UNII: 1F3X9DRV9X)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TARTARIC ACID (UNII: W4888I119H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
VANILLYL BUTYL ETHER (UNII: S2ULN37C9R)	
UREA (UNII: 8W8T17847W)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75568-024-	1 in 1 PATCH		

1	01	1 in 1 PATCH		
1		0.37 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/15/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	02/15/2025	

Labeler - Scholl's Wellness Company LLC (117174744)

Establishment

Name	Address	ID/FEI	Business Operations
Shanghai Sabei Cosmetic Co., Ltd.		542969967	pack(75568-024)

Revised: 12/2024

Scholl's Wellness Company LLC