

OXYGEN- oxygen gas
Institutional Care Pharmacy

Oxygen

Oxygen Compressed USP Label

OXYGEN, COMPRESSED USP UN1072 OXYGEN 2

DANGER: COMPRESSED, OXIIZING GAS, MAY CAUSE OR INTENSIFY FIRE; OXIDIZER, CONTAINS GAS UNDER PRESSURE; MAY EXPLODE IF HEATED.

In case of fire: stop leak if safe to do so. Keep reduction valves free from grease and oil. Open valve slowly. Store and use in a well-ventilated place. Use a back-flow preventative device in the piping. Close valve after each use and when empty. Protect from sunlight when ambient temperature exceeds 52° C/125° F. Container temperature should not exceed 52° C/125° F. Do not handle until all safety precautions have been read and understood. Keep/store away from combustible materials. Use only with equipment cleaned for oxygen service and rated for cylinder pressure.

DO NOT REMOVE THIS PRODUCT LABEL.

For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications. **Rx only.**

Uninterrupted use of high concentrations of oxygen over a long duration without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment. Produced by Air Liquefaction.

Contents	Size
170 Liters	__ M6
248 Liters	__ C
415 Liters	__ D
682 Liters	__ E
6910 Liters	__ H
__ Liters	__ Other

CAS 7782-44-7

Institutional Care Pharmacy

1815 West County Road 54

Tiffin, OH 44883

1-800-228-8278

OXYGEN, COMPRESSED USP

UN 1072

DANGER:

COMPRESSED, OXIDIZING GAS.
MAY CAUSE OR INTENSIFY FIRE; OXIDIZER.
CONTAINS GAS UNDER PRESSURE;
MAY EXPLODE IF HEATED.

In case of fire: stop leak if safe to do so. Keep reduction valves free from grease and oil. Open valve slowly. Store and use in a well ventilated place. Use a back flow preventative device in the piping. Close valve after each use and when empty. Protect from sunlight when ambient temperature exceeds 52 °C/125 °F. Container temperature should not exceed 52 °C/125°F. Do not handle until all safety precautions have been read and understood. Keep/store away from combustible materials. Use only with equipment cleaned for oxygen service and rated for cylinder pressure.

DO NOT REMOVE THIS PRODUCT LABEL.

For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **Rx only**.

Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment. Produced by Air Liquefaction.



Contents	Size
170 Liters	<input type="checkbox"/> M6
248 Liters	<input type="checkbox"/> C
415 Liters	<input type="checkbox"/> D
682 Liters	<input type="checkbox"/> E
6910 Liters	<input type="checkbox"/> H
_____ Liters	<input type="checkbox"/> Other

CAS NO: 7782-44-7

Institutional Care Pharmacy

1815 West County Road 54

Tiffin, OH 44883

1-800-228-8278

Oxygen Label

OXYGEN, COMPRESSED USP

UN1072

OXYGEN 2

DANGER: COMPRESSED, OXIDIZING GAS, MAY CAUSE OR INTENSIFY FIRE; OXIDIZER, CONTAINS GAS UNDER PRESSURE; MAY EXPLODE IF HEATED.

In case of fire: stop leak if safe to do so. Keep reduction valves free from grease and oil. Open valve slowly. Store and use in a well-ventilated place. Use a back-flow preventative device in the piping. Close valve after each use and when empty. Protect from sunlight when ambient temperature exceeds 52° C/125° F. Container temperature should not exceed 52° C/125° F. Do not handle until all safety precautions have been read and understood. Keep/store away from combustible materials. Use only with equipment cleaned for oxygen service and rated for cylinder pressure.

DO NOT REMOVE THIS PRODUCT LABEL.

For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications. **Rx only.**

Uninterrupted use of high concentrations of oxygen over a long duration without monitoring its effect on oxygen content of arterial blood, may be harmful. Use only with pressure reducing equipment and apparatus designed for oxygen. Do not attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment. Produced by Air Liquefaction.

Contents	Size
170 Liters	__ M6
248 Liters	__ C
415 Liters	__ D
682 Liters	__ E
6910 Liters	__ H
__ Liters	__ Other

CAS 7782-44-7

Institutional Care Pharmacy

Southern Region

7537 Easy Street

Mason, OH 45040

513-573-9625

OXYGEN, COMPRESSED USP

UN 1072

DANGER:

MAY CAUSE OR INTENSIFY FIRE; OXIDIZER.
CONTAINS GAS UNDER PRESSURE;
MAY EXPLODE IF HEATED.



Do not handle until all safety precautions have been read and understood. Keep and store away from clothing and other combustible materials. Keep valves and fittings free from grease and oil. Use and store only outdoors or in a well-ventilated place. In case of fire: Stop leak if safe to do so. Use a back flow preventive device in the piping. Use only with equipment of compatible materials of construction and rated for cylinder pressure. Use only with equipment cleaned for oxygen service. Open valve slowly. Close valve after each use and when empty. Protect from sunlight when ambient temperature exceeds 52 °C (125 °F). Read and follow the Safety Data Sheet (SDS) before use.

CAS NO: 7782-44-7

DO NOT REMOVE THIS PRODUCT LABEL.

WARNING: For emergency use only when administered by properly trained personnel for oxygen deficiency and resuscitation. For all other medical applications, **Rx only**. Uninterrupted use of high concentrations of oxygen over a long duration, without monitoring its effect on oxygen content of arterial blood, may be harmful. Do not attempt to use on patients who have stopped breathing unless used in conjunction with resuscitative equipment. Produced by Air Liquefaction. ©Applied Home Healthcare Equipment 2014

Contents	Size
113 Liters	<input type="checkbox"/> M4
165 Liters	<input type="checkbox"/> M6
255 Liters	<input type="checkbox"/> C
425 Liters	<input type="checkbox"/> D
680 Liters	<input type="checkbox"/> E
7080 Liters	<input type="checkbox"/> H
____ Liters	<input type="checkbox"/> Other

Institutional Care Pharmacy Southern Region

7537 Easy Street
Mason, OH 45040
513-573-9625

OXYGEN

oxygen gas

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:44363-1815
Route of Administration	RESPIRATORY (INHALATION)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYGEN (UNII: S88TT14065) (OXYGEN - UNII:S88TT14065)	OXYGEN	99 L in 100 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:44363-1815-1	170 L in 1 CYLINDER; Type 0: Not a Combination Product	01/01/1988	
2	NDC:44363-1815-2	248 L in 1 CYLINDER; Type 0: Not a Combination Product	01/01/1988	
3	NDC:44363-1815-3	415 L in 1 CYLINDER; Type 0: Not a Combination Product	01/01/1988	
4	NDC:44363-1815-4	682 L in 1 CYLINDER; Type 0: Not a Combination Product	01/01/1988	
5	NDC:44363-1815-5	6910 L in 1 CYLINDER; Type 0: Not a Combination Product	01/01/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA205767	01/01/1988	

Labeler - Institutional Care Pharmacy (068899806)

Registrant - Institutional Care Pharmacy (068899806)

Establishment

Name	Address	ID/FEI	Business Operations
Institutional Care Pharmacy		068899806	manufacture(44363-1815)

Establishment

Name	Address	ID/FEI	Business Operations
Institutional Care Pharmacy, Inc		078368225	manufacture(44363-1815)

