ASPIRIN- aspirin tablet Athlete's Needs Inc

Clear Choice 44-249

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• do not take more than directed

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredient

corn starch

Questions or comments?

1-800-426-9391

Principal Display Panel

Clear Choice

ASPIRIN

- Pain Reliever
- Fever Reducer (NSAID)


PAIN & FEVER RELIEF CAFFEINE FREE SODIUM FREE

Actual Size

100 Tablets 325 mg Each

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Distributed by: A.N.I. Toledo, OH 43623 50844 REV0122C24912



| ASPIRIN | | | | | |
|--|----------------|--------------------|--------------------------|---------------|----------|
| aspirin tablet | | | | | |
| | | | | | |
| Product Information | | | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | | NDC:53185-249 | |
| Route of Administration | ORAL | | | | |
| | | | | | |
| | - | | | | |
| Active Ingredient/Active | Moiety | | | | |
| Ingredient Name | | | Basis of Strength | | Strength |
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | | | ASPIRIN | | 325 mg |
| | | | | | |
| In a stirre I norme dia mta | | | | | |
| Inactive Ingredients | | | | | |
| li li | ngredient Name | | | Stre | ength |

| STARCH, CORN (UNII | l: 08232NY3SJ) | | | | | |
|---|---|----------------------|-------------------|----------------|-----------------------|--|
| | | | | | | |
| Product Charac | teristics | | | | | |
| Color | white | Score | no | no score | | |
| Shape | ROUND | Size | | 10mm | | |
| Flavor | | Imprint Code | | ASPIRIN;44;249 | | |
| Contains | | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| Packaging | | | | | | |
| | Package | Description | Marketing Date | Start | Marketing End Date | |
| # Item Code 1 NDC:53185-249- 1 | - | 0: Not a Combination | | Start | - | |
| # Item Code 1 NDC:53185-249- 1 | .00 in 1 BOTTLE; Type | 0: Not a Combination | Date | Start | - | |
| # Item Code 1 NDC:53185-249- 1 | .00 in 1 BOTTLE; Type | 0: Not a Combination | Date | Start | - | |
| # Item Code 1 NDC:53185-249- 1 | .00 in 1 BOTTLE; Type Product | 0: Not a Combination | Date | Start | - | |
| # Item Code 1 NDC:53185-249- 12 1 P | 00 in 1 BOTTLE; Type Product nformation Application Nu | 0: Not a Combination | Date | g Start | - | |

Labeler - Athlete's Needs Inc (619703945)

| Establishment | | | |
|-------------------------|---------|-----------|----------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 038154464 | pack(53185-249) |
| | | | |
| Establishment | | | |
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 832867837 | manufacture(53185-249) |
| | | | |
| Establishment | | | |
| | | | |

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|----------------------------|
| LNK International, Inc. | | 967626305 | pack(53185-249) |

Revised: 10/2023

Athlete's Needs Inc