# HEALTH MART PHARMACY CHILDRENS LORATADINE- loratadine solution Health Mart

\_\_\_\_\_

### $\textbf{Health Mart}_{\texttt{R}} \textbf{ Pharmacy Children's Loratadine}$

Drug Facts

# Active ingredient (in each 5 mL teaspoonful)

Loratadine 5 mg

# Purpose

Antihistamine

# Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

# Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product** do not take more than directed. Taking more than directed may cause drowsiness.

#### Stop use and ask a doctor if

• an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **(1-800-222-1222)** 

# Directions

• use only with enclosed dosing cup

adults and children 6	2 teaspoonfuls (TSP) daily; do not take more than 2
years and over	teaspoonfuls (TSP) in 24 hours
children 2 to under 6	1 teaspoonful (TSP) daily; do not take more than 1
years of age	teaspoonful (TSP) in 24 hours
children under 2 years	ask a doctor
of age	
consumers with liver	ask a doctor
or kidney disease	dSK d UUCIUI

#### Other information

- safety sealed: do not use if imprinted safety seal is torn or missing
- store between 20° and 25°C (68° and 77°F)

#### **Inactive ingredients**

artificial grape flavor, citric acid monohydrate, glycerin, propylene glycol, purified water, sodium benzoate, sodium metabisulfite, sucrose

#### Questions?

#### Call 1-866-923-4914

Distributed by McKesson One Post Street San Francisco, CA 94104

# PRINCIPAL DISPLAY PANEL - 120 ml Bottle Carton

NDC 62011-0072-1

#### Health Mart<sub>®</sub> PHARMACY

compare to Children's Claritin<sup>®</sup> active ingredient\*

#### children's loratadine syrup

(loratadine oral solution) 5 mg/5 mL antihistamine

indoor & outdoor allergies original prescription strength

24 hour non-drowsy<sup>†</sup> relief of: sneezing; runny nose; itchy, watery eyes; itchy throat or nose

2 Yrs. & older


grape flavor dosing cup included alcohol free • gluten free

#### 4 FL OZ (120 ml)

<sup>†</sup>when taken as directed. see drug facts panel.

		Health Mart
		children' loratadine syrup
5		(loratadine oral solution 5 mg/5 mL antihistamin
SEALED WITH	I PRINTED NECKBAND	NDC 62011-00
Drug Facts Active ingredie (in each 5 mL		B Health Mar
Loratadine 5 mg	elieves these symptoms due to hay	compare Children's Clariti active ingredier
<ul> <li>sneezing</li> </ul>	itching of the nose or throat	children
Warnings Do not use if you have	ve ever had an allergic reaction to this	loratadine
	aredients	loiuluun
product or any of its in Ask a doctor before	ngredients. use if you have liver or kidney disease. termine if you need a different dose.	
product or any of its in Ask a doctor before Your doctor should det When using this pro	use if you have liver or kidney disease. termine if you need a different dose. duct do not take more than directed.	syrup
product or any of its in Ask a doctor before Your doctor should det When using this pro Taking more than direc Stop use and ask a product occurs. Seek	use if you have liver or kidney disease. termine if you need a different dose. duct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away.	(loratadine oral solution 5 mg/5 mL antihistamin
product or any of its in Ask a doctor before Your doctor should def When using this pro Taking more than direc Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or contact	use if you have liver or kidney disease. termine if you need a different dose. duct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this	(loratadine oral solution
product or any of its in Ask a doctor before of Your doctor should def When using this pro Taking more than direct Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or contact (1-800-222-1222)	use if you have liver or kidney disease. termine if you need a different dose. oduct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get ct a Poison Control Center right away.	Syrup (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng 24 hour non-drowsy†relief
product or any of its in Ask a doctor before of Your doctor should def When using this pro Taking more than direct Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or contact (1-800-222-1222)	use if you have liver or kidney disease. termine if you need a different dose. oduct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get a Poison Control Center right away. e only with enclosed dosing cup 2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls	Syrup (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng
product or any of its in Ask a doctor before Your doctor should det When using this pro Taking more than direct Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or contar (1-800-222-1222) Directions • use adults and children	use if you have liver or kidney disease. termine if you need a different dose. oduct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get a Poison Control Center right away. e only with enclosed dosing cup 2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls (TSP) in 24 hours 1 teaspoonful (TSP) daily; do not take more than 1 teaspoonful	Syrup (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng 24 hour non-drowsytrelief sneezing; runny nos itchy, watery eyes; itchy throat or no
product or any of its in Ask a doctor before of Your doctor should def When using this pro Taking more than direct Stop use and ask a product occurs. Seek If pregnant or breast before use. Keep out of reach of medical help or contact (1-800-222-1222) Directions • use adults and children 6 years and over children 2 to under 6 years of age	use if you have liver or kidney disease. termine if you need a different dose. induct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get ct a Poison Control Center right away. e only with enclosed dosing cup 2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls (TSP) in 24 hours 1 teaspoonful (TSP) daily; do not	Syrup (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng 24 hour non-drowsy trelief sneezing; runny nos itchy, watery eyes; itchy throat or no 2 Yrs. & old
product or any of its in Ask a doctor before of Your doctor should det When using this pro Taking more than dired Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or conta- (1-800-222-1222) Directions • use adults and children 6 years and over children 2 to under 6 years of age children under 2 years of age	use if you have liver or kidney disease. termine if you need a different dose. oduct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get ct a Poison Control Center right away. e only with enclosed dosing cup 2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls (TSP) in 24 hours 1 teaspoonful (TSP) daily; do not take more than 1 teaspoonful (TSP) in 24 hours	Syrug (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng 24 hour non-drowsytrelief sneezing; runny nos itchy, watery eyes; itchy throat or no 2 Yrs. & old
product or any of its in Ask a doctor before of Your doctor should det When using this pro Taking more than dired Stop use and ask a product occurs. Seek If pregnant or breas before use. Keep out of reach of medical help or conta- (1-800-222-1222) Directions • use adults and children 6 years and over children 2 to under 6 years of age	use if you have liver or kidney disease. termine if you need a different dose. oduct do not take more than directed. cted may cause drowsiness. doctor if • an allergic reaction to this medical help right away. st-feeding, ask a health professional of children. In case of overdose, get ct a Poison Control Center right away. e only with enclosed dosing cup 2 teaspoonfuls (TSP) daily; do not take more than 2 teaspoonfuls (TSP) in 24 hours 1 teaspoonful (TSP) daily; do not take more than 1 teaspoonful (TSP) in 24 hours ask a doctor	Syrup (loratadine oral solution 5 mg/5 mL antihistamin indoor & outdoor allergie original prescription streng 24 hour non-drowsytrelief sneezing; runny nos itchy, watery eyes; itchy throat or no





# HEALTH MART PHARMACY CHILDRENS LORATADINE

loratadine solution

Product Information					
Product T ype	HUMAN OTC DRUG	Item Code (Source)		NDC:62011-0072	
Route of Administration	ORAL				
Active Ingredient/Active Moie	ety				
Ingredient Name		Basis of Strength		ength	Strength
Loratadine (UNII: 7AJO3BO7QN) (Lor	atadine - UNII:7AJO3BO7QN)		Loratadine		5  mg  in  5  mL
Inactive Ingredients					
	Ingredient Name				Strength

citric acid monohyd	rate (UNII: 2968PHW8QP)				
glycerin (UNII: PDC6	A3C0OX)				
propylene glycol (U	NII: 6 DC9 Q 16 7 V3)				
water (UNII: 059QF0	KO0R)				
<b>sodium benzoate</b> (U	NII: OJ245FE5EU)				
sodium metabisulfit	e (UNII: 4VON5FNS3C)				
sucrose (UNII: C151H	3M554)				
Product Charac	teristics				
	YELLOW (colorless to slightly yellow)	Score			
Shape		Size			
Flavor	GRAPE Imprint C		ode		
Contains					
Packaging					
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>		
<b>1</b> NDC:62011-0072-1	1 in 1 CARTON	08/20/2004			
1	120 mL in 1 BOTTLE; Type 0: Not a Combination Product				
Marketing Information					
Marketing Catego	ry Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA076805	08/20/2004			

# Labeler - Health Mart (177667227)

# Establishment

Name	Address	ID/FEI	Business Operations
Taro Pharmaceutical Industries, Ltd.		600072078	MANUFACTURE(62011-0072)

Revised: 1/2019

Health Mart