


METHOCARBAMOL- methocarbamol tablet, coated

Bostal LLC

DESCRIPTION

Methocarbamol Tablets USP, 500 mg, 750 mg, and 1000 mg, a carbamate derivative of guaifenesin, is a central nervous system (CNS) depressant with sedative and musculoskeletal relaxant properties.

The chemical name of methocarbamol is a 3-(2 methoxyphenoxy)-1,2-propanediol 1-carbamate and has the empirical formula C₁₁H₁₅NO₅. Its molecular weight is 241.24. The structural formula is shown below.



Methocarbamol is a white powder, sparingly soluble in water and chloroform, soluble in alcohol (only with heating) and propylene glycol, and insoluble in benzene and n-hexane.

Methocarbamol Tablet, 500 mg is available as an orange, film coated, round concave tablet containing 500 mg of methocarbamol, USP for oral administration. The inactive ingredients present are microcrystalline cellulose, croscarmellose sodium, FD&C Yellow 6 aluminum lake, hydroxypropyl cellulose, hypromellose, magnesium stearate, polyethylene glycol, triacetin, titanium dioxide.

Methocarbamol Tablet, 750 mg is available as a yellow, film coated, modified capsule shaped tablet containing 750 mg of methocarbamol, USP for oral administration. The inactive ingredients present are microcrystalline cellulose, croscarmellose sodium, iron oxide yellow, iron oxide red, hydroxypropyl cellulose, hypromellose, magnesium stearate, polyethylene glycol, triacetin, titanium dioxide.

Methocarbamol tablet, 1000 mg is available as an orange, film coated, oblong-shaped tablet containing 1000 mg of methocarbamol, USP for oral administration. The inactive ingredients present in methocarbamol tablets 1000 mg are same as those present in methocarbamol tablets 500 mg.

Clinical Pharmacology

The mechanism of action of methocarbamol in humans has not been established, but may be due to general central nervous system (CNS) depression. It has no direct action on the contractile mechanism of striated muscle, the motor end plate or the nerve fiber.

Pharmacokinetics

In healthy volunteers, the plasma clearance of methocarbamol ranges between 0.20 and 0.80 L/h/kg, the mean plasma elimination half-life ranges between 1 and 2 hours, and the plasma protein binding ranges between 46% and 50%.

Methocarbamol is metabolized via dealkylation and hydroxylation. Conjugation of methocarbamol also is likely. Essentially all methocarbamol metabolites are eliminated in the urine. Small amounts of unchanged methocarbamol also are excreted in the urine.

Special populations

Elderly

The mean (\pm SD) elimination half-life of methocarbamol in elderly healthy volunteers (mean (\pm SD) age, 69 (\pm 4) years) was slightly prolonged compared to a younger (mean (\pm SD) age, 53.3 (\pm 8.8) years), healthy population (1.5 (\pm 0.4) hours versus 1.1 (\pm 0.27) hours, respectively). The fraction of bound methocarbamol was slightly decreased in the elderly versus younger volunteers (41 to 43% versus 46 to 50%, respectively).

Renally impaired

The clearance of methocarbamol in 8 renally-impaired patients on maintenance hemodialysis was reduced about 40% compared to 17 normal subjects, although the mean (\pm SD) elimination half-life in these two groups was similar: 1.2 (\pm 0.6) versus 1.1 (\pm 0.3) hours, respectively.

Hepatically impaired

In 8 patients with cirrhosis secondary to alcohol abuse, the mean total clearance of methocarbamol was reduced approximately 70% compared to that obtained in 8 age- and weight-matched normal subjects. The mean (\pm SD) elimination half-life in the cirrhotic patients and the normal subjects was 3.38 (\pm 1.62) hours and 1.11 (\pm 0.27) hours, respectively. The percent of methocarbamol bound to plasma proteins was decreased to approximately 40 to 45% compared to 46 to 50% in the normal subjects.

INDICATIONS AND USAGE

Methocarbamol is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of methocarbamol has not been clearly identified, but may be related to its sedative properties. Methocarbamol does not directly relax tense skeletal muscles in man.

CONTRAINDICATIONS

Methocarbamol is contraindicated in patients hypersensitive to methocarbamol or to any of the tablet components.

WARNINGS

Since methocarbamol may possess a general CNS depressant effect, patients receiving methocarbamol tablets should be cautioned about combined effects with alcohol and

other CNS depressants.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol tablets should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see PRECAUTIONS, Pregnancy).

Use In Activities Requiring Mental Alertness

Methocarbamol may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle. Patients should be cautioned about operating machinery, including automobiles, until they are reasonably certain that methocarbamol therapy does not adversely affect their ability to engage in such activities.

PRECAUTIONS

Information for Patients

Patients should be cautioned that methocarbamol may cause drowsiness or dizziness, which may impair their ability to operate motor vehicles or machinery.

Because methocarbamol may possess a general CNS-depressant effect, patients should be cautioned about combined effects with alcohol and other CNS depressants.

Drug Interactions

See WARNINGS and PRECAUTIONS for interaction with CNS drugs and alcohol.

Methocarbamol may inhibit the effect of pyridostigmine bromide. Therefore, methocarbamol should be used with caution in patients with myasthenia gravis receiving anticholinesterase agents.

Drug/Laboratory Test Interactions

Methocarbamol may cause a color interference in certain screening tests for 5-hydroxyindoleacetic acid (5-HIAA) using nitrosonaphthol reagent and in screening tests for urinary vanillylmandelic acid (VMA) using the Gitlow method.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies to evaluate the carcinogenic potential of methocarbamol have not been performed. No studies have been conducted to assess the effect of methocarbamol on mutagenesis or its potential to impair fertility.

Pregnancy

Teratogenic Effects — Pregnancy Category C

Animal reproduction studies have not been conducted with methocarbamol. It is also not known whether methocarbamol can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Methocarbamol should be given to a pregnant woman only if clearly needed.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see WARNINGS).

Nursing Mothers

Methocarbamol and/or its metabolites are excreted in the milk of dogs; however, it is not known whether methocarbamol or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when methocarbamol is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of methocarbamol in pediatric patients below the age of 16 have not been established.

ADVERSE REACTIONS

Adverse reactions reported coincident with the administration of methocarbamol include:

Body as a whole: Anaphylactic reaction, angioneurotic edema, fever, headache

Cardiovascular system: Bradycardia, flushing, hypotension, syncope, thrombophlebitis

Digestive system: Dyspepsia, jaundice (including cholestatic jaundice), nausea and vomiting

Hemic and lymphatic system: Leukopenia

Immune system: Hypersensitivity reactions

Nervous system: Amnesia, confusion, diplopia, dizziness or lightheadedness, drowsiness, insomnia, mild muscular incoordination, nystagmus, sedation, seizures (including grand mal), vertigo

Skin and special senses: Blurred vision, conjunctivitis, nasal congestion, metallic taste, pruritus, rash, urticaria

To report SUSPECTED ADVERSE REACTIONS, contact AustarPharma LLC at 1-844-375-5410 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

OVERDOSAGE

Limited information is available on the acute toxicity of methocarbamol. Overdose of methocarbamol is frequently in conjunction with alcohol or other CNS depressants and includes the following symptoms: nausea, drowsiness, blurred vision, hypotension, seizures, and coma.

In post-marketing experience, deaths have been reported with an overdose of methocarbamol alone or in the presence of other CNS depressants, alcohol or psychotropic drugs.

DOSAGE AND ADMINISTRATION

Methocarbamol, 500 mg — Adults:

Initial dosage: 3 tablets 4 times daily.

Maintenance dosage: 2 tablets 4 times daily

Methocarbamol, 750 mg — Adults:

Initial dosage: 2 tablets 4 times daily.

Maintenance dosage: 1 tablet 4 times daily or 2 tablets 3 times daily

Methocarbamol, 1000 mg — Adults:

Initial dosage: 1 ½ tablets 4 times daily.

Maintenance dosage: 1 tablets 4 times daily

Six grams a day are recommended for the first 48 to 72 hours of treatment. (For severe conditions 8 grams a day may be administered). Thereafter, the dosage can usually be reduced to approximately 4 grams a day.

Treatment

Management of overdose includes symptomatic and supportive treatment. Supportive measures include maintenance of an adequate airway, monitoring urinary output and vital signs, and administration of intravenous fluids if necessary. The usefulness of hemodialysis in managing overdose is unknown.

HOW SUPPLIED

Methocarbamol Tablets USP, 500 mg — Orange, film coated, round concave tablets with one side debossed 'AP212', the other side bisected. They are supplied as follows:

Bottles of 100, NDC 35561-212-12

Bottles of 500, NDC 35561-212-13

Methocarbamol Tablets USP, 750 mg — Yellow, film coated, modified capsule shaped tablets; one side debossed 'AP211' and other side blank. They are supplied as follows:

Bottles of 100, NDC 35561-211-1

Bottles of 500, NDC 35561-211-13

Methocarbamol Tablets USP, 1000 mg — Orange, film coated, oblong-shaped tablets;

one side debossed 'AP349' and other side bisected. They are supplied as follows:

Bottles of 100, NDC 35561-349-12

Bottles of 500, NDC 35561-349-13

Store at controlled room temperature, between 20° to 25°C (68° to 77°F).

Dispense in tight container.

LBL136

REV050622

revised: May 2022

Manufactured by:
AustarPharma, LLC

18 Mayfield Ave
Edison, NJ 08837, USA

Principal Display Panel - 500mg 500counts

NDC: 35561-212-13

Methocarbamol
500mg 500 tablet (s)

RX only



NDC 35561-212-13

**Methocarbamol
Tablets USP**
500 mg

500 TABLETS
Rx Only

USUAL ADULT DOSAGE: Two to four tablets four times daily.
See accompanying descriptive literature.
STORE at 20° to 25°C (68° to 77°F) [See USP controlled room temperature].
Dispense in a tight container.
Manufactured by: AustarPharma, LLC,
Edison, NJ 08837

LBL133
REV091118


N 3 35561 21213 6

LOT NO.:
EXP. DATE:

Principal Display Panel - 500mg 100 counts

NDC: 35561-212-12

Methocarbamol
500mg 100 tablet (s)

RX only



NDC 35561-212-12

Methocarbamol Tablets USP

500 mg

100 TABLETS
Rx Only

USUAL ADULT DOSAGE: Two to four tablets four times daily.

See accompanying descriptive literature.

STORE at 20° to 25°C (68° to 77°F) [See USP controlled room temperature].

Dispense in a tight container.

Manufactured by: AustarPharma, LLC,
Edison, NJ 08837

LBL132
REV091118



LOT NO.:
EXP. DATE:

Principal Display Panel - 750mg 100counts

NDC: 35561-211-12

Methocarbamol
750mg 100 tablet (s)

RX only



NDC 35561-211-12

Methocarbamol Tablets USP

750 mg

100 TABLETS
Rx Only

USUAL ADULT DOSAGE: Two tablets three times daily.

See accompanying descriptive literature.

STORE at 20° to 25°C (68° to 77°F) [See USP controlled room temperature].

Dispense in a tight container.

Manufactured by: AustarPharma, LLC,
Edison, NJ 08837

LBL134
REV091118



LOT NO.:
EXP. DATE:

Principal Display Panel - 750mg 500counts

NDC: 35561-211-13

Methocarbamol
750mg 500tablet (s)


RX only


NDC 35561-211-13
**Methocarbamol
Tablets USP**
750 mg

500 TABLETS
Rx Only

USUAL ADULT DOSAGE: Two tablets three times daily.
See accompanying descriptive literature.
STORE at 20° to 25°C (68° to 77°F) [See USP controlled room temperature].
Dispense in a tight container.
Manufactured by: AustarPharma, LLC,
Edison, NJ 08837

LBL135
REV091118



LOT NO.:
EXP. DATE:

Principal Display Pane - 1000 mg 100 counts

NDC: 35561-349-12

Methocarbamol
1000mg 100 tablet(s)

Rx Only



NDC 35561-349-12



Methocarbamol Tablets USP

1000 mg

100 TABLETS
Rx Only

Each film coated tablet contains: 1000 mg Methocarbamol USP
Dosage: See package insert for full prescribing information.
Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).
Dispense in tight container.
Keep this and all drugs out of the reach of children.
Manufactured by: AustarPharma LLC
Edison, NJ 08837

LBL414
REV052022



Principal Display Panel - 1000 mg 500 counts

NDC: 35561-349-13

Methocarbamol

1000 mg 5000 tablets

RX only



NDC 35561-349-13



Methocarbamol Tablets USP

1000 mg

500 TABLETS
Rx Only

Each film coated tablet contains: 1000 mg Methocarbamol USP
Dosage: See package insert for full prescribing information.
Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).
Dispense in tight container.
Keep this and all drugs out of the reach of children.
Manufactured by: AustarPharma LLC
Edison, NJ 08837

LBL415
REV052022



METHOCARBAMOL

methocarbamol tablet, coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:35561-349
---------------------	-------------------------	---------------------------	---------------

Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	1000 mg	
Product Characteristics				
Color	orange	Score	2 pieces	
Shape	RECTANGLE	Size	21mm	
Flavor		Imprint Code	AP349	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35561-349-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/16/2022	
2	NDC:35561-349-13	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/16/2022	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA200958	07/16/2022		

METHOCARBAMOL			
methocarbamol tablet, coated			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:35561-211
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	750 mg
Product Characteristics			
Color	white	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	AP211

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35561-211-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/09/2018	
2	NDC:35561-211-13	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/09/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200958	10/09/2018	

METHOCARBAMOL

methocarbamol tablet, coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:35561-212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	500 mg

Product Characteristics

Color	orange	Score	2 pieces
Shape	ROUND	Size	13mm
Flavor		Imprint Code	AP212
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35561-212-12	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/10/2018	
2	NDC:35561-212-13	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/10/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200958	10/10/2018	

Labeler - Bostal LLC (362785011)

Revised: 2/2026

Bostal LLC