

LEADER LIQUID BANDAGE- benzethonium chloride dyclonine hydrochloride liquid
LEADER/ Cardinal Health 110, Inc.

Leader Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic
Topical analgesic

Uses

- first aid to prevent bacteria contamination in minor cuts and scrapes

Warnings

For external use only.

Flammable:

do not use while smoking or near heat or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor
- on infected areas or wounds that are draining
- with other first aid products such as lotions and creams
- over sutures
- on mucous membranes

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious bleeding
- diabetes
- poor circulation
- serious burns

When using this product

- do not store at temperature above 120°F

Intentional misuse by deliberately concentrating and inhaling the contents can be harmful or fatal.

Stop use and ask a doctor if

- condition persists or gets worse
- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- Clean affected area
- apply a small amount on the area 1-3 times daily
- let dry
- a second coating may be applied for extra protection
- to remove, apply more Liquid Bandage and quickly wipe off
- finger nail polish remover may dissolve Liquid Bandage

Other information

- **cap bottle tightly**
- store at room temperature away from heat
- may temporarily sting upon application
- do not allow to come in contact with floors, countertops, or other finished surfaces - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

antiseptic

liquid

bandage

helps prevent infection

for small cuts and wounds

invisible, flexible, waterproof, easy to use

UP & UP

NET WT 1 FL OZ (29.6 mL)



DO NOT USE IF NECKBAND IS BROKEN

LEADER™

Drug Facts

Active Ingredients
Benzethonium chloride 0.2%.....
Dyclonine hydrochloride 0.75%.....
Purpose
First aid antiseptic
Topical analgesic

Uses
■ first aid to prevent bacterial contamination in minor cuts, scrapes and burns
■ for temporary relief of pain

Warnings
For external use only.
Flammable: Keep away from fire or flame.

Do not use ■ in the eyes ■ over large areas of the body
■ longer than 1 week unless directed by a doctor
Ask a doctor before use if you have ■ deep or puncture wounds
■ deep cuts ■ animal bites ■ serious burns
When using this product ■ avoid contact with eyes
Stop use and ask a doctor if ■ condition persists or gets worse
■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ infection occurs

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions ■ for adults and children 2 years of age and older ■ clean affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ second coating may be applied for extra protection
■ children under 2 years of age: consult a doctor

Other information ■ cap bottle tightly ■ store between 20° to 25°C

OVERLAY

DO NOT USE IF NECKBAND IS BROKEN

Active ingredients: Benzethonium Chloride 0.2% and Dyclonine Hydrochloride 0.75% **Purpose:** First aid antiseptic and topical analgesic **Uses:** First aid to prevent bacterial contamination in minor cuts, scrapes, and burns. For the temporary relief of pain.

Warnings: For external use only. Flammable: Keep away from fire or flame. **Do not use** in eyes, over large areas of the body, and longer than 1 week unless directed by a doctor.

Liquid Bandage

Benzethonium Chloride, 0.2%
Dyclonine Hydrochloride, 0.75%
First Aid Antiseptic
Topical Analgesic

Seals to Protect
For Small Cuts & Wounds
Water-Resistant

1.0 FL OZ (29.6 mL)

Ask a doctor before use if you have deep or puncture wounds, deep cuts, animal bites and serious burns.

When using this product avoid contact with eyes.

Stop use and ask a doctor if condition persists or gets worse, symptoms persist for more than 7 days or clear up and occur again within a few days, and infection occurs.

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

Directions: For adults and children 2 years of age and older. Clean affected area. Apply a small amount on the area 1 to 3 times daily and let dry. A second coating may be applied for extra protection. Children under 2 years of age: consult a doctor.

Other information: Cap bottle tightly. Store between 20° to 25°C (68° to 77°F). May temporarily sting upon application.

To remove, apply more Liquid Bandage and quickly wipe off. **Inactive ingredients:** acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions? Call 1-800-200-6313

Do not allow to come into contact with fabrics, floors, countertops or other surfaces -- will stain.

DIST. BY CAH DUBLIN OH, 43017 ©2022 Cardinal Health

www.myleader.com

1-800-200-6313

100% Money Back Guarantee

Return to place of purchase if not satisfied.

CIN 5768684 REV. 3/22

50-033LD

LEADER LIQUID BANDAGE

benzethonium chloride dyclonine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0599
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 10 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0599-1	10 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product	04/12/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/12/2022	

Labeler - LEADER/ Cardinal Health 110, Inc. (063997360)

Revised: 2/2024

LEADER/ Cardinal Health 110, Inc.