

LEADER LIQUID BANDAGE- benzethonium chloride dyclonine hydrochloride liquid
LEADER/ Cardinal Health 110, Inc.

Leader Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%
Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic
Topical analgesic

Uses

- first aid to prevent bacterial contamination in minor cuts, scrapes and burns
- for the temporary relief of pain

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep puncture wounds
- deep cuts
- animal bites
- serious burns

When using this product

- avoid contact with eyes

Stop use and ask a doctor if

- condition persists or gets worse
- symptoms persist for more than 7 days or clear up and occur again within a few days

- infection occurs

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1-3 times daily and let dry
- a second coating may be applied for extra protection
- children under 2 years of age: consult a doctor

Other information

- **cap bottle tightly**
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

LEADER

Liquid

Bandage

Benzethonium Chloride, 0.2%
First Aid Antiseptic

Dyclonine Hydrochloride, 0.75%
Topical Analgesic

**Seals to Protect
For Small Cuts & Wounds**

Prevents Infection

Invisible & Flexible

Water-Resistant

Built-In Applicator Brush

1.0 FL OZ (29.6 mL)



Liquid Bandage

**Benzethonium Chloride, 0.2%
Dyclonine Hydrochloride, 0.75%
First Aid Antiseptic | Topical Analgesic**

**Seals to Protect
For Small Cuts & Wounds**
Prevents Infection
Invisible & Flexible
Water-Resistant
Built-In Applicator Brush

**COMPARE TO
NEW-SKIN®
active ingredient***

100% Money Back Guarantee

1.0 FL OZ (29.6 mL)

Use Liquid Bandage For:
Blisters, Chapped & Cracked Fingertips, Hangnails, Paper Cuts,
Shaving Nicks & Help in Preventing the Formation of Calluses.

Great For:
Athletes, Active Lifestyles, Outdoor Sportsmen & Musicians.

53-033LD



Liquid Bandage

Benzethonium Chloride, 0.2%
Dyclonine Hydrochloride, 0.75%
First Aid Antiseptic
Topical Analgesic

Seals to Protect
For Small Cuts & Wounds



Liquid Bandage

Benzethonium Chloride, 0.2% | Dyclonine Hydrochloride, 0.75%
First Aid Antiseptic | Topical Analgesic

Seals to Protect
For Small Cuts & Wounds

**NO VARNISH
AREA
(WHITE ONLY)**

©2022 Cardinal Health. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY
CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com
1-800-200-6313

Essential to Care™ since 1979

100% Money Back Guarantee
Return to place of purchase if not satisfied.



CIN 5768684 REV. 3/22



*This product is not manufactured or distributed by Advance Health, LLC, owner of the registered trademark New-Skin®.


Care
■ Wipe top of bottle clean after each use.
■ Do not allow to come into contact with fabrics, floors, countertops or other surfaces – will stain.

Questions? Call 1-800-200-6313

Inactive ingredients nitrocellulose, SD alcohol 40, acetone, amyl acetate, castor oil, ethyl acetate, more Liquid Bandage and quickly wipe off

Other information ■ cap bottle tightly ■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting upon application ■ to remove, apply affected area ■ apply a small amount on the area 1 to 3 times daily and let dry ■ a second coating may be applied for extra protection ■ children under 2 years of age: consult a doctor

Directions ■ for adults and children 2 years of age and older ■ clean ■ infection occurs ■ few days ■ Control Center right away. ■ Keep out of reach of children. If swallowed, get medical help or call a Poison



LEADER LIQUID BANDAGE

benzethonium chloride dyclonine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0599
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	0.2 mg in 10 mL
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	0.75 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ACETONE (UNII: 1364PS73AF)	
AMYL ACETATE (UNII: 92Q24NH7AS)	
CASTOR OIL (UNII: D5340Y2I9G)	
ETHYL ACETATE (UNII: 76845O8NMZ)	
PYROXYLIN (UNII: KYR8BR2X6O)	
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0599-1	10 mL in 1 BOTTLE, WITH APPLICATOR; Type 0: Not a Combination Product	04/12/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	04/12/2022	

Labeler - LEADER/ Cardinal Health 110, Inc. (063997360)

Revised: 12/2025

LEADER/ Cardinal Health 110, Inc.