

GRAVOL- dimenhydrinate capsule, liquid filled
Church & Dwight Co., Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Gravol™ Motion Sickness Relief - Liquid Gels

Active ingredient (in each Liquid Gel)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

For prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase dizziness
- be careful when driving a motor vehicle or operating machinery or any activity that requires alertness

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

adults and children 12 years and over:	<ul style="list-style-type: none">• take 1 to 2 liquid gels every 4-6 hours• do not take more than 8 liquid gels in 24 hours, or as directed by a doctor
children 6 to under 12 years:	<ul style="list-style-type: none">• give 1 liquid gel every 6-8 hours• do not give more than 3 liquid gels in 24 hours, or as directed by a doctor

Other information

- store at room temperature 59°-77°F (15°-25°C)
- **do not use if packet is torn**

Inactive ingredients

Gelatin, glycerin, mannitol, polyethylene glycol 400, povidone, purified water, Red 33, sorbitan, sorbitol

Questions or comments?

1-800-952-5080 M-F 9am to 5pm ET

PRINCIPAL DISPLAY PANEL

OVER 70 YEARS OF
QUALITY & TRUST†

Prevents and Treats:

Nausea & Vomiting

Dizziness*

*Prevention and treatment of nausea,
vomiting and dizziness due to motion sickness

Ages 6+

Gravol™ LIQUID GELS

Gravol™

MOTION SICKNESS RELIEF

Dimenhydrinate 50 mg - Antiemetic

8

LIQUID GELS

(Liquid Filled

Capsules)

TASTE FREE

EASY TO SWALLOW

Prevents and Treats:

Nausea & Vomiting

Dizziness*

*Prevention and treatment of nausea, vomiting and dizziness due to motion sickness

Ages 6+



Gravol™ LIQUID GELS

Gravol™

MOTION SICKNESS RELIEF

Dimenhydrinate 50 mg – Antiemetic

8

LIQUID GELS
(Liquid Filled Capsules)

TASTE FREE
EASY TO SWALLOW



Gravol™ LIQUID GELS

Drug Facts	
Active ingredient (in each Liquid Gel) Dimenhydrinate 50 mg	Purpose Antiemetic
Use For the prevention and treatment of these symptoms associated with motion sickness: • nausea • vomiting • dizziness	
Warnings Do not use for children under 2 years of age unless directed by a doctor. Ask a doctor before use if you have: • a breathing problem such as emphysema or chronic bronchitis • glaucoma • trouble urinating due to an enlarged prostate gland Ask a doctor or pharmacist before use if you are taking: sedatives or tranquilizers When using this product: • marked drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • be careful when driving a motor vehicle or operating machinery or any activity that requires alertness If pregnant or breast-feeding, ask a doctor before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions • To prevent motion sickness, the first dose should be taken 1/2 to 1 hour before starting activity. • To prevent or treat motion sickness, see below: • adults and children 12 years and over: take 1 to 2 liquid gels every 4-6 hours • do not take more than 8 liquid gels in 24 hours, or as directed by a doctor • children 6 to under 12 years: • give 1 liquid gel every 6-8 hours • do not give more than 3 liquid gels in 24 hours, or as directed by a doctor Other information • store at room temperature 59°-77°F (15°-25°C) • do not use if packet is torn	
Inactive ingredients Gelatin, glycerin, mannitol, polyethylene glycol 400, powders, purified water, FD-33, sorbitan, sorbitol	
Questions? Call us at 1-800-952-5080 M-F 9am to 5pm ET	

Made in Canada
Church & Dwight Co., Inc., Ewing, NJ 08628
Gravol is a trademark of Church & Dwight Canada Corp.



Gravol™

GRC-00163-02 1202277

GRAVOL

dimenhydrinate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:10237-940
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C00X)	
MANNITOL (UNII: 3OWL53L36A)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
WATER (UNII: 059QF0KO0R)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	pink	Score	score with uneven pieces
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10237-940-08	1 in 1 CARTON	03/02/2022	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	03/02/2022	

Labeler - Church & Dwight Co., Inc. (001211952)

Establishment

Name	Address	ID/FEI	Business Operations
Church & Dwight Canada Corp.		253933600	manufacture(10237-940)

Revised: 3/2022

Church & Dwight Co., Inc.