

HYDROXYZINE HYDROCHLORIDE- hydroxyzine hydrochloride tablet Direct Rx

Hydroxyzine HCL

Hydroxyzine Hydrochloride Tablets, USP are available as follows:

10 mg tablets: round, film coated white tablets. Debossed H on one side and 500 on the reverse side. They are available as follows:

Bottles of 100: NDC 23155-500-01

Bottles of 500: NDC 23155-500-05

Bottles of 1000: NDC 23155-500-10

25 mg tablets: round, film coated white tablets. Debossed H over 501 on one side and plain on the reverse side. They are available as follows:

Bottles of 100: NDC 23155-501-01

Bottles of 500: NDC 23155-501-05

Bottles of 1000: NDC 23155-501-10

50 mg tablets: round, film coated white tablets. Debossed H over 502 on one side and plain on the reverse side. They are available as follows:

Bottles of 100: NDC 23155-502-01

Bottles of 500: NDC 23155-502-05

Bottles of 1000: NDC 23155-502-10

Dispense in a tight container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Distributed by:

Avet Pharmaceuticals Inc.

East Brunswick, NJ 08816

1.866.901.DRUG (3784)

logo

51U000000193US07

Rev. 10/2019

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested: adults, 50 to 100 mg q.i.d.; children under 6 years, 50 mg daily in divided doses; children over 6 years, 50 to 100 mg daily in divided doses.

For use in the management of pruritus due to allergic conditions such as chronic

urticaria and atopic and contact dermatoses and in histamine-mediated pruritus: adults, 25 mg t.i.d. or q.i.d.; children under 6 years, 50 mg daily in divided doses; children over 6 years, 50 to 100 mg daily in divided doses.

As a sedative when used as a premedication and following general anesthesia: 50 to 100 mg for adults and 0.6 mg/kg of body weight in children.

When treatment is initiated by the intramuscular route of administration, subsequent doses may be administered orally.

As with all potent medication, the dosage should be adjusted according to the patient's response to therapy.

The most common manifestation of hydroxyzine overdose is hypersedation. Other reported signs and symptoms were convulsions, stupor, nausea and vomiting. As in the management of overdose with any drug, it should be borne in mind that multiple agents may have been taken.

If vomiting has not occurred spontaneously, it should be induced. Immediate gastric lavage is also recommended. General supportive care, including frequent monitoring of the vital signs and close observation of the patient, is indicated. Hypotension, though unlikely, may be controlled with intravenous fluids and levarterenol or metaraminol. Do not use epinephrine as hydroxyzine counteracts its pressor action.

Hydroxyzine overdose may cause QT prolongation and Torsade de Pointes. ECG monitoring is recommended in cases of hydroxyzine overdose.

There is no specific antidote. It is doubtful that hemodialysis would be of any value in the treatment of overdose with hydroxyzine. However, if other agents such as barbiturates have been ingested concomitantly, hemodialysis may be indicated. There is no practical method to quantitate hydroxyzine in body fluids or tissue after its ingestion or administration.

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested.

Useful in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses and in histamine-mediated pruritus.

As a sedative when used as a premedication and following general anesthesia, hydroxyzine may potentiate meperidine and barbiturates, so their use in pre-anesthetic adjunctive therapy should be modified on an individual basis. Atropine and other belladonna alkaloids are not affected by the drug. Hydroxyzine is not known to interfere with the action of digitalis in any way and it may be used concurrently with this agent.

The effectiveness of hydroxyzine as an antianxiety agent for long term use, that is more than 4 months, has not been assessed by systematic clinical studies. The physician should reassess periodically the usefulness of the drug for the individual patient.

Hydroxyzine hydrochloride is unrelated chemically to the phenothiazines, reserpine, meprobamate or the benzodiazepines. Hydroxyzine is not a cortical depressant, but its action may be due to a suppression of activity in certain key regions of the subcortical area of the central nervous system.

Primary skeletal muscle relaxation has been demonstrated experimentally. Bronchodilator activity, and antihistaminic and analgesic effects have been

demonstrated experimentally and confirmed clinically. An antiemetic effect, both by the apomorphine test and the veriloid test, has been demonstrated.

Pharmacological and clinical studies indicate that hydroxyzine in therapeutic dosage does not increase gastric secretion or acidity and in most cases has mild antisecretory activity.

Hydroxyzine is rapidly absorbed from the gastrointestinal tract and hydroxyzine's clinical effects are usually noted within 15 to 30 minutes after oral administration.

Hydroxyzine hydrochloride, USP has the chemical name of (±)-2-[2-[4-(p-Chloro-α-phenylbenzyl)-1-piperaziny]ethoxy]ethanol dihydrochloride.

structure

Molecular Formula: C₂₁H₂₇ClN₂O₂ · 2HCl Molecular Weight: 447.83

Hydroxyzine hydrochloride, USP occurs as a white, odorless powder which is very soluble in water.

Each tablet for oral administration contains 10 mg, 25 mg, or 50 mg hydroxyzine hydrochloride, USP. Inactive ingredients include anhydrous lactose, colloidal silicon dioxide, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate and titanium dioxide.

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed. Dosage: See package insert. Store between 68-77 degrees F. For RX ONLY. Keep out of reach of children.



Lot# 03MA0218
Hydroxyzine HCL 10mg
Packaged and
Distributed By: DIRECT
72189-333-30

Hydroxyzine HCL
10mg
30 Tabs

Generic For: Atarax
Each tablet contains Hydroxyzine hydrochloride,
USP 10 mg

Discontinued
72189-333-30
03MA0218
Davenport, GA 30534
81277

Dist. By Anet Pharmaceuticals Inc.
Emt Brunswick, NJ 08816
NDC 22189-333-30
Mfg NDC 23155-500-05

Mfg Lot: 20921021A
TR 3/3/2022 9277472

Hydroxyzine HCL 10mg
NDC 22189-333-30 30 Tabs
Lot 03MA0218 Exp 6/30/23
Mfg NDC 23155-500-05

Hydroxyzine HCL 10mg
NDC 22189-333-30 30 Tabs
Lot 03MA0218 Exp 6/30/23
Mfg NDC 23155-500-05

Hydroxyzine HCL 10mg
NDC 22189-333-30 30 Tabs
Lot 03MA0218 Exp 6/30/23
Mfg NDC 23155-500-05

Caution: Federal law prohibits transfer of this drug to any person other than the patient for whom it was prescribed. Dosage: See package insert. Store between 68-77 degrees F. For RX ONLY. Keep out of reach of children.



NDC 72189-333-72

Hydroxyzine HCL

10mg

120 Tabs

Generic For: **Atarax**

Each tablet contains: Hydroxyzine hydrochloride, USP 10 mg

Lot# 060C2504
 Prod# 4206-010-72
 Packaged and Distributed By: **DIRECT Rx**



Discard After: 5/31/27
 72189-333-72
 060C2504 Dawsonville, GA 30534
 5/31/27
 CR7EF

Dist. By Avet Pharmaceuticals Inc.
 East Brunswick, NJ 08816
 NDC 23155-500-10

Mfg Lot: 209250177A
 KS 10/6/2025 3107237

Hydroxyzine HCL 10mg
 NDC 72189-333-72 120 Tabs
 Lot 060C2504 Exp 5/31/27
 Mfg NDC 23155-500-10

Hydroxyzine HCL 10mg
 NDC 72189-333-72 120 Tabs
 Lot 060C2504 Exp 5/31/27
 Mfg NDC 23155-500-10

Hydroxyzine HCL 10mg
 NDC 72189-333-72 120 Tabs
 Lot 060C2504 Exp 5/31/27
 Mfg NDC 23155-500-10

Hydroxyzine HCL 10mg
 NDC 72189-333-72 120 Tabs
 Lot 060C2504 Exp 5/31/27
 Mfg NDC 23155-500-10

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:72189-333(NDC:23155-500)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROXYZINE DIHYDROCHLORIDE (UNII: 76755771U3) (HYDROXYZINE - UNII:30S50YM8OG)	HYDROXYZINE DIHYDROCHLORIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CROSPVIDONE (UNII: 2S7830E561)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	7mm

Flavor		Imprint Code	H;500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72189-333-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	03/03/2022	
2	NDC:72189-333-72	120 in 1 BOTTLE; Type 0: Not a Combination Product	03/03/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204279	03/03/2022	

Labeler - Direct Rx (079254320)

Registrant - Direct Rx (079254320)

Establishment

Name	Address	ID/FEI	Business Operations
DirectRx		079254320	repack(72189-333)

Revised: 10/2025

Direct Rx