

RITE AID LAXATIVE SALINE LAXATIVE- sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema
RITE AID

RITE AID LAXATIVE SALINE ENEMA LAXATIVE

Drug Facts

<i>Active ingredients (in each 118 mL delivered dose)</i>	<i>Purpose</i>
Dibasic Sodium Phosphate 7 g	Saline laxative
Monobasic Sodium Phosphate 19 g	Saline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

For rectal use only.

Dosage warning

Using more than one enema in 24 hours can be harmful.

Do not use

- on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- **do not use more than directed. Serious side effects may occur from excess dosage**

- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours)

Do not use if taking another sodium phosphate product.

Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other information

- **each 118 ml contains** 4.3 g sodium
- store at 15 to 30°C (59 to 86°F)
- additional liquids by mouth are recommended while using this product
- **Tamper Evident:** DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.

Inactive ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments?

866-323-0107

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,
CAMP HILL, PA 17011

PRINCIPAL DISPLAY PANEL - 133 mL Bottle Carton

NDC 11822-0604-1

FREE
FROM
PARABEN
FREE

Compare to the active
ingredients in Fleet[®] Enema*

LAXATIVE
SALINE
ENEMA

DIBASIC SODIUM PHOSPHATE 7 g
MONOBASIC SODIUM PHOSPHATE 19 g

READY TO USE

not made with natural rubber latex

for relief of occasional constipation

easy, soft squeeze bottle

2
BOTTLES

4.5 FL OZ (133 mL) EACH

PL1016B
V18

CAUTION:
REMOVE
PROTECTIVE
SHIELD FROM
ENEMA TIP
BEFORE
INSERTING

LAXATIVE SALINE ENEMA

DIBASIC SODIUM PHOSPHATE 7 g
MONOBASIC SODIUM PHOSPHATE 19 g

NDC 11822-0604-1

FREE
F R M

**PARABEN
FREE**

Compare to the active
ingredients in **Fleet® Enema***

POSITIONS FOR USING THIS ENEMA

Left side position: Lie on left side with knees bent and arms resting comfortably.



Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.



REMOVE PROTECTIVE SHIELD FROM ENEMA TIP BEFORE INSERTING:

Hold bottle upright, grasping bottle cap with fingers. Grasp protective shield with other hand and pull gently to remove.



ADMINISTERING ENEMA:

- With steady pressure, gently insert enema tip into rectum with a slight side-to-side movement, with tip pointing toward navel. Insertion may be easier if person receiving enema bears down, as if having a bowel movement. Doing so helps relax muscles around anus.
- Do not force enema tip into rectum. Stop using if tip is hard to insert. Forcing the tip into the rectum can cause injury (especially if you have hemorrhoids). If enema tip causes rectal bleeding or pain, get immediate medical care.
- Squeeze bottle until nearly all liquid is gone. It is not necessary to empty bottle completely, as it contains more liquid than needed.
- Remove tip from rectum and stay in position until urge to evacuate is strong (usually 1 to 5 minutes). If no urge is felt after 5 minutes of using, try to empty bowel. Call a doctor promptly if no liquid comes out of the rectum after 30 minutes because dehydration could occur.

LAXATIVE SALINE ENEMA

DIBASIC SODIUM PHOSPHATE 7 g
MONOBASIC SODIUM PHOSPHATE 19 g

READY TO USE

not made with natural rubber latex
for relief of occasional constipation
easy, soft squeeze bottle



2
BOTTLES

4.5 FL OZ (133 mL) EACH

Lot# FPO

Exp: FPO

CARTON SEALED
FOR SAFETY. DO
NOT USE IF TOP
OR BOTTOM
FLAP OF CARTON
IS TORN OR
MISSING.

COMPLETE ENEMA IN A DISPOSABLE SQUEEZE BOTTLE WITH SOFT, PRE-LUBRICATED TIP

Protective
shield prevents
contamination

Comfortable,
lubricated tip
for ease of
insertion

One-way
safety valve
controls flow
and prevents
reflux

Easy, soft
squeeze
bottle



BOTTLE, CAP AND SHIELD ARE
NOT MADE WITH NATURAL
RUBBER LATEX

ALLERGIC REACTION TO
MATERIALS CONTAINING LATEX IS
A GROWING AND SERIOUS
MEDICAL PROBLEM

Drug Facts

Active Ingredients (in each 118 mL contains sodium 4.3 g)

Dibasic Sodium Phosphate 7 gSaline laxative
Monobasic Sodium Phosphate 19 gSaline laxative

Use

- relieves occasional constipation
- this product generally produces bowel movement in 1 to 5 minutes

Warnings

Dosage warning: Using more than one enema in 24 hours can be harmful.

For rectal use only.

Do not use ■ on children under 2 years of age

Ask a doctor before use if you

- have already used a laxative for more than 3 days
- have kidney disease, have heart problems, or are dehydrated
- are 55 years of age or older
- are on a sodium-restricted diet
- have abdominal pain, nausea, or vomiting
- have a sudden change in bowel habits lasting more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

When using this product

- do not use more than directed. Serious side effects may occur from excess dosage**
- do not use for more than 3 days, without asking a doctor

Stop use and ask a doctor if

- you have rectal bleeding
- you have no bowel movement within 30 minutes of enema use
- you have symptoms of dehydration (thirstiness, dizziness, vomiting, urinating less often than normal)

These symptoms may indicate a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away: 800-222-1222.

Directions (or as directed by a doctor)

Single daily dosage (per 24 hours)

Do not use if taking another sodium phosphate product.

Do not use more unless directed by a doctor. See Warnings.

Adults and children 12 years of age and older	1 bottle once daily
Children 2 to under 12 years of age	Use Pediatric Enema
Children under 2 years of age	DO NOT USE

Other Information

*This product is not manufactured or distributed by C.B. Fleet Company Inc., owner of the registered trademark Fleet®.

Other information

- each 118 ml contains sodium 4.3 g
- store at 15° to 30°C (59° to 86°F)
- additional liquids by mouth are recommended while using this product
- **Tamper Evident:** DO NOT USE IF TOP OR BOTTOM FLAP OF CARTON IS TORN OR MISSING.

Inactive Ingredients

benzalkonium chloride, disodium EDTA, purified water

Questions or comments? 866-323-0107



DISTRIBUTED BY:
 RITE AID, 30 HUNTER LANE,
 CAMP HILL, PA 17011
www.riteaid.com
MADE IN USA

**SATISFACTION
 GUARANTEE**
 If you're not satisfied, we'll
 happily refund your money.

RITE AID LAXATIVE SALINE LAXATIVE

sodium phosphate, dibasic, unspecified form and sodium phosphate, monobasic, unspecified form enema

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0604
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74) (PHOSPHATE ION - UNII:NK08V8K8HR, SODIUM CATION - UNII:LYR4MONH37)	SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM	7 g in 118 mL
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW) (PHOSPHATE ION - UNII:NK08V8K8HR, SODIUM CATION - UNII:LYR4MONH37)	SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM	19 g in 118 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0604-1	2 in 1 CARTON	12/03/2019	
1		133 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:11822-0604-2	6 in 1 CARTON	12/03/2019	
2		133 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:11822-0604-3	12 in 1 CARTON	12/03/2019	
3		133 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph drug	M007	12/03/2019	

Labeler - RITE AID (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
Natureplex, LLC		062808196	MANUFACTURE(11822-0604)

Revised: 12/2024

RITE AID