# NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled WinCo Foods, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

## **Drug Facts**

# Active ingredients (in each softgel)

# Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

# Purpose

#### Pain reliever/Fever Reducer

**Cough Suppressant** 

Antihistamine

#### Uses

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - o minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

# **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**: If sore throat is severe, persists more than 2 days, is accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor of pharmacist before taking this product.

# Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

# Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

# When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

# Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

# If pregnant or breast-feeding,

ask a health professional before use.

### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

#### Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

# **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

# Questions or comments?

Call 1-800-824-1706 Monday-Friday 9AM-5PM MST

# **Principal Display Panel**

†Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu LiquiCaps®

MULTI-SYMPTOM

Cold & Flu Relief

ACETAMINOPHEN / PAIN RELIEVER - FEVER REDUCER

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT

DOXYLAMINE SUCCINATE / ANTIHISTAMINE

Relieves: Aches, Fever, Sore throat, Cough, Sneezing & Runny Nose

Alcohol-free

Softgels

\*\*Liquid-Filled Capsules

†This product is not manufactured or distributed by the Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademark of the Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

#### **DISTRIBUTED BY:**

WINCO FOODS, LLC,

**BOISE, ID 83704** 

www.wincofoods.com

#### **Package Label**

TZM mq4 - me9 yebiri - yebnoM 80r1-428-008 lls3 usuyeq quowaruesa wsi\ occnr ■ SVOID SICOUORIC DIOVE Questions or comments? excuspinty may occur, especially in children When using this product **a** do not exceed recommended dosage sorbitam, sorbitol, white ink ■ psking sedatives or transmizers wannidol, polyednýtene glycol, povidone, propytene glycol, pumhed water, ■ taking the blood thinning drug warfarin inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, Wak a doctor or pharmacist before use if you are ■ store between 15-30°C (59-86°F)
■ avoid excessive heat conôµ дият осспиз миди доо шлеци byjeðuu (шлепа) Отнег іптогталью amoking, astima, chronic bronchibs or emphysema a breathing problem or chronic cough that lasts such as occurs with label to ensure correct dosing puej6 ajejsoud pe6jejue we of eub griñennu elduori ... when using other Daytime or Mighttime products, carefully read each ■ djancowa Wax a doctor before use if you have I liver disease ■ cyliqueu nugeu 15 years: do not use ■ swallow whole; do not crush, chew, or dissolve pharmacist before taking this product. not lonow if your prescription drug contains an MAOI, ask a doctor or adults and children 12 years and over: take 2 softgels with water every Parkinson's disease), or for 2 weeks after stopping the MADI drug. If you do ■ do not take more than 4 doses in 24 hours (certain drugs for depression, psychiathc, or emotional conditions, or DILECTIONS ■ do not take more than directed (see Overdose warming) If you are now taking a prescription monoamine oxidase inhibitor (MAN) acetaminophen, ask a doctor or pharmacist. adults as well as for children even if you do not notice any signs or symptoms. uoubusecubgiou): It you are not sure whether a drug contains Genter (1-800-222-1 222) nght away. Quick medical afternoon is critical for миди зий одлет сило сощений эсеранновией (въезсифрой от damage, in case of overdose, get medical help or contact a Poison Control asn 100 0g Overdose warming: Taking more than the recommended dose can cause liver coverig a doctor promptly. Keep out of reach of children. accompanied or followed by fever, headache, rash, nausea, or vomiling, If pregnant or breast-feeding, ask a health professional before use. Sore throat warming: If sore throat is severe, persists for more than 2 days, is These could be signs of a serious condition. If a skin reaction occurs, stop use and seek medical help right away. condu course pack or occurs with resh or headache that lasts useu ■ wsy include: 

skin reddening 

plisters ■ redness or swelling is present ■ new symptoms occur ■ tever gets worse or lasts more than 3 days

yllergy alert: Acetaminophen may cause severe sidn reactions. Symptoms 3 or more alcoholic drinks every day while using this product

■ with other drugs containing acetaminophen

may occur if you take: In more than 4,000 mg of acetaminophen in 24 hours Liver warning: This product contains acetaminophen. Severe liver damage

Drug Facts (continued)

Doxylamine succinate 6.25 mg. animetaildhA cough due to minor throat and bronchial irritation pondy anbuessaut Burzaeus pue asou Áuuru . Dextromethorphan HBr 15 mg ■ sove throat ■ headache ■ minnor aches and pains naes 

I peur significación de la principa del principa de la principa del principa de la principa del principa de la principa del principa de la principa del sasoding Active ingredients (in each softgel) gund years (continued) Drug Facts

WinCo

■ beju ou condu ôspa wouse ou jesta mous grau \ qe\a

pe caustri when driving a motor vehicle or operating machinery

sicohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

**Drug Facts** (continued)

# **MULTI-SYMPTOM** Cold & Flu Relief

**Nighttime** 

Compare to the active ingredients in VICKS® NYQUIL® Cold & Flu LiquiCaps®

> Relieves: Aches, Fever, Sore Throat, Cough, Sneezing & Runny Nose

ACTUAL SIZE

Alcohol-free

\*Liquid-Filled Capsules

ACETAMINOPHEN/ PAIN RELIEVER - FEVER REDUCER DEXTROMETHORPHAN HBr/COUGH SUPPRESSANT DOXYLAMINE SUCCINATE/ANTIHISTAMINE

'This product is not manufactured or distributed by The Procter & Gamble Company

AMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704





# WINCO FOODS Nighttime Multi Symptom Cold & Flu Relief

# NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67091-338
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg		
DO XYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DO XYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg		

Inactive Ingredients		
Ingredient Name	Strength	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
GELATIN (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
PO VIDO NE (UNII: FZ989 GH94E)		
PROPYLENE GLYCOL (UNII: 6 DC9 Q16 7 V3)		
WATER (UNII: 059QF0KO0R)		
SORBITAN (UNII: 6O92ICV9RU)		
SORBITOL (UNII: 506T60A25R)		
MANNITOL (UNII: 3OWL53L36A)		

Product Characteristics			
Color	GREEN (Clear)	Score	no score
Shape	OVAL	Size	20 mm
Flavor		Imprint Code	P30
Contains			

Packaging			
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>

1	NDC:67091-338-24	24 in 1 CARTON	06/01/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
	M 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Annliestian Number or Managraph Citation	Mauliating Ctart Data	Marketing End Date
	Marketing Catego	ry Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	Marketing Catego FC MONOGRAPH FIN		06/01/2019	Marketing End Date

# Labeler - WinCo Foods, LLC (056098817)

Revised: 6/2019 WinCo Foods, LLC