

IBUPROFEN- ibuprofen tablet, film coated
Rite Aid Corporation

Rite Aid 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - backache
 - the common cold
 - headache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- shock
- hives
- facial swelling
- skin reddening
- blisters
- asthma (wheezing)
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- are age 60 or older
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart

failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist

- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

**RITE
AID®
PHARMACY**

****Compare to the active ingredient in Advil® Tablets**

**pain relief
ibuprofen**

ibuprofen tablets USP, 200 mg
pain reliever • fever reducer
(NSAID)

EASY TO OPEN CAP

**NOT FOR HOUSEHOLDS
WITH YOUNG CHILDREN**

ACTUAL SIZE

**150 COATED
TABLETS**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**


**This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil® Tablets.

50844 REV0618B29129

**DISTRIBUTED BY: RITE AID
30 HUNTER LANE
CAMP HILL, PA 17011**

**IF YOU'RE NOT
SATISFIED, WE'LL
HAPPILY REFUND**

YOUR MONEY.




****Compare to the active ingredient in Advil® Tablets**

pain relief ibuprofen

ibuprofen tablets USP, 200 mg
pain reliever • fever reducer (NSAID)

EASY TO OPEN CAP
NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN



ACTUAL SIZE

150 COATED TABLETS

Drug Facts (continued)

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor


Other information

- store between 20°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients carnitina wax, coloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?
Call 1-800-426-3391 9:30 AM-4:00 PM ET, Monday-Friday

ITEM 0370300



0 11822 37300 5

SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ACTUATION GUARANTEE

RITE AID PHARMACY

IF YOU'RE NOT SATISFIED, HAPPILY RETURN YOUR MONEY.

**** This product is not manufactured or distributed by Wyeth LLC, owner of the registered trademark Advil® tablets. 58844 REV0618829129**

DISTRIBUTED BY:
RITE AID, 30 HUNTER LANE
CAMP HILL, PA 17011

Directions

Take 1 tablet every 4 to 6 hours while symptoms persist. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Warnings

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- when using this product
 - take with food or milk if stomach upset occurs
 - stop use and ask a doctor if
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better

Precautions

- if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Do not take more than directed

- do not take more than directed
- the smallest effective dose should be used

Drug Facts (continued)

Warnings

- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed
- allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin redness
 - rash
 - blisters
 - hives
- if an allergic reaction occurs, stop use and seek medical help right away.

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - menstrual cramps
 - toothache
 - the common cold
 - headache
 - minor pain of arthritis
- temporarily reduces fever

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Drug Facts (continued)

Warnings

- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed
- allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin redness
 - rash
 - blisters
 - hives
- if an allergic reaction occurs, stop use and seek medical help right away.

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - menstrual cramps
 - toothache
 - the common cold
 - headache
 - minor pain of arthritis
- temporarily reduces fever

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient

ibuprofen USP
200 mg (NSAID) Pain reliever/fever reducer
(in each brown tablet)

Purpose

temporarily relieves minor aches and pains due to: backache, muscular aches, menstrual cramps, toothache, the common cold, headache, minor pain of arthritis, temporarily reduces fever

Drug Facts (continued)

Warnings

- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed
- allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin redness
 - rash
 - blisters
 - hives
- if an allergic reaction occurs, stop use and seek medical help right away.

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - menstrual cramps
 - toothache
 - the common cold
 - headache
 - minor pain of arthritis
- temporarily reduces fever

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed


RITE AID PHARMACY

****Compare to the active ingredient in Advil® Tablets**

pain relief ibuprofen

ibuprofen tablets USP, 200 mg
pain reliever • fever reducer (NSAID)

EASY TO OPEN CAP
NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN



ACTUAL SIZE

150 COATED TABLETS

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient

ibuprofen USP
200 mg (NSAID) Pain reliever/fever reducer
(in each brown tablet)

Purpose

temporarily relieves minor aches and pains due to: backache, muscular aches, menstrual cramps, toothache, the common cold, headache, minor pain of arthritis, temporarily reduces fever

Drug Facts (continued)

Warnings

- Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed
- allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 - facial swelling
 - asthma (wheezing)
 - shock
 - skin redness
 - rash
 - blisters
 - hives
- if an allergic reaction occurs, stop use and seek medical help right away.

Uses

- temporarily relieves minor aches and pains due to:
 - backache
 - muscular aches
 - menstrual cramps
 - toothache
 - the common cold
 - headache
 - minor pain of arthritis
- temporarily reduces fever

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- taking any other drug

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to any other NSAID, which may cause severe stomach bleeding. The chance is higher if you
 - are age 60 or older
 - take a blood thinning (anticoagulant) or steroid drug
 - take more or for a longer time than directed

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-0291
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0291-0	1 in 1 CARTON	05/24/1988	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-0291-3	1 in 1 CARTON	05/24/1988	
2		200 in 1 BOTTLE; Type 0: Not a Combination Product		
	NDC:11822-0291			

3	NDC:11822-0291-9	1 in 1 CARTON	05/24/1988	
3		150 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:11822-0291-5	1 in 1 CARTON	05/24/1988	
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:11822-0291-6	1 in 1 CARTON	05/24/1988	
5		16 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:11822-0291-2	1 in 1 CARTON	05/24/1988	
6		12 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:11822-0291-8	1 in 1 CARTON	05/24/1988	
7		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:11822-0291-1	1 in 1 CARTON	05/24/1988	
8		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
9	NDC:11822-0291-7	1 in 1 CARTON	05/24/1988	02/06/2022
9		250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
10	NDC:11822-0291-4	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - Rite Aid Corporation (014578892)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(11822-0291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(11822-0291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(11822-0291)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		832867837	PACK(11822-0291)
-------------------------	--	-----------	------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(11822-0291)

Revised: 6/2019

Rite Aid Corporation