

**NAPROXEN SODIUM- naproxen sodium tablet, coated**  
**Chain Drug Marketing Association**

-----  
**QCH - 1144 - 2019-1004**

***Drug Facts***

**Active ingredient (in each tablet)**

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID <sup>1</sup>)

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - menstrual cramps
  - headache
  - toothache
  - the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert**

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

### **Stop use and ask a doctor if**

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- you develop heartburn
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn

child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

adults and children 12 years and older	<ul style="list-style-type: none"><li>▪ take 1 tablet every 8 to 12 hours while symptoms last</li><li>▪ for the first dose you may take 2 tablets within the first hour</li><li>▪ do not exceed 2 tablets in any 8- to 12-hour period</li><li>▪ do not exceed 3 tablets in a 24-hour period</li></ul>
children under 12 years	<ul style="list-style-type: none"><li>▪ ask a doctor</li></ul>

### Other information

- **each tablet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).
- retain carton for complete product information

### Inactive ingredients

croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide

### PRINCIPAL DISPLAY PANEL

NDC 63868-465-50

QUALITY CHOICE

†Compare to the active ingredient in ALEVE®

Naproxen Sodium

Non-Prescription Strength

Naproxen Sodium Tablets, 220mg

# Pain Reliever / Fever Reducer (NSAID)

50 Tablets

COATING FREE AREA

INK AND COATING FREE FOR LOT AND EXPIRATION STAMPING

6 7 4 7 6 5 1 9



DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

NDC 63868-465-50



## Naproxen Sodium

Non-Prescription Strength

Naproxen Sodium Tablets, 220mg  
Pain Reliever | Fever Reducer (NSAID)

### 50 Tablets

**'Compare to the active ingredient in ALEVE'**



**Drug Facts (continued)**  
**Inactive ingredients** croscarmellose sodium, FD&C blue #2, macrogol, magnesium stearate, polyvinyl alcohol, povidone, pregelatinized starch, talc, titanium dioxide



© Distributed by C.D.M.A., Inc.  
43157 W. Nine Mile  
Novi, MI 48376-0995  
[www.qualitychoice.com](http://www.qualitychoice.com)  
Questions: 248-449-9300

F114402OCH\_R2

**Drug Facts (continued)**  
**Do not use** ■ If you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery  
**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you are taking a diuretic  
■ you have asthma  
■ you have problems or serious side effects from taking pain relievers or fever reducers  
**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking any other drug  
**When using this product**  
■ take with food or milk if stomach upset occurs  
■ long term continuous use may increase the risk of heart attack or stroke  
**Stop use and ask a doctor if**  
■ side effects occur. You may report side effects to FDA at 1-800-FDA-1088.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ you have difficulty swallowing  
■ it feels like the pill is stuck in your throat  
■ you develop heartburn  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**Drug Facts**  
**Active ingredient (in each tablet)** Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)  
**Purpose** Pain reliever/fever reducer  
\* nonsteroidal anti-inflammatory drug  
**Uses**  
■ temporarily relieves minor aches and pains due to:  
■ minor pain of arthritis  
■ headache  
■ backache  
■ menstrual cramps  
■ toothache  
■ the common cold  
■ temporarily reduces fever  
**Warnings**  
**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives  
■ facial swelling  
■ asthma (wheezing)  
■ shock  
■ skin reddening  
■ rash  
■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Drug Facts (continued)**  
**if pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
■ do not take more than directed  
■ the smallest effective dose should be used  
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)  
■ drink a full glass of water with each dose  
adults and children 12 years and older  
■ take 1 tablet every 8 to 12 hours while symptoms last  
■ for the first dose you may take 2 tablets within the first hour  
■ do not exceed 2 tablets in any 8- to 12-hour period  
■ do not exceed 3 tablets in a 24-hour period  
■ ask a doctor  
children under 12 years  
**Other information**  
■ each tablet contains: sodium 20 mg  
■ store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F).  
■ retain carton for complete product information

## NAPROXEN SODIUM

naproxen sodium tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-465
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

### Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	I3
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-465-50	1 in 1 CARTON	09/03/2009	12/31/2027
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63868-465-01	1 in 1 CARTON	09/03/2009	12/31/2027
2		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079096	09/03/2009	12/31/2027

