

IBUPROFEN- ibuprofen tablet, coated
Family Dollar (FAMILY WELLNESS)

Drug Facts

Active ingredient (in each Brown tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - minor pain of arthritis
 - toothache
 - backache
 - the common cold
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - slurred speech
 - leg swelling
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and older:
 - take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, iron oxide red, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

SEE NEW WARNINGS

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS†

IBUPROFEN

Tablets, USP 200 mg

Pain Reliever/Fever Reducer (NSAID)*

GLUTEN FREE

COATED TABLETS

*nonsteroidal anti-inflammatory drug

**TAMPER-EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT
INFORMATION.**

†This product is not manufactured or distributed by Haleon Group of Companies,
distributor of Advil® Tablets.

DISTRIBUTED BY:

MIDWOOD BRANDS, LLC

510 VOLVO PARKWAY,

CHESAPEAKE, VA 23320

Package Label

SKU 900556
 0 32251 93516 9
 PLO-8808A F0009648 Lot No.: Exp. Date:
 NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.
 DISTRIBUTED BY: MIDWOOD BRANDS, LLC
 510 VOLVO PKWY, CHESAPEAKE, VA 23320 USA
 PRODUCT OF INDIA

Drug Facts (continued)

Symptoms of heart problems or stroke: ■ chest pain ■ slurred speech ■ leg swelling ■ trouble breathing ■ weakness in one part or side of body ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and older: ■ take 1 tablet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 tablet, 2 tablets may be used ■ do not exceed 6 tablets in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

Other information ■ store at 20°-25°C (68°-77°F)

Inactive ingredients colloidal silicon dioxide, corn starch, hypromellose, magnesium stearate, ▶

Drug Facts (continued)

■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have ▶

Drug Facts (continued)

microcrystalline cellulose, pregelatinized starch, red iron oxide, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or comments?
 Call 1-877-763-3935 Monday-Friday 9AM-5PM EST

†This product is not manufactured or distributed by Haleon Group of Companies, distributor of Advil® Tablets.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts

Active ingredient (in each Brown tablet)
 ibuprofen 200 mg (NSAID)*
 *nonsteroidal anti-inflammatory drug

Purposes
 Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache ■ backache ■ the common cold ■ menstrual cramps ■ temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ▶

FAMILY Wellness™

SEE NEW WARNINGS

COMPARE TO THE ACTIVE INGREDIENT IN ADVIL® TABLETS†

IBUPROFEN
 Tablets, USP 200 mg
 Pain Reliever/Fever Reducer (NSAID)*
 GLUTEN FREE

24 COATED TABLETS
 *nonsteroidal anti-inflammatory drug


 ACTUAL SIZE

NDC 55319-252-24

FAMILY WELLNESS Ibuprofen 200 mg Tablets

IBUPROFEN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-252(NDC:59556-857)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND (round shaped)	Size	10mm
Flavor		Imprint Code	IBU200
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-252-25	1 in 1 BOX	09/15/2023	
1		250 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:55319-252-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	09/15/2023	
3	NDC:55319-252-24	1 in 1 BOX	09/15/2023	
3		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:55319-252-50	1 in 1 BOX	09/15/2023	
4		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:55319-252-10	1 in 1 BOX	09/15/2023	
5		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079129	09/15/2023	

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)

Revised: 10/2025

Family Dollar (FAMILY WELLNESS)