

**TOPCARE ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated**  
**Topco Associates LLC**

-----  
**Topco Associates LLC. Anti-Diarrheal Tablets Drug Facts**

**Active ingredient (in each caplet)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

**Inactive ingredients**

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

**Questions or comments?**

**1-888-423-0139**

**Principal Display Panel**

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

Anti-Diarrheal Tablets

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg

Anti-Diarrheal

● Controls the Symptoms of Diarrhea

12 CAPLETS†

†CAPSULE-SHAPED TABLETS

actual size

OPEN OTHER END  
**CODE AREA**

22453 88 C8

**Drug Facts (continued)**  
 D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch  
**Drug Facts (continued)**  
 Questions or comments?  
 1-888-423-0139

**Drug Facts (continued)**

**Drug Facts**  
 (in each caplet)  
 Loperamide HCl 2 mg..... Anti-diarrheal

**Purpose**  
 Use controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**  
 Do not use if you have ever had a severe allergic reaction to loperamide HCl  
 Heart alert: Taking more than directed can cause serious heart problems or death  
 Do not use if you have bloody or black stool  
 Ask a doctor before use if you have:  
 ■ fever ■ mucus in the stool  
 ■ a history of liver disease  
 ■ a history of abnormal heart rhythm  
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**  
 ■ symptoms get worse  
 ■ diarrhea lasts for more than 2 days  
 ■ you get abdominal swelling or bloating.  
 These may be signs of a serious condition.  
 If pregnant or breastfeeding, ask a health professional before use.

**Other information**  
 ■ see end panel for lot number and expiration date  
 ■ store at 20°-25°C (68°-77°F) (up to 33 lbs)  
 children under 2 years do not use  
 children 2-5 years (34 to 47 lbs) ask a doctor  
 children 6-8 years (48-59 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours  
 children 9-11 years (60-95 lbs) 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours  
 children 12 years and over 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours

**Directions**  
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ find night dose on chart if possible, use weight to dose; otherwise, use age.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).



**DO NOT USE IF  
 BLISTER UNIT  
 IS BROKEN  
 OR TORN**

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

DISTRIBUTED BY  
 TOPCO ASSOCIATES LLC  
 ELK GROVE VILLAGE  
 IL 60007  
 ©TOPCO PERA0521  
 QUESTIONS?  
 1-888-423-0139  
 topcare@topco.com  
 www.topcarebrand.com



Scan here for more information or call 1-888-423-0139



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.




NDC 36800-224-53

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT\*

# Anti-Diarrheal Tablets





**TOPCARE ANTI DIARRHEAL**  
 loperamide hydrochloride tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-224
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

**Product Characteristics**

<b>Color</b>	GREEN	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm

<b>Flavor</b>		<b>Imprint Code</b>	L2	
<b>Contains</b>				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-224-91	6 in 1 CARTON	08/07/2003	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:36800-224-53	12 in 1 CARTON	02/26/2003	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:36800-224-62	24 in 1 CARTON	04/22/2003	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:36800-224-80	1 in 1 CARTON	11/12/2007	03/31/2020
4		96 in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075232	02/26/2003		

**Labeler** - Topco Associates LLC (006935977)

Revised: 12/2025

Topco Associates LLC