

TOPCARE ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
Topco Associates LLC

Topco Associates LLC. Anti-Diarrheal Tablets Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-888-423-0139

Principal Display Panel

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

Anti-Diarrheal Tablets

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg

Anti-Diarrheal

● Controls the Symptoms of Diarrhea

12 CAPLETS†

†CAPSULE-SHAPED TABLETS

actual size

OPEN OTHER END
CODE AREA

22453 88 C8

Drug Facts (continued)
 D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch
Drug Facts (continued)
 1-888-423-0139
Questions or comments?

Drug Facts (continued)

Drug Facts
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ find night dose on chart if possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information
 ■ store at 20°-25°C (68°-77°F)
 ■ see end panel for lot number and expiration date

Warnings
 Do not use if you have ever had a severe allergic reaction to loperamide HCl or other allergic reaction to loperamide HCl
 ■ a history of liver disease
 ■ a history of abnormal heart rhythm
 ■ fever
 ■ mucus in the stool
 ■ Ask a doctor before use if you have
 ■ Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
 When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating.
 These may be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.

Use
 Use controls symptoms of diarrhea, including Travelers' Diarrhea

Active ingredient (in each caplet)
 Loperamide HCl 2 mg..... Anti-diarrheal

Purpose
 Loperamide HCl 2 mg..... Anti-diarrheal



DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

DISTRIBUTED BY
 TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE
 IL 60007
 ©TOPCO PERA0521
 QUESTIONS?
 1-888-423-0139
 topcare@topco.com
 www.topcarebrand.com



Scan here for more information or call 1-888-423-0139



This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.




NDC 36800-224-53

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT*

Anti-Diarrheal Tablets





TOPCARE ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm

Flavor		Imprint Code	L2	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-224-91	6 in 1 CARTON	08/07/2003	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:36800-224-53	12 in 1 CARTON	02/26/2003	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:36800-224-62	24 in 1 CARTON	04/22/2003	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:36800-224-80	1 in 1 CARTON	11/12/2007	03/31/2020
4		96 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075232	02/26/2003		

Labeler - Topco Associates LLC (006935977)

Revised: 11/2021

Topco Associates LLC