

**DAYTIME NIGHTTIME MULTI SYMPTOM COLD CHILDRENS- dextromethorphan hbr, guaifenesin, phenylephrine hcl, acetaminophen, diphenhydramine hcl, phenylephrine hcl  
Walgreens**

-----  
**Drug Facts**

**Active ingredients Daytime (in each 5 mL)**

Dextromethorphan HBr 5 mg

Guaifenesin 100 mg

Phenylephrine HCl 2.5 mg

**Active ingredients Night Time (in each 10 mL)**

**Acetaminophen 325 mg**

Diphenhydramine HCl 12.5 mg

Phenylephrine HCl 5 mg

**Purposes for Day Time**

Cough suppressant

Expectorant

Nasal decongestant

**Purposes for Night Time**

**Pain reliever/fever reducer**

Antihistamine/Cough suppressant

Nasal decongestant

**Uses**

Daytime

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - nasal congestion due to a cold

- stuffy nose

## Nighttime

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - sneezing
  - runny nose
  - nasal congestion
  - cough
- controls cough to help your child get to sleep
- temporarily reduces fever

## Warnings

### Nighttime only

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

#### DAYTIME

- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

#### NIGHTTIME

- with other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this

product.

- with any other drug containing diphenhydramine even one used on the skin
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- to make a child sleepy

### **Ask a doctor before use your if child has**

#### **DAYTIME**

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such occurs with asthma

#### **NIGHTTIME**

- liver disease
- heart disease
- high blood disease
- diabetes
- thyroid disease
- glaucoma
- a breathing problem such as chronic bronchitis
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if your child is**

#### **NIGHTTIME**

- taking sedatives or tranquilizers.
- taking the blood thinning drug warfarin

### **When using this product,**

#### **Daytime**

**do not use more than directed.**

#### **Nighttime**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

#### **DAYTIME**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

## NIGHTTIME

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

## Keep out of reach of children.

### Daytime

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Nighttime

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you not notice any signs or symptoms.

## Directions

### Daytime

- do not take more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor
- mL= milliliter

age	dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

### Nighttime

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see Overdose warning)**
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device

- mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

## **Other information**

### Daytime

- **each 5 mL contains:** sodium 5 mg
- store between 20-25°C (68°-77°F). Do not refrigerate.

### Nighttime

- **each 10 mL contains:** sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

## **Inactive ingredients**

### **Inactive ingredients for Day Time**

citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### **Inactive ingredients for Night Time**

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

## **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

Compare to the active ingredients in Children's Mucinex® Multi-Symptom Cold & Night Time Multi-Symptom Cold††

### **MULTI-SYMPTOM**

#### **Children's Cold**

DEXTROMETHORPAN HBr 5 mg /

COUGH SUPPRESSANT

GUAIFENESIN 100 mg /

EXPECTORANT

PHENYLEPHRINE HCl 2.5 mg /

## NASAL DECONGESTANT

Daytime

Alcohol Free

- Relief from chest congestion, cough & stuffy nose
- Breaks up mucus

AGES 4 - 11 YEARS

Berry flavor

FL OZ (mL)

children's

## **MULTI-SYMPTOM**

### **Children's Cold**

ACETAMINOPHEN 325 mg /

PAIN RELIEVER / FEVER REDUCER

DIPHENHYDRAMINE HCl 12.5 mg /

ANTIHISTAMINE / COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg /

NASAL DECONGESTANT

Nighttime

Alcohol Free

- Relief from stuffy nose, cough, fever, sore throat, runny nose & sneezing

AGES 6 - 11 YEARS

Berry flavor

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

††This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold and Children's Mucinex® Multi-Symptom Cold

DISTRIBUTED BY; WALGREEN CO.

DEERFIELD, IL 60015

walgreens.com

**Product Label**

### Nighttime Multi-Symptom Cold Drug Facts

**Drug Facts (continued)**

**When using this product:**

- do not use more than directed
- irritability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if:**

- irritation, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

**For adult use:**

- do not give more than directed (see Overdose warning)
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 days unless directed by a doctor

**Measure only with dosing cup provided.** Do not use any other dosing device. ■ mL = milliliter

**Children 6 to under 12 years of age:** 10 mL every 4 hours

**Children under 6 years of age:** do not use

**Other information**

- each 10 mL contains sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Daytime Multi-Symptom Cold Drug Facts

**Drug Facts**

**Active ingredients (in each 5 mL)**

Ingredient	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughing more productive
- temporarily relieves
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep
- nasal congestion due to a cold
- stuffy nose

**Warnings**

**Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI)** (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if your child has:**

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if:**

- irritation, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough lasts more than 7 days, comes back, or occurs with a fever, rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**In case of overdose, get medical help or contact a Poison Control Center 1-800-222-1222 right away.**

**Directions**

- do not take more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor
- mL = milliliter

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

**Drug Facts (continued)**

**Other information**

- each 5 mL contains sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?**  
Call 1-877-753-3888 Monday-Friday 9AM-5PM EST

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DOSAGE CUP PROVIDED**

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.**

**This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Night Time Multi-Symptom Cold and Children's Mucinex® Multi-Symptom Cold.**

DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015

**100% SATISFACTION GUARANTEED**

[www.walgreens.com](http://www.walgreens.com)  
©2021 Walgreen Co.

## DAY & NIGHT PACK

NDC 0363-4200-08

Compare to the active ingredients in Children's Mucinex® Multi-Symptom Cold & Night Time Multi-Symptom Cold\*

### MULTI-SYMPTOM Children's Cold

DEXTROMETHORPHAN HBr 5 mg / COUGH SUPPRESSANT  
GUAIFENESIN 100 mg / EXPECTORANT  
PHENYLEPHRINE HCl 2.5 mg / NASAL DECONGESTANT

**Daytime**

**Alcohol Free**

- Relief from chest congestion, cough & stuffy nose
- Breaks up mucus

### MULTI-SYMPTOM Children's Cold

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER  
DIPHENHYDRAMINE HCl 12.5 mg / ANTIHISTAMINE / COUGH SUPPRESSANT  
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

**Nighttime**

**Alcohol Free**

- Relief from stuffy nose, cough, fever, sore throat, runny nose & sneezing

**AGES 4 - 11 YEARS** Berry flavor      **AGES 6 - 11 YEARS** Berry flavor

**2 - 4 FL OZ (118 mL) BOTTLES / TOTAL 8 FL OZ (236 mL)**

ITEM 357645 W11042-0326-L

PLD-HS4A PLS-64A FDC0830

Lot No.:  
Exp. Date:

## WALGREENS Multi-Symptom Children's Cold Daytime Nighttime Berry Flavor

### Product label

**Drug Facts (continued)**

- do not give more than 5 days unless directed by a doctor
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- dose as follows or as directed by a doctor
- children 6 to under 12 years of age: 10 mL every 4 hours
- children under 6 years of age: do not use

**Other information**

- each 10 mL contains sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients:** citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Walgreens Pharmacist Recommended.  
Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
[www.walgreens.com](http://www.walgreens.com) ©2021 Walgreen Co.

PLD-6543A LB010842 ITEM 357645 W11042-0326-F W30RG0222-F

MULTI-SYMPTOM Children's Cold

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER  
DIPHENHYDRAMINE HCl 12.5 mg / ANTIHISTAMINE / COUGH SUPPRESSANT  
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

**Nighttime**

**Alcohol Free**

**AGES 6 - 11 YEARS**

**4 FL OZ (118 mL)**

Berry flavor

NDC 0363-4060-04

**Drug Facts**

**Active ingredients (in each 10 mL)**

Ingredient	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold and flu symptoms
- minor aches and pains
- headache
- sore throat
- sneezing
- runny nose
- nasal congestion
- cough
- controls cough to help your child get to sleep
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see Overdose warning)
- do not give more than 5 doses in any 24-hour period
- if needed, repeat dose every 4 hours while symptoms last

## WALGREENS Nighttime Multi-Symptom Children's Cold Berry Flavor

### Package label

**Drug Facts** (continued)

**Other information**

- each 5 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate
- Inactive ingredients** citric acid, disodium EDTA, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO.  
DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2021 Walgreen Co.

PLD-F544A LB010843 ITEM 357645 W11042-0326-F

REV-0623  
W3ORG0222-F



**Walgreens**

**MULTI-SYMPTOM Children's Cold**

DEXTROMETHORPHAN HBr 5 mg / COUGH SUPPRESSANT  
GUAIFENESIN 100 mg / EXPECTORANT  
PHENYLEPHRINE HCl 2.5 mg / NASAL DECONGESTANT

Daytime Alcohol Free

- Relief from chest congestion, cough & stuffy nose
- Breaks up mucus

AGES 4 - 11 YEARS

4 FL OZ (118 mL)

Berry flavor

NDC 0363-4070-04

**Drug Facts**

**Active ingredients (in each 5 mL)**

- Dextromethorphan HBr 5 mg.....Cough suppressant
- Guaifenesin 100 mg.....Expectorant
- Phenylephrine HCl 2.5 mg.....Nasal decongestant

**Purposes**

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep
  - nasal congestion due to a cold
  - stuffy nose

**Warnings**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- do not take more than 6 doses in any 24-hour period
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- keep dosing cup with product
- dose as follows or as directed by a doctor

Age	Dose
children 6 years to under 12 years	10 mL every 4 hours
children 4 years to under 6 years	5 mL every 4 hours
children under 4 years	do not use

**WALGREENS Daytime Multi-Symptom Children's Cold Berry Flavor**

**DAYTIME NIGHTTIME MULTI SYMPTOM COLD CHILDRENS**

dextromethorphan hbr, guaifenesin, phenylephrine hcl, acetaminophen, diphenhydramine hcl, phenylephrine hcl kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-4200
---------------------	----------------	---------------------------	---------------

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4200-08	1 in 1 KIT; Type 0: Not a Combination Product	12/31/2014	

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	118 mL
Part 2	1 BOTTLE, PLASTIC	118 mL

**Part 1 of 2**

**DAYTIME MULTI SYMPTOM COLD CHILDRENS**

dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

**Product Information**

<b>Item Code (Source)</b>	NDC:0363-4070
---------------------------	---------------

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4070-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2014	

## Part 2 of 2

### NIGHTTIME MULTI SYMPTOM COLD CHILDRENS

acetaminophen,diphenhydramine hcl,phenylephrine hcl liquid

#### Product Information

Item Code (Source) NDC:0363-4060

Route of Administration ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 [arb'U] in 10 mL

#### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
XANTHAN GUM (UNII: TTV12P4NEE)	

#### Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a		

4060-04	Combination Product		
---------	---------------------	--	--

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2014	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/31/2014	

**Labeler** - Walgreens (008965063)

Revised: 3/2026

Walgreens