

**ACETAMINOPHEN- acetaminophen tablet**  
**PD-Rx Pharmaceuticals, Inc.**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Acetaminophen 325mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning:**This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children.

Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:**acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if the user has**

liver disease

**Ask a doctor or pharmacist before use if the user is**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of the reach of children.**

**Overdose warning:**

In the case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"><li>• take 2 tablets, every 4 to 6 hours while symptoms last</li><li>• do not take more than 10 tablets in 24 hours, unless directed by a doctor</li><li>• do not use for more than 10 days unless directed by a doctor</li></ul>
children 6 to under 12 years	<ul style="list-style-type: none"><li>• take 1 tablet every 4 to 6 hours while symptoms last</li><li>• do not take more than 5 tablets in 24 hours</li><li>• do not use for more than 5 days unless directed by a doctor</li></ul>
children under 6 years	ask a doctor

**Other information**

- store between 20-25°C (68-77°F)
- TAMPER EVIDENT: DO NOT USE IF SEAL IS BROKEN OR MISSING FROM BOTTLE

### Inactive ingredients

povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid


### Questions or comments?

1-800-645-2158 (Mon-Fri 8 AM to 4 PM EST)


### Regular Strength

### Acetaminophen

325 mg

<b>Drug Facts</b> <b>Active Ingredient</b> (in each tablet) Acetaminophen 325mg		<b>Purpose</b> Pain reliever/fever reducer	
<b>USES</b> • temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps • temporarily reduces fever			
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if: • Adult takes more than 4,000 mg of acetaminophen in 24 hours • Child takes more than 5 doses in 24 hours, which is the maximum daily amount • Taken with other drugs containing acetaminophen • Adult has 3 or more alcoholic drinks every day while using this product.			
<b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.			
<b>Do not use</b> • with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product.			
 GTIN: 00372789267203 SNO: H22A09000004 EXP: 08/2024 LOT: H22A09		Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. <b>Stop use and ask a doctor if:</b> Pain gets worse or lasts more than 10 days in adults • Pain gets worse or lasts more than 5 days in children under 12 years • Fever gets worse or lasts more than 3 days • New symptoms occur • Redness or swelling is present. These could be signs of a serious condition. <b>If pregnant or breast-feeding,</b> ask a health professional before use. <b>Keep out of reach of children. Overdose warning:</b> Taking more than the recommended dose (overdose) may cause liver damage. In the case of accidental overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
<b>Directions:</b> • Do not take more than directed (see overdose warning).			
Adults and children 12 years and over	• take 2 tablets every 4 to 6 hours while symptoms last • do not take more than 10 tablets in 24 hours, unless directed by a doctor • do not use for more than 10 days unless directed by a doctor		
Children 6 to 11 years	• take 1 tablet every 4 to 6 hours while symptoms last • do not take more than 5 tablets in 24 hours • do not use for more than 5 days unless directed by a doctor		
Children under 6 years	Ask a doctor		
Other information: Store between 20-25 °C (68-77°F) Inactive ingredients Povidone, pregelatinized corn starch, sodium starch glycolate, stearic acid. Questions or comments? 1-800-645-2158			

NDC 72789-267-20



**ACETAMINOPHEN**  
**325 mg**

Marketed and Packaged By:  
 PD-Rx Pharmaceuticals, Inc  
 Oklahoma City, OK 73127  
 1-405-942-3040 v.8.19.0

**20 Tablets**  
 TAMPER EVIDENT: DO NOT USE IF  
 SEAL IS BROKEN OR MISSING FROM BOTTLE.

## ACETAMINOPHEN

acetaminophen tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72789-267(NDC:0536-1327)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	PH020
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72789-267-20	20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/01/2022	
2	NDC:72789-267-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/05/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/11/2021	

**Labeler** - PD-Rx Pharmaceuticals, Inc. (156893695)

**Registrant** - PD-Rx Pharmaceuticals, Inc. (156893695)

## Establishment

Name	Address	ID/FEI	Business Operations
PD-Rx Pharmaceuticals, Inc.		156893695	repack(72789-267)

Revised: 2/2025

PD-Rx Pharmaceuticals, Inc.