

**DAIRYLAND BRAND CHG TEAT DIP- chlorhexidine gluconate,
didecyldimethylammonium chloride liquid
Stearns Packaging Corporation**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

CHG Teat Dip

TEAT DIPPING is an important part of a MASTITIS CONTROL PLAN:

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

TEAT DIPPING IN COLD WEATHER:

When the temperature falls below 10 degrees F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outside areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

DIRECTIONS FOR USE:

- DO NOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder), IMMEDIATELY after every milking.

TAKE TIME

OBSERVE LABEL

DIRECTIONS

- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour used dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

FIRST-AID:

EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.

INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water or milk. DO NOT induce vomiting. Get medical attention immediately.

24 HOUR EMERGENCY CONTACT CHEM-TEL, 1-800-255-3924

DAIRYLAND BRAND

CHG TEAT DIP

A SANITIZING TEAT DIP NPE-Free Aid in Reducing the Spread of Organisms Which May Cause Mastitis.

ACTIVE INGREDIENTS:

Chlorhexidine Digluconate.....0.35%
Didecyl Dimethyl Ammonium Chloride.....0.20%

Excellent skin conditioning properties.

KEEP OUT OF THE REACH OF CHILDREN

See side panel for precautionary and First-Aid Statements.

STEARNS PACKAGING CORPORATION
4200 Sycamore Ave. (53714), PO Box 3216, Madison, WI 53704
(800) 655-5008 www.stearnspkg.com

NET CONTENTS: 1 U.S. FL. GALLON (3.78 LITERS)

DIRECTIONS FOR USE:

- DONOT DILUTE.
- Dip teats immediately after milking as a routine part of milking procedures.
- Dip to THOROUGHLY COVER at least the lower one-third of the teat (preferably to the base of the udder). IMMEDIATELY after every milking.
- Empty and clean dip cups after each milking, or during milking, if they become contaminated.
- Do not pour used dip back into original container.
- KEEP FROM FREEZING
- Keep container closed to prevent contamination.
- Store container in an upright position.

TEAT DIPPING IS AN IMPORTANT PART OF A MASTITIS CONTROL PLAN:

1. Dip teats immediately after milking.
2. Use good milking and environmental management procedures.
3. Use properly functioning milking equipment.
4. Treat every quarter of every cow at dry off.
5. Identify and treat or manage clinical cases promptly.
6. Cull chronically infected cows.

TEAT DIPPING IN COLD WEATHER:
When the temperature falls below 10°F, or when wind chill is significant, special precautions should be taken to avoid chapped and frozen teats.

1. Teats should be dry before turning cows out into cold weather.
2. When teats are dipped after milking, allow 30 seconds contact, and blot off any excess teat dip with a single service towel.
3. Warm the teat dip to reduce drying time.
4. Provide windbreaks in outdoor areas for cows.
5. Monitor fresh cows with swollen udders and teats since they are more susceptible to chapped and frozen teats.

DAIRYLAND BRAND

**CHG
TEAT DIP**

**A SANITIZING TEAT DIP
NPE-Free Aid in Reducing the Spread of
Organisms Which May Cause Mastitis**

ACTIVE INGREDIENTS:
Chlorhexidine Digluconate 0.35%
Didecyl Dimethyl Ammonium Chloride 0.20%

Excellent skin conditioning properties.

KEEP OUT OF THE REACH OF CHILDREN
See side panel for Precautionary and First-Aid Statements.

STEARNS PACKAGING CORPORATION
4200 Sycamore Ave, 53714, PO Box 3216, Madison, WI 53704
(800) 655-5008 • www.stearnspkg.com

NET CONTENTS: 1 U.S. GALLON (3.78 LITERS)

FIRST-AID:
EYES: If contact with eyes occurs, flush with plenty of cool water for 15 minutes. Consult a physician.
INTERNAL: May be harmful if swallowed. If ingested, drink large amounts of water. DO NOT induce vomiting. Get medical attention immediately.

PRIMEROS AUXILIOS:
OJOS: Si ocurre contacto con los ojos, lavar con abundante agua fría durante 15 minutos. Consulte a un médico.
INTERNO: Puede ser nocivo en caso de ingestión. Beba grandes cantidades de agua si se ingiere este producto. NO induzca el vómito. Obtenga atención médica inmediatamente.

24 HOUR EMERGENCY CONTACT
24 horas servicio telefónico de repuesta a emergencias:
1-800-255-3924

**#1205342
(ST-80)
REORDER NUMBER**

LOT NO.:
EXP. DATE:

INSTRUCCIONES POR EL USO:

- NO lo dilúas.
- Sumerja las pezones inmediatamente después del ordeño como parte rutinaria de procedimientos de ordeño.
- Sumerja para cubrir a fondo al menos el tercio inferior de la tetina (preferiblemente a la base de la teta), inmediatamente después de cada ordeño.
- Vacío y limpie las tazas sumergiendo después de cada ordeño, o durante el ordeño, si contaminan.
- No vierta las pezones utilizadas volver a la recipiente de origen.
- NO DEJE QUE SE CONGELÉ.
- Mantenga el recipiente bien cerrado para evitar la contaminación.
- Mantenga el recipiente en una posición vertical.

BARIO DE PEZONES ES UNA PARTE IMPORTANTE DE UN PLAN PARA CONTROL DE LA MASTITIS:

1. Sumerja los pezones inmediatamente después del ordeño.
2. Utilice buenas prácticas de gestión para el ordeño y medio ambiente.
3. Utilice equipo de ordeño que funcione correctamente.
4. Trate a cada cuarto de la ubre de cada vaca al dentro de fase de secado.
5. Identifique y trate o maneje casos clínicos con prontitud.
6. Entesque las vacas que están infectados crónicamente.

BARIO DE PEZONES EN TIEMPO FRÍO:
Cuando la temperatura cae por debajo de 10°F (-12°C), o cuando el factor de enfriamiento del viento es importante tome precauciones especiales para evitar las pezones agrietadas y congeladas.

1. Los pezones deben estar secos antes de soñar las vacas en el frío.
2. Cuando los pezones se sumergen después del ordeño, espere 30 segundos de contacto, y posteriormente seque el exceso de baño de pezones con una toalla desechable.
3. Caliente el baño de pezones para reducir el tiempo de secado.
4. Proporcione protección contra el viento en las áreas fuera para las vacas.
5. Vigile las vacas frescas con ubres hinchadas y pezones ya que son más susceptibles a los pezones agrietados y congelados.

**TAKE TIME
OBSERVE
LABEL DIRECTIONS**

DAIRYLAND BRAND CHG TEAT DIP

chlorhexidine gluconate, didecyl dimethyl ammonium chloride liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:60282-2080
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	3.5 g in 1 L
DIDECYLDIMONIUM CHLORIDE (UNII: JXN40O9Y9B) (DIDECYLDIMONIUM - UNII:Z7F472XQPA)	DIDECYLDIMONIUM CHLORIDE	2 g in 1 L

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60282-2080-1	3.78 L in 1 BOTTLE, PLASTIC		
2	NDC:60282-2080-2	18.93 L in 1 PAIL		
3	NDC:60282-2080-3	56.78 L in 1 DRUM		
4	NDC:60282-2080-4	208.2 L in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/02/1985	

Labeler - Stearns Packaging Corporation (006069256)

Registrant - Stearns Packaging Corporation (006069256)

Establishment

Name	Address	ID/FEI	Business Operations
Stearns Packaging Corporation		006069256	manufacture

Establishment

Name	Address	ID/FEI	Business Operations
Lab Express International		800875106	api manufacture

Revised: 12/2025

Stearns Packaging Corporation