## COLD AND FLU RELIEF NIGHTTIME- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled AmerisourceBergen (Good Neighbor Pharmacy) 46122

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Drug Facts**

#### Active ingredients (in each softgel)

#### Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

#### Purpose

#### Pain reliever/Fever Reducer

**Cough Suppressant** 

Antihistamine

#### Uses

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**: If sore throat is severe, persists more than 2 days, is accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor prompt

#### Do not use

• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor of pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

#### When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

#### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.%3

#### If pregnant or breast-feeding,

ask a health professional before use.

#### Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **Directions**

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

#### Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

#### **Inactive ingredients**

D&C yellow #10, edible white ink, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol

#### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

#### **Principal Display Panel**

\*\*Compare to VICKS® NYQUIL® LIQUICAPS® active ingredients

#### **Nighttime Liquid Caps**

Acetaminophen, Dextromethorphan HBr, Doxylamine succinate

#### Cold & Flu

Pain Reliever-FeverReducer

Cough Suppressant-Antihistamine

Relieves Cold and Flu-like Symptoms

So You Can Rest

Softgels

\*\*This product is not manufactured or distributed by the Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademark of the Procter & Gamble Company

### TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

#### KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Distributed by AmerisouceBergen

1300 Morris Drive, Chesterbrook, PA 19087

Questions or Concerns? Visit us at www.mygnp.com

#### **Package Label**

<sup>\*</sup>may contain this ingredient

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST Questions or comments?

> may contain this ingredient sorbitan\*, sorbitol

đejstju, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, Inactive ingredients D&C yellow #10, edible white ink, FD&C blue #1,

> ■ SAOID EXCESSIVE NEST ■ store between 15-30°C (59-86°F) Other information

label to ensure correct dosing ■ when using other Daytime or Nighttime products, carefully read each

- children under 12 years: do not use
- swallow whole; do not crush, chew, or dissolve
  - e ponus
- squitz sud children 12 years and over: take 2 softgels with water every
  - do not take more than 4 doses in 24 hours
  - do not take more than directed (see Overdose warning)

#### DIrections

children even it you do not notice any signs or symptoms. Center right away. Quick medical attention is critical for adults as well as for damage. In case of overdose, get medical help or contact a Poison Control Overdose warning: Taking more than the recommended dose can cause liver keep out of reach of children.

it pregnant or breast-feeding, ask a health professional before use.

pe sidus ot a serious condition.

- condu comes pack or occurs with rash or headache that lasts. These could
  - Lequess or swelling is present
     Inew symptoms occur
    - Tever gets worse or lasts more than 3 days
    - beju or cough gets worse or lasts more than 7 days

Stop use and ask a doctor if

**Drug Facts** (continued)

#### ■ pe careţnţ when driving a motor vehicle or operating machinery

- alcohol, sedatives, and tranquilizers may increase drowsiness
- wsrked growsiness may occur ■ SAOID SICOUOIIC DUUKS
  - excitability may occur, especially in children

#### When using this product do not exceed recommended dosage

- ESKING THE BIOOD THINNING DRUG WAITARIN ESKING SECRETIVES OF Transplicers Ask a doctor or pharmacist before use if you are
  - condu tust occurs with too much phiegm (mucus) суколіс рконсуірг ок втрпузета
- s preathing problem or chronic cough such as occurs with smoking, asthma,
  - trouble urinating due to an enlarged prostate gland

■ ðjøncoma Ask a doctor before use if you have 

liver disease

pharmacist before taking this product. not know if your prescription drug contains an MAOI, ask a doctor or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do (certain drugs for depression, psychiatric, or emotional conditions, or

- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) sak a doctor or pharmacist.
- nonprescription). If you are not sure whether a drug contains acetaminophen, ■ with any other drug containing acetaminophen (prescription or

s qoctor promptly.

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult Sore throat warming: If sore throat is severe, persists for more than 2 days, is It a skin reaction occurs, stop use and seek medical help right away.

may include: 

skin reddening 

blisters 

rash

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms 3 or more alcoholic drinks every day while using this product

with other drugs containing acetaminophen

occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours Liver warming: This product contains acetaminophen. Severe liver damage may Sbuiwem

Drug Facts (continued)

Doxylamine succinate 6.25 mg. **Antihistamine** Cough suppressant Dextromethorphan HBr 15 mg. ■ sore throat Pain reliever/fever reducer Active ingredients (in each softgel) SƏSN Purpose Drug Facts

 cough due to minor throat and bronchial irritation ■ LINUIN HOSE SUG SUGESING ■ µesqscµe ■ minor sches and pains femborarily relieves common cold and flu symptoms

nund racts (continued)

\*\*Compare to Vicks° NyQuil° LiquiCaps° active ingredients

NDC 46122-326-73



# Night Time Liquid Caps

acetaminophen, dextromethorphan HBr, doxylamine succinate

Cold & Flu

Pain Reliever-Fever Reducer Cough Suppressant-Antihistamine Relieves Cold and Flu-like Symptoms So You Can Rest



EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROXEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.



"This product is not manufactured or distributed by The Procter & Gamble Company, Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



PLD-D41R FC002972

Distributed By AmerisourceBergen 1300 Morris Drive, Chesterbrook, PA 19087 Questions or Concerns? Visit us at www.mygnp.com





### Good Neighbor Pharmacy Night Time Cold & Flu Liquid Caps

### COLD AND FLU RELIEF NIGHTTIME

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-326	
Route of Administration	ORAL			

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
ACETAMINO PHEN (UNII: 36209 ITL9D) (ACETAMINO PHEN - UNII: 36209 ITL9D)	ACETAMINOPHEN	325 mg			
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg			
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg			

Inactive Ingredients			
Ingredient Name	Strength		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
PO VIDO NE (UNII: FZ989 GH94E)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SORBITAN (UNII: 6O92ICV9RU)			
SORBITOL (UNII: 506T60A25R)			

Due dont Characteristics				
Product Characteristics				
Color	GREEN	Score	no score	
Shape	OVAL	Size	20 mm	
Flavor		Imprint Code	P30;94A;SCU1;215;P120	
Contains				

Packaging					
# Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>		
1 NDC:46122-326-73	16 in 1 CARTON	11/30/2015	12/31/2020		
1	$1\ \text{in}\ 1\ \text{BLISTER}$ PACK; Type 0: Not a Combination Product				
Marketing Information					
Marketing Catego	ory Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC MONOGRAPH FI	NAL part341	11/30/2015	12/31/2020		

Labeler - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 11/2019 AmerisourceBergen (Good Neighbor Pharmacy) 46122