

**ANTACID- aluminum hydroxide, magnesium hydroxide, simethicone suspension**  
**Shopko Stores Operating Co., LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Shopko Antacid Drug Facts**

**Active ingredients (in each 10 mL)**

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg

Magnesium hydroxide 800 mg

Simethicone 80 mg

**Purposes**

Antacid

Antigas

**Uses**

relieves

- heartburn
- acid indigestion
- sour stomach
- upset stomach due to these symptoms
- pressure and bloating commonly referred to as gas

**Warnings**

**Ask a doctor before use if you have**

- kidney disease
- a magnesium-restricted diet

**Ask a doctor or pharmacist before use if you are**

presently taking a prescription drug. Antacids may interact with certain prescription drugs.

**When using this product**

do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- shake well before use

- measure with dosing cup provided
- **adults and children 12 years and over:** 10 mL – 20 mL between meals, at bedtime or as directed by a doctor
- do not take more than 60 mL in any 24 hour period
- do not use the maximum dosage for more than 2 weeks
- **children under 12 years:** ask a doctor
- mL = milliliter

### **Other information**

- **each 10 mL contains:** magnesium 350 mg, potassium 20 mg and sodium 3 mg
- **does not meet USP requirements for preservative effectiveness**
- store at 20-25°C (68-77°F), do not freeze

### **Inactive ingredients**

butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

Compare To Maximum Strength Mylanta<sup>®</sup> active ingredients

MAXIMUM STRENGTH

ANTACID

Antacid/Antigas

Tough on heartburn

Works on contact

Classic Original Flavor

12 FL OZ (355 mL)

NDC 37012-340-40

\*Compare To  
Maximum Strength Mylanta®  
active ingredients



# MAXIMUM STRENGTH ANTACID

Antacid/Antigas

Tough on heartburn  
Works on contact  
Classic Original Flavor

12 FL OZ (355 mL)

: 34040 & C F3

**DO NOT USE IF  
PRINTED NECKBAND  
IS BROKEN OR  
MISSING**

\*This product is not  
manufactured or  
distributed by Infirst  
Healthcare Inc.,  
distributor of Maximum  
Strength Mylanta®.



4 00135 59018 4

**NO COPY AREA**

### Drug Facts

Active ingredients (in each 10 mL)	Purposes
Aluminum hydroxide (equiv. to dried gel, USP) 800 mg.....	Antacid
Magnesium hydroxide 800 mg.....	Antacid
Simethicone 80 mg.....	Antigas

**Uses** relieves ■ heartburn ■ acid indigestion ■ sour stomach  
■ upset stomach due to these symptoms  
■ pressure and bloating commonly referred to as gas

**Warnings**  
Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product do not exceed 60 mL in a 24-hour period, or use the maximum dosage for more than 2 weeks

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ shake well before use ■ measure with dosing cup provided  
■ adults and children 12 years and over: 10 mL – 20 mL between meals, at bedtime or as directed by a doctor  
■ do not take more than 60 mL in any 24 hour period  
■ do not use the maximum dosage for more than 2 weeks  
■ children under 12 years: ask a doctor ■ mL = milliliter

**Other information**  
■ each 10 mL contains: magnesium 350 mg, potassium 20 mg and sodium 3 mg  
■ does not meet USP requirements for preservative effectiveness  
■ store at 20-25°C (68-77°F), do not freeze

**Inactive ingredients** butylparaben, flavor, hypromellose, microcrystalline cellulose and carboxymethylcellulose sodium, potassium citrate, propylparaben, purified water, simethicone emulsion, sorbitol, sorbitol solution ▶

**NO COPY AREA**

**Drug Facts  
(continued)  
Questions or  
comments?  
1-800-719-9260**

Packaged By Perrigo®,  
Allegan, MI 49010  
1-800-719-9260  
©2017 Shopko Stores  
Operating Co., LLC  
www.shopko.com



By Refund or Exchange

: 34040 & C B3

# ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:370 12-340
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ALUMINUM HYDRO XIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDRO XIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDRO XIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	80 mg in 10 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)	
BUTYL PARABEN (UNII: 3QP1IU3FV8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	WHITE (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	LEMON	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:370 12-340-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/05/2013	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part331	06/05/2013	

**Labeler** - Shopko Stores Operating Co., LLC (023252638)

Revised: 12/2019

Shopko Stores Operating Co., LLC