

RITE AID PHARMACY ANTIBIOTIC WITH PAIN RELIEF- antibiotic with pain relief cream
RITE AID

Drug Facts

Active ingredient

Neomycin Sulfate (3.5mg)
Polymixin B Sulfate (10,000 units)
Pramoxine Hydrochloride (10 mg)

Purpose

Neomycin Sulfate	First Aid Antibiotic
Polymixin B Sulfate	First Aid Antibiotic
Pramoxine Hydrochloride	External Analgesic

Uses

first aid to prevent infection and for the temporary relief of pain or discomfort in minor:
cuts
scapes
burns

Warnings

For external use only.

Keep this and all drugs out of the reach of children

If swallowed, get medical help or contact Poison Control Center right away.

Directions

Adults and children 2 years of age and older:

- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other Information

- store at 20° to 25°C (68° to 77°F)

Inactive Ingredients

Cetareth-20, Cetearyl Alcohol, Glyceryl Stearate SE, Methylparaben, Mineral Oil, Phosphoric Acid, Propylene Glycol, Purified Water, White Petrolatum

Questions ?

Principal Primary Panel -Tube

Rite Aid Multi Antibiotic Cream with Pain Relief

NDC 11822-2163-1

- Neomycin Sulfate
- Polymixin B Sulfate
- Pramoxine HCL

First aid Antibiotic Pain Relieving Cream

NET WT 1.0 OZ (28g)

NDC 11822-2163-1

MAXIMUM STRENGTH

ANTIBIOTIC

+ PAIN RELIEF CREAM

NEOMYCIN SULFATE
POLYMYXIN B SULFATE • PRAMOXINE HCl

FIRST AID ANTIBIOTIC PAIN RELIEVING CREAM
 Helps prevent infection & relieves pain in minor cuts, scrapes & burns

 **NET WT 1.0 OZ (28 g)**

Active ingredients (in each gram) Neomycin 3.5mg, Polymyxin B 10,000 units, and Pramoxine hydrochloride 10 mg

Purposes First aid antibiotic and External analgesic

Uses first aid to help prevent infection in and temporarily relieves pain due to • minor cuts • scrapes • burns.

Warnings • For external use only. **Allergy alert:** • do not use if allergic to any of the ingredients.

Do not use • in or near the eyes • on large areas of the body. **Ask a doctor before use if you have** • deep or puncture wounds • animal bites • serious burns. **When using this product** • do not use longer than 1 week.

Stop use and ask a doctor if • condition lasts or gets worse • symptoms last for more than 7 days or clear up and come back within a few days • a rash or other allergic reaction occurs. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center immediately. 1-800-222-1222

Directions
 Adults and children 2 years and older: • clean affected area • apply a small amount (equal to the surface area of tip of finger) on area 1 to 3 times daily • may be covered with a sterile bandage. Children under 2 years: ask a doctor.

Other information • Store at controlled room temperature 68°-77°F (20°-25°C).

Questions? Call 1-800-719-9260 *This product is not manufactured or distributed by ©Johnson & Johnson Consumer Products Company - A Division of Johnson & Johnson Consumer Companies, Inc., owner of the registered trademark Neosporin® + Pain Relief

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 www.riteaid.com MADE IN THE USA OF U.S. & IMPORTED INGREDIENTS AND COMPONENTS 255968



Principal Primary Panel -Carton

Rite Aid Multi Antibiotic Cream with Pain Relief

NDC 11822-2163-1

- Neomycin Sulfate
- Polymixin B Sulfate
- Pramoxine HCL

First aid Antibiotic Pain Relieving Cream

NET WT 1.0 OZ (28g)

Drug Facts

Active Ingredients (in each gram) Purpose
Neomycin 3.5 mg First aid antibiotic
Polymyxin B 10,000 units First aid antibiotic
Pramoxine hydrochloride 10 mg External analgesic
Uses First aid to help prevent infection in and temporarily
relieve pain due to minor cuts • scrapes • burns

Warnings For external use only
Allergy alert: • do not use if allergic to any of the ingredients
Do not use • in or near the eyes • on large areas of the body
Ask a doctor before use if you have • deep or puncture
wounds • animal bites • serious burns

Directions
Apply a small amount (equal to the surface area of lip
of finger) on area 1 to 3 times daily.

Other Information
Store at controlled room temperature 68°-77° (20°-25°C).

Drug Facts

Active Ingredients Celebreth-20, Celestrol Alcohol,
Glyceryl Stearate SE, Methylparaben, Mineral Oil, Phosphoric
Acid, Propylene Glycol, Purified Water, White Petrolatum

Questions? Call 1-800-719-8280

Helps prevent infection &
relieves pain in minor
cuts, scrapes & burns

LOT
EXP

DISTRIBUTED BY:
J&J PHARMACEUTICALS, INC.
CINCINNATI, OH 45228
www.jandj.com
MADE IN U.S.A. OF U.S.
INGREDIENTS AND COMPONENTS

SATISFACTION
GUARANTEE
If you're not satisfied, we'll
refund your money.



NDC 11822-2163-1

MAXIMUM STRENGTH
ANTIBIOTIC + PAIN
RELIEF CREAM
POLYMYXIN B SULFATE PRAMOXINE HCl
FIRST AID ANTIBIOTIC
PAIN RELIEVING CREAM

MAXIMUM STRENGTH
**ANTIBIOTIC +
PAIN RELIEF
CREAM**

NEOMYCIN SULFATE
POLYMYXIN B SULFATE
PRAMOXINE HCl

FIRST AID ANTIBIOTIC
PAIN RELIEVING CREAM



NET WT 1.0 OZ (28 g)

*Compare to the active ingredients
of Neosporin® + Pain Relief



RITE AID PHARMACY ANTIBIOTIC WITH PAIN RELIEF

antibiotic with pain relief cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-2163
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP'U] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g

PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)		PRAMOXINE HYDROCHLORIDE	10 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)				
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)				
GLYCERYL STEARATE SE (UNII: FCZ5MH785I)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
MINERAL OIL (UNII: T5L8T28FGP)				
PHOSPHORIC ACID (UNII: E4GA8884NN)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
WHITE PETROLATUM (UNII: B6E5W8RQJ4)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-2163-1	1 in 1 CARTON	10/01/2022	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M004	10/01/2022	

Labeler - RITE AID (014578892)

Revised: 11/2024

RITE AID