

**ACID REDUCER MAXIMUM STRENGTH- famotidine tablet, film coated**  
**Rite Aid Corporation**

-----  
**Rite Aid Corporation Acid Reducer Drug Facts**

**Active ingredient (in each tablet)**

Famotidine 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture and light

**Inactive ingredients**

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

SEE NEW WARNINGS

Compare to the active ingredient of Maximum Strength Pepcid® AC

ACID REDUCER

Famotidine Tablets, 20 mg

acid reducer

MAXIMUM STRENGTH

Just one tablet prevents & relieves heartburn due to acid indigestion

ACTUAL SIZE

25 TABLETS



\* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., as produced by Maximum Strength Pepcid® AC.

GLUTEN FREE

**Drug Facts (continued)**  
**Questions or comments? 1-800-719-9260**

**Drug Facts (continued)**

**Active ingredient**  
 (in each tablet) Famotidine 20 mg..... Acid reducers

**Purpose**  
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**  
 your heartburn continues or worsens  
 you need to take this product for more than 14 days  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away. (1-800-222-1222)

**Directions**  
 adults and children 12 years and over: swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn  
 Do not chew.  
 to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn  
 do not use more than 2 tablets in 24 hours  
 children under 12 years: ask a doctor

**Other information**  
 read the directions and warnings before use  
 keep the carton. It contains important information.  
 store at 20°-25°C (68°-77°F)  
 protect from moisture

**Inactivating ingredients**  
 silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

CONVENIENT RECLOSING TAB

OPEN OTHER END

© 2009 RITE AID

19447 83 C6

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

**DO NOT USE IF PRINTED BLISTER UNIT IS BROKEN OR TORN**

DISTRIBUTED BY: RITE AID  
 30 HUNTER LANE,  
 CAMP HILL, PA 17011  
 www.riteaid.com

**SATISFACTION GUARANTEE**  
 If you're not satisfied, we'll happily refund your money.

SEE NEW WARNINGS

NDC 11822-0194-5

Compare to the active ingredient of Maximum Strength Pepcid® AC\*

# ACID REDUCER

Famotidine Tablets, 20 mg

acid reducer

**MAXIMUM STRENGTH**

Just one tablet prevents & relieves heartburn due to acid indigestion

ACTUAL SIZE





25  
TABLETS

## ACID REDUCER MAXIMUM STRENGTH

famotidine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0194
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FAMOTIDINE</b> (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	L194
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0194-3	1 in 1 CARTON	09/29/2006	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11822-0194-5	25 in 1 CARTON	09/29/2006	

2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11822-0194-7	2 in 1 CARTON	09/29/2006	
3		85 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:11822-0194-2	30 in 1 CARTON	04/13/2008	07/29/2015
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	09/29/2006	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 7/2021

Rite Aid Corporation