

GLYCERIN- glycerin suppository
Rising Pharma Holdings, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Glycerin Suppositories 2 g

Drug Facts

Active ingredient (in each suppository)

Glycerin USP, 2 grams

Purpose

Laxative

Uses

- for relief of occasional constipation
- this product usually produces bowel movement in 1/4 to 1 hour

Warnings

For rectal use only

Do not use for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks

When using this product it may cause rectal discomfort or a burning sensation

Stop use and ask a doctor if you have

- rectal bleeding
- no bowel movement after using this product

These may indicate a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 6 years of age and over : 1 suppository once daily
- children under 6 years : do not use
- insert 1 suppository well up into rectum and retain for 15 minutes
- it need not melt to produce laxative action

Other information

keep away from excessive heat.

Inactive ingredients

purified water, sodium stearate

Questions or Comments?

Call **1-844-474-7464**

TAMPER EVIDENT: DO NOT USE IF SEALED WRAPPER AROUND SUPPOSITORIES IS BROKEN OR MISSING

Distributed by:

Rising Pharma Holdings, Inc.
East Brunswick, NJ 08816

Made in India

CODE NO.: MH/DRUGS/25-KD/724



Issued: 08/2023


NDC 57237-325-52 (5 blisters of 5 suppositories)

GLYCERIN SUPPOSITORIES 2 g

LAXATIVE

For fast, gentle relief of occasional constipation

		LOT: XXXXXXXX EXP: YYYY/MM								
		Unvarnished area for 2D Code								
<table border="1"><tr><td>Drug Facts Active ingredient (in each suppository) Purpose Glycerin USP, 2 grams Laxative</td><td>Drug Facts (continued) Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.</td></tr><tr><td>Uses ■ for relief of occasional constipation ■ this product usually produces bowel movement in 1/4 to 1 hour</td><td>Directions ■ adults and children 6 years of age and over : 1 suppository once daily ■ children under 6 years : do not use ■ insert 1 suppository well up into the rectum and retain for 15 minutes ■ it need not melt to produce laxative action</td></tr><tr><td>Warnings For rectal use only Do not use for longer than 1 week unless directed by a doctor Ask a doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits lasting more than 2 weeks When using this product it may cause rectal discomfort or a burning sensation Stop use and ask a doctor if you have ■ rectal bleeding ■ no bowel movement after using this product These may indicate a serious condition,</td><td>Other information Keep away from excessive heat. Inactive ingredients purified water, sodium stearate</td></tr><tr><td colspan="2" style="text-align: right;">Questions or Comments? Call 1-844-474-7464</td></tr></table>			Drug Facts Active ingredient (in each suppository) Purpose Glycerin USP, 2 grams Laxative	Drug Facts (continued) Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	Uses ■ for relief of occasional constipation ■ this product usually produces bowel movement in 1/4 to 1 hour	Directions ■ adults and children 6 years of age and over : 1 suppository once daily ■ children under 6 years : do not use ■ insert 1 suppository well up into the rectum and retain for 15 minutes ■ it need not melt to produce laxative action	Warnings For rectal use only Do not use for longer than 1 week unless directed by a doctor Ask a doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits lasting more than 2 weeks When using this product it may cause rectal discomfort or a burning sensation Stop use and ask a doctor if you have ■ rectal bleeding ■ no bowel movement after using this product These may indicate a serious condition,	Other information Keep away from excessive heat. Inactive ingredients purified water, sodium stearate	Questions or Comments? Call 1-844-474-7464	
Drug Facts Active ingredient (in each suppository) Purpose Glycerin USP, 2 grams Laxative	Drug Facts (continued) Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.									
Uses ■ for relief of occasional constipation ■ this product usually produces bowel movement in 1/4 to 1 hour	Directions ■ adults and children 6 years of age and over : 1 suppository once daily ■ children under 6 years : do not use ■ insert 1 suppository well up into the rectum and retain for 15 minutes ■ it need not melt to produce laxative action									
Warnings For rectal use only Do not use for longer than 1 week unless directed by a doctor Ask a doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits lasting more than 2 weeks When using this product it may cause rectal discomfort or a burning sensation Stop use and ask a doctor if you have ■ rectal bleeding ■ no bowel movement after using this product These may indicate a serious condition,	Other information Keep away from excessive heat. Inactive ingredients purified water, sodium stearate									
Questions or Comments? Call 1-844-474-7464										
GLYCERIN SUPPOSITORIES 2 g LAXATIVE		TAMPER EVIDENT: DO NOT USE IF SEALED WRAPPER AROUND SUPPOSITORIES IS BROKEN OR MISSING								



NDC 57237-325-51 (5 suppositories in 1 blister pack)

Front Foil



**GLYCERIN
SUPPOSITORIES 2 g**

For rectal use only



NDC 57237-325-51



**GLYCERIN
SUPPOSITORIES 2 g**

For rectal use only



NDC 57237-325-51

Code No.:
MH/DRUGS/25-KD/724



**GLYCERIN
SUPPOSITORIES 2 g**

For rectal use only



NDC 57237-325-51

A
12004708

Back Foil

Dist. By: Rising Pharma Holdings, Inc.

Keep away from excessive heat.

Dist. By: Rising Pharma Holdings, Inc.

Keep away from excessive heat.

Dist. By: Rising Pharma Holdings, Inc.

Keep away from excessive heat.

A
12004707

GLYCERIN

glycerin suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57237-325
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	2 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

SODIUM STEARATE (UNII: QU7E2XA9TG)

Product Characteristics

Color	white (clear)	Score	
Shape	BULLET	Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57237-325-21	2 in 1 CARTON	08/24/2023	
1	NDC:57237-325-16	6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:57237-325-52	5 in 1 CARTON	08/24/2023	
2	NDC:57237-325-51	5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	08/24/2023	

Labeler - Rising Pharma Holdings, Inc. (116880195)

Establishment

Name	Address	ID/FEI	Business Operations
Bliss GVS Pharma Limited		675630493	manufacture(57237-325) , analysis(57237-325) , pack(57237-325)

Revised: 9/2023

Rising Pharma Holdings, Inc.