

**FEXOFENADINE- fexofenadine tablet, film coated**  
**Wockhardt Limited**

-----  
**Fexofenadine Tablets**

**Active ingredient (in each tablet)**

**For 30 mg:**

Fexofenadine HCl 30 mg

**For 60 mg:**

Fexofenadine HCl 60 mg

**For 180 mg:**

Fexofenadine HCl 180 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use**

if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have**

kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if**

an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

#### For 30 mg:

adults and children 12 years of age and over	take two 30 mg tablets with water every 12 hours; do not take more than 4 tablets in 24 hours
children 6 to under 12 years of age	take one 30 mg tablet with water every 12 hours; do not take more than 2 tablets in 24 hours
children under 6 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### For 60 mg:

adults and children 12 years of age and over	take one 60 mg tablet with water every 12 hours; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### For 180 mg:

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn
- safety sealed: do not use if carton is open or if individual blister unit is torn or open
- store between 20° to 25°C (68° to 77°F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

### Questions or comments?

Call toll-free **1-800-346-6854**

### PACKAGE LABEL PRINCIPAL DISPLAY PANEL

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-744-08

**Children's  
Fexofenadine HCl Tablet, USP**

**30 mg** 12 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR



Children's

**Fexofenadine HCl Tablets, USP**

**30 mg**

**12HOUR**

**ALLERGY**

Ages 6 Years and Older

**Drug Facts**

**Active ingredient (in each tablet)**

Fexofenadine HCl 30 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 12 years of age and over	take two 30 mg tablets with water every 12 hours; do not take more than 4 tablets in 24 hours
children 6 to under 12 years of age	take one 30 mg tablet with water every 12 hours; do not take more than 2 tablets in 24 hours
children under 6 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

**Other information**

- safety sealed: do not use if carton is open or if individual blister unit is torn or open
- store at 20° to 25°C (68° to 77°F)
- protect from excessive moisture

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

**Questions or comments?**

Call toll-free 1-800-346-6854



Non Varnish Area

Lot:  
Exp.:

Children's

**Fexofenadine HCl Tablets, USP**

**30 mg**

**12HOUR**

**ALLERGY**

Ages 6 Years and Older

DO NOT USE IF CARTON IS OPEN OR IF INDIVIDUAL BLISTER UNIT IS TORN OR OPEN

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-744-09

**Children's**

**Fexofenadine HCl Tablets, USP**

**30 mg/antihistamine**

**ALLERGY**

**12HOUR**

Manufactured by: Wockhardt Limited, Mumbai, India. Phone: +91 22 25112111  
Distributed by: Wockhardt USA LLC, 20 Waterview Blvd., Parsippany, NJ 07054, USA.

## Indoor and Outdoor Allergies

**Ages 6 Years and Older**

**RELIEF OF:**

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**6 Tablets 30 mg each**

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-982-08

**Fexofenadine HCl Tablet, USP**

**60 mg**

**12 Hour**

**ALLERGY**

Distributed by: Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR



TEAR

TEAR



# Fexofenadine HCl Tablets, USP

60 mg

**ALLERGY**

**12HOUR**

### Drug Facts

#### Active ingredient (in each tablet)

Fexofenadine HCl 60 mg .....Antihistamine

#### Purpose

#### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

#### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

adults and children 12 years of age and over	take one 60 mg tablet with water every 12 hours; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

#### Other information

- safety sealed: do not use if carton is open or if individual blister unit is torn or open
- store at 20° to 25°C (68° to 77°F)
- protect from excessive moisture

#### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

#### Questions or comments?

Call toll-free 1-800-346-6854



**Non Varnish Area**

Lot:  
Exp.:

# Fexofenadine HCl Tablets, USP

60 mg

**ALLERGY**

**12HOUR**

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-982-09

# Fexofenadine HCl Tablets, USP

**60 mg/antihistamine**

**ALLERGY**

**12HOUR**

**Indoor and Outdoor Allergies**

**RELIEF OF:**

- ✓ Sneezing
- ✓ Runny Nose

DO NOT USE IF CARTON IS OPEN OR IF  
INDIVIDUAL BLISTER UNIT IS TORN OR OPEN

Manufactured by  
Wockhardt  
Mumbai, Inc  
Iss. 151211

red by: **Distributed by:**  
 Limited, Wockhardt USA LLC,  
 dia, 20 Waterview Blvd,  
 Parsippany, NJ 07054  
 USA

- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**12 Tablets 60 mg each**

<p>ORIGINAL PRESCRIPTION STRENGTH NON-DROWSY NDC-64679-987-12</p> <p><b>Fexofenadine HCl Tablets, USP</b></p> <p><b>180 mg/antihistamine</b></p> <p><b>ALLERGY</b> Indoor and Outdoor Allergies</p> <p><b>24HOUR</b></p> <p><b>30 Tablets 180 mg each</b></p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS MISSING OR TORN</p>	<p><b>Active ingredient (in each tablet)</b> Fexofenadine HCl 180 mg.....Antihistamine</p> <p><b>Purpose</b> temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat</p> <p><b>Warnings Do not use</b> if you ever had an allergic reaction to this product or any of its ingredients. <b>Ask a doctor before use if you have kidney disease.</b> Your doctor should determine if you need a different dose. <b>When using this product</b> ■ do not take more than directed. ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) <b>Stop use and ask a doctor if</b> an allergic reaction to this product occurs. Seek medical help right away. <b>If pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center ►</p>	<p>right away.</p> <p><b>Directions</b> ■ <b>adults and children 12 years of age and over</b> take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ <b>children under 12 years of age</b> do not use ■ <b>adults 65 years of age and older</b> ask a doctor ■ <b>consumers with kidney disease</b> ask a doctor <b>Other information</b> ■ safety sealed: do not use if carton is open or if inner foil seal imprinted with "sealed for your protection" is missing or torn ■ store at 20° to 25°C (68° to 77°) ■ protect from excessive moisture</p> <p><b>Inactive ingredients</b> colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide</p> <p><b>Questions or comments?</b> Call toll free 1-800-346-6854</p>	<p>5 64679 98712 4</p> <p><b>Manufactured by:</b> Wockhardt Limited Mumbai, India.</p> <p><b>Distributed by:</b> Wockhardt USA LLC, 20 Waterview Blvd, Parsippany, NJ 07054, USA.</p> <p>Iss. 151211</p> <p>Lot: Exp.:</p> <p><b>Non Varnish Area</b></p>

ORIGINAL, PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-987-12

# Fexofenadine HCl Tablets, USP

180 mg/antihistamine

**ALLERGY**

Indoor and Outdoor  
Allergies

**24-HOUR**

**RELIEF OF:**

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**30 Tablets 180 mg each**

**Safety Sealed: Do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn.**

## Drug Facts

**Active ingredient (in each tablet) Purpose**  
Fexofenadine HCl 180 mg.....Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▲

## Drug Facts (continued)

### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn
- store at 20° to 25° C (68° to 77° F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

### Questions or comments?

Call 1-800-346-6854

# Fexofenadine HCl Tablets, USP

**ALLERGY**

**1 BOTTLE  
INSIDE**

**Manufactured by:**  
Wockhardt Limited,  
Mumbai, India.

**Distributed by:**  
Wockhardt USA LLC,  
20 Waterview Blvd.  
Parsippany, NJ 07054  
USA.

Iss:151211



N 3 64679 98712 4

LOT:

EXP:



DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH  
"SEALED FOR YOUR PROTECTION" IS MISSING OR TORN

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC-64679-987-13

# Fexofenadine HCl Tablets, USP

180 mg/antihistamine

## ALLERGY

Indoor and Outdoor Allergies

24HOUR



45 Tablets 180 mg each

### Active ingredient (in each tablet)

Fexofenadine HCl 180 mg.....Antihistamine

### Purpose

**Uses** temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat

**Warnings Do not use** if you ever had an allergic reaction to this product or any of its ingredients. **Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose. **When using this product** ■ do not take more than directed. ■ do not take at the same time as aluminum or magnesium antacids ■ do not take with fruit juices (see Directions) **Stop use and ask a doctor** if an allergic reaction to this product occurs. Seek medical help right away. **If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center ►

right away.

**Directions** ■ adults and children 12 years of age and over take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours ■ children under 12 years of age do not use ■ adults 65 years of age and older ask a doctor ■ consumers with kidney disease ask a doctor **Other information** ■ safety sealed: do not use if carton is open or if inner foil seal imprinted with "sealed for your protection" is missing or torn ■ store at 20° to 25°C (68° to 77°) ■ protect from excessive moisture

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide



Manufactured by:  
Wockhardt Limited  
Mumbai, India.

Distributed by:  
Wockhardt USA, LLC  
20 Valerius Blvd  
Parsippany, NJ 07054, USA.

Iss. 151211

Lot:  
Exp.:  
**Non Varnish Area**

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-987-13

# Fexofenadine HCl Tablets, USP

180 mg/antihistamine

**ALLERGY**

Indoor and Outdoor  
Allergies

**24HOUR**

**RELIEF OF:**

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**45 Tablets 180 mg each**

**Safety Sealed: Do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn.**

## Drug Facts

**Active ingredient (in each tablet)** Purpose  
Fexofenadine HCl 180 mg.....Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▲

## Drug Facts (continued)

### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn
- store at 20° to 25° C (68° to 77° F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

### Questions or comments?

Call 1-800-346-6854

# Fexofenadine HCl Tablets, USP

**ALLERGY**

**1 BOTTLE  
INSIDE**

**Manufactured by:**  
Wockhardt Limited,  
Mumbai, India.

**Distributed by:**  
Wockhardt USA LLC,  
20 Waterview Blvd,  
Parsippany, NJ 07054  
USA.

Iss:151211



LOT:

EXP:



**Safety Sealed:** Do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn.

# Fexofenadine HCl Tablets, USP

**ALLERGY**

**2 BOTTLES  
INSIDE**

**45 tablets in each Bottle**



LOT:

EXP:

## Drug Facts

**Active ingredient (in each tablet)** Purpose  
Fexofenadine HCl 180 mg.....Antihistamine

### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.  
**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▲

## Drug Facts (continued)

### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

### Other information

- safety sealed: do not use if carton is open or if inner seal imprinted with "Sealed for Your Protection" is missing or torn
- store at 20° to 25° C (68° to 77°F)
- protect from excessive moisture

### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

### Questions or comments?

Call 1-800-346-6854

# Fexofenadine HCl Tablets, USP

**ALLERGY**

**2 BOTTLES  
INSIDE**

**45 tablets in each Bottle**

**Manufactured by:**  
Wockhardt Limited,  
Mumbai, India.

**Distributed by:**  
Wockhardt USA LLC,  
20 Waterview Blvd.  
Parsippany, NJ 07054  
USA.

Iss:151211

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-987-14

# **Fexofenadine HCl** **Tablets, USP**

**180 mg/antihistamine**

**ALLERGY**

**Indoor and Outdoor Allergies**

**24HOUR**

**RELIEF OF:**

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**WOCKHARDT**

**90 Tablets 180 mg each**

NDC 64679-987-09

**Fexofenadine HCl Tablet, USP**

**180 mg** 24 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-987-09

**Fexofenadine HCl Tablet, USP**

**180 mg** 24 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-987-09

**Fexofenadine HCl Tablet, USP**

**180 mg** 24 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-987-09

**Fexofenadine HCl Tablet, USP**

**180 mg** 24 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR

NDC 64679-987-09

**Fexofenadine HCl Tablet, USP**

**180 mg** 24 Hour

**ALLERGY**

**Distributed by:** Wockhardt USA LLC.  
Parsippany, NJ 07054, USA.

Lot : Exp.:

TEAR



TEAR

TEAR



Fexofenadine HCl Tablets, USP

180 mg

24HOUR

ALLERGY

**Drug Facts**

**Active ingredient (in each tablet)**

Fexofenadine HCl 180 mg

**Purpose**

Antihistamine

**Uses**

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- Itching of the nose or throat

**Warnings**

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**if pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

**Other information**

- safety sealed: do not use if carton is open or if individual blister unit is torn or open
- store at 20° to 25°C (68° to 77°F)
- protect from excessive moisture

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

**Questions or comments?**

Call toll-free 1-800-346-6854



Non Varnish Area

Lot:  
Exp.:

Fexofenadine HCl Tablets, USP

180 mg

24HOUR

ALLERGY

DO NOT USE IF CARTON IS OPEN OR IF  
INDIVIDUAL BLISTER UNIT IS TORN OR OPEN

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-987-10

Fexofenadine HCl Tablets, USP

180 mg/antihistamine

ALLERGY

24HOUR

## Indoor and Outdoor Allergies

### RELIEF OF:

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**5 Tablets 180 mg each**

**Manufactured by:** Wockhardt Limited,  
Mumbai, India. s. 151211

**Distributed by:** Wockhardt USA LLC,  
20 Waterview Blvd., Parsippany,  
NJ 07054, USA.



# Fexofenadine HCl Tablets, USP

180 mg

ALLERGY

24HOUR

### Drug Facts

#### Active ingredient (in each tablet)

Fexofenadine HCl 180 mg

#### Purpose

Antihistamine

#### Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

#### Warnings

**Do not use** if you have ever had an allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** kidney disease. Your doctor should determine if you need a different dose.

#### When using this product

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)

**Stop use and ask a doctor if** an allergic reaction to this product occurs. Seek medical help right away.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

adults and children 12 years of age and over	take one 180 mg tablet with water once a day; do not take more than 1 tablet in 24 hours
--	---

children under 12 years of age	do not use
--------------------------------	------------

adults 65 years of age and older	ask a doctor
----------------------------------	--------------

consumers with kidney disease	ask a doctor
-------------------------------	--------------

#### Other information

- safety sealed: do not use if carton is open or if individual blister unit is torn or open
- store at 20° to 25°C (68° to 77°F)
- protect from excessive moisture

#### Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone K-30, talc, titanium dioxide

#### Questions or comments?

Call toll-free 1-800-346-6854



Non Varnish Area

Lot:  
Exp.:

# Fexofenadine HCl Tablets, USP

180 mg

ALLERGY

24HOUR

ORIGINAL PRESCRIPTION STRENGTH  
NON-DROWSY

NDC 64679-987-11

# Fexofenadine HCl Tablets, USP

180 mg/antihistamine

ALLERGY

24HOUR

Indoor and Outdoor Allergies

RELIEF OF:

DO NOT USE IF CARTON IS OPEN OR IF  
INDIVIDUAL BLISTER UNIT IS TORN OR OPEN

Manufactured by  
Wockhardt  
Mumbai, Inc  
Iss. 151211

red by: **Distributed by:**  
 Limited, Wockhardt USA LLC,  
 .lla, 20 Waterview Blvd.  
 Parsippany, NJ 07054  
 USA.

- ✓ Sneezing
- ✓ Runny Nose
- ✓ Itchy, Watery Eyes
- ✓ Itchy Nose or Throat



**15 Tablets 180 mg each**

## FEXOFENADINE

fexofenadine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55648-744
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXO FENADINE HYDRO CHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	30 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND (round)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	W;30
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55648-744-09	1 in 1 CARTON		
1	NDC:55648-744-08	6 in 1 BLISTER PACK		

  

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079112	02/08/2012	

**FEXOFENADINE**  
fexofenadine tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55648-982
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
FEXO FENADINE HYDRO CHLORIDE (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg

Inactive Ingredients	
Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	WHITE	Score	no score
Shape	CAPSULE (Capsule)	Size	11mm
Flavor		Imprint Code	W;982
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:55648-982-09	2 in 1 CARTON	
1	NDC:55648-982-08	6 in 1 BLISTER PACK	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079112	02/08/2012	

## FEXOFENADINE

fexofenadine tablet, film coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55648-987
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FEXO FENADINE HYDRO CHLORIDE (UNII: 2S068B75ZU) (FEXO FENADINE - UNII:E6582LOH6V)	FEXO FENADINE HYDROCHLORIDE	180 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELOLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	WHITE	Score	no score
Shape	CAPSULE (Capsule)	Size	17mm
Flavor		Imprint Code	W;987
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55648-987-12	1 in 1 CARTON		

1		30 in 1 BOTTLE		
2	NDC:55648-987-13	1 in 1 CARTON		
2		45 in 1 BOTTLE		
3	NDC:55648-987-14	2 in 1 CARTON		
3		45 in 1 BOTTLE		
4	NDC:55648-987-10	1 in 1 CARTON		
4	NDC:55648-987-09	5 in 1 BLISTER PACK		
5	NDC:55648-987-11	3 in 1 CARTON		
5	NDC:55648-987-09	5 in 1 BLISTER PACK		
6	NDC:55648-987-05	2000 in 1 POUCH		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079112	02/08/2012	

**Labeler** - Wockhardt Limited (650069115)

**Registrant** - Wockhardt Limited (650069115)

## Establishment

Name	Address	ID/FEI	Business Operations
Wockhardt Limited		676257570	Manufacture

Revised: 5/2012

Wockhardt Limited