

NICOTINE TRANSDERMAL SYSTEM STEP 1- nicotine patch, extended release

NICOTINE TRANSDERMAL SYSTEM STEP 2- nicotine patch, extended release

NICOTINE TRANSDERMAL SYSTEM STEP 3- nicotine patch, extended release

CARDINAL HEALTH 110, LLC. DBA LEADER

Drug Facts

Active ingredient Step 1 (in each patch)

Nicotine, 21 mg delivered over 24 hours

Active ingredient Step 2 (in each patch)

Nicotine, 14 mg delivered over 24 hours

Active ingredient Step 3 (in each patch)

Nicotine, 7 mg delivered over 24 hours

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product

- if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if

- skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash
- irregular heartbeat or palpitations occur.
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat.
- you have symptoms of an allergic reaction (such as difficulty breathing or rash).

Keep out of reach of children and pets.

Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away(1-800-222-1222). Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke **more than 10 cigarettes per day**, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1	STEP 2	STEP 3
Use one 21 mg patch/day	Use one 14 mg patch/day	Use one 7 mg patch/day

- if you smoke **10 or less cigarettes per day**, start with **Step 2** for 6 weeks, then **Step 3** for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning

- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- It is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information

store at 20-25°C (68-77°F)

Inactive ingredients

acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

Questions or comments?

Call **1-800-585-8682** Weekdays (8am-8pm EST)

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use one patch daily according to directions.
3. It is important to complete treatment.
4. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.
5. Use patch with a behavioral support program, such as the one described in the enclosed booklet.

SPL UNCLASSIFIED SECTION

For your family's protection, patches are supplied in child-resistant pouches. **Do not use if individual pouch is open or torn.**

- Not for sale to persons under 18 years of age.
- Proof of age required.
- Not for sale in vending machines or from any source where proof of age cannot be verified.

See self-help guide inside.

DISTRIBUTED BY

CARDINAL HEALTH

DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313 Essential to Care™ since 1979

*Habitrol® is a registered trademark of Dr. Reddy's Laboratories SA.

©2026 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

All LEADER™ Brand Products Have A 100% Money Back Guarantee

Return to place of purchase if not satisfied.

CIN 6107585

REV. 02/26

Principal Display Panel

70000-0762-1

Nicotine Transdermal System Patch

21 mg delivered over 24 hours

STOP SMOKING AID

21 mg

STEP 1

Includes: **Behavior Support Program with SELF-HELP GUIDE**

If You Smoke **MORE** than 10 Cigarettes per Day: Start with Step 1.

If You Smoke 10 or **LESS** Cigarettes per Day: Start with Step 2.

7 PATCHES (1 WEEK KIT)

The full treatment course for STEP 1 is 28 patches (4 weeks); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

COMPARE TO HABITROL® NICOTINE TRANSDERMAL SYSTEM PATCH active ingredient*



Principal Display Panel

NDC 70000-0761-1

Nicotine Transdermal System Patch

14 mg delivered over 24 hours

STOP SMOKING AID

14 mg

STEP 2

Includes: Behavior Support Program with SELF-HELP GUIDE


If You Smoke **MORE** than 10 Cigarettes per Day: Use After Completing Step 1.

If You Smoke 10 or **LESS** Cigarettes per Day: Start with Step 2.

7 PATCHES (1 WEEK KIT)

The full course of treatment for STEP 2 is 2 or 6 weeks (depending on how many cigarettes you smoke per day); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

COMPARE TO HABITROL® NICOTINE TRANSDERMAL SYSTEM PATCH active ingredient*



Principal Display Panel

NDC 70000-0760-1

Nicotine Transdermal System Patch

7 mg Delivered Over 24 Hours

STOP SMOKING AID

7 mg

STEP 3

Includes: **Behavior Support Program with SELF-HELP GUIDE**

For Use After Completing Step 2

7 PATCHES (1 WEEK KIT)

The full treatment course for STEP 3 is 14 patches (2 weeks); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

COMPARE TO HABITROL® NICOTINE TRANSDERMAL SYSTEM PATCH active ingredient*

E Z 0 1 5 1 0 5 1

Drug Facts

Active ingredient (in each patch) **Purpose**
Nicotine, 7 mg delivered over 24 hours. Stop smoking aid

Use reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking.

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.

- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- an allergy to adhesive tape or have skin problems, because you are more likely to get rashes.
- stomach ulcer or diabetes.
- history of seizures.

Ask a doctor or pharmacist before use if you are using a non-nicotine stop smoking drug or taking a prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

When using this product if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if skin redness caused by the patch does not go away after four days, or if your skin swells, or you get a rash

- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose, such as nausea, vomiting, dizziness, weakness and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Used patches have enough nicotine to poison children and pets. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222). Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.

Directions If you are under 18 years of age, ask a doctor before use

- before using this product, read the enclosed self-help guide for complete directions and other information
- begin using the patch on your quit day
- if you smoke more than 10 cigarettes per day, use the following schedule below:

Weeks 1 thru 4	Weeks 5 and 6	Weeks 7 and 8
STEP 1 use one 21 mg patch/day	STEP 2 use one 14 mg patch/day	STEP 3 use one 7 mg patch/day

- if you smoke 10 or less cigarettes per day, start with Step 2 for 6 weeks, then Step 3 for 2 weeks
- apply one new patch every 24 hours on skin that is dry, clean and hairless
- remove backing from patch and immediately press onto skin. Hold for 10 seconds.
- wash hands after applying or removing patch. Save pouch to use for patch disposal. Dispose of the used patches by folding sticky ends together and putting in pouch.
- the used patch should be removed and a new one applied to a different skin site at the same time each day
- if you have vivid dreams, you may remove the patch at bedtime and apply a new one in the morning.
- do not wear more than one patch at a time
- do not cut patch in half or into smaller pieces
- do not leave patch on for more than 24 hours because it may irritate your skin and loses strength after 24 hours
- to avoid possible burns, remove patch before undergoing any MRI (magnetic resonance imaging) procedures
- it is important to complete treatment. If you feel you need to use the patch for a longer period to keep from smoking, talk to your health care provider.

Other information store at 20-25°C (68-77°F)

Inactive ingredients acrylate adhesive, aluminumized polyester, cellulose paper, methacrylic acid copolymer

Comments or Questions? Call 1-800-585-8882 Weekdays (8am-5pm EST)

See bottom panel for lot number and expiration date.
*Habitrol® is a registered trademark of Dr. Reddy's Laboratories SA.

Nicotine Transdermal System Patch
7 mg Delivered Over 24 Hours
STOP SMOKING AID

7 mg

STEP 3

For your family's protection, patches are supplied in child-resistant pouches. Do not use if individual pouch is open or torn.

Not for sale to persons under 18 years of age. Proof of age required. Not for sale in vending machines or from any source where proof of age cannot be verified.

See self-help guide inside.

LEADER[™]

NDC 70000-0760-1

Nicotine Transdermal System Patch
7 mg Delivered Over 24 Hours | **STOP SMOKING AID**

7 mg

STEP 3

Includes:
Behavior Support Program with SELF-HELP GUIDE
For Use After Completing Step 2

COMPARE TO HABITROL[®] NICOTINE TRANSDERMAL SYSTEM PATCH active ingredient*

100% Money Back Guarantee

The full treatment course for STEP 3 is 14 patches (2 weeks); this package contains 7 patches (1 week course) only. Read the enclosed self-help guide for additional information.

7 PATCHES (1 WEEK KIT)

CIN 6107569 REV. 02/26

0 96295 14598 4

Nicotine Transdermal System Patch
7 mg Delivered Over 24 Hours
STOP SMOKING AID

7 mg

STEP 3

Cardinal Health[™]
DISTRIBUTED BY: CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myladder.com 1-800-210-4393
Essential to Care[™] since 1970

©2025 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the LEADER LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

100% Money Back Guarantee
Returns to place of purchase. Not cashed.

NICOTINE TRANSDERMAL SYSTEM STEP 1

nicotine patch, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0762
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	21 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
Methacrylic Acid (UNII: 1CS02G8656)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date

1	NDC:70000-0762-1	7 in 1 CARTON	03/20/2026	
1		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:70000-0762-2	14 in 1 CARTON	03/20/2026	
2		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	03/20/2026	

NICOTINE TRANSDERMAL SYSTEM STEP 2

nicotine patch, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0761
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	14 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
Methacrylic Acid (UNII: 1CS02G8656)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0761-1	7 in 1 CARTON	03/20/2026	
1		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:70000-0761-2	14 in 1 CARTON	03/20/2026	
2		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
NDA	NDA020076	03/20/2026	

NICOTINE TRANSDERMAL SYSTEM STEP 3

nicotine patch, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0760
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	7 mg in 24 h

Inactive Ingredients

Ingredient Name	Strength
Methacrylic Acid (UNII: 1CS02G8656)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0760-1	7 in 1 CARTON	03/20/2026	
1		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		
2	NDC:70000-0760-2	14 in 1 CARTON	03/20/2026	
2		24 h in 1 PATCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020076	03/20/2026	

Labeler - CARDINAL HEALTH 110, LLC. DBA LEADER (063997360)