

MUCINEX FAST-MAX DAY COLD AND FLU AND MUCINEX NIGHTSHIFT NIGHT SEVERE COLD AND FLU MAXIMUM STRENGTH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and triprolidine hydrochloride
Reckitt Benckiser LLC

Mucinex® Fast-Max® Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu

Maximum Strength

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purposes</i>
<i>Mucinex Fast-Max Day Cold & Flu</i>	
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

<i>Active ingredients (in each caplet)</i>	<i>Purposes</i>
<i>Mucinex Nightshift Night Severe Cold & Flu</i>	
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Triprolidine HCl 1.25 mg	Antihistamine

Uses

Mucinex Fast-Max Day Cold & Flu

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial

passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Night Severe Cold & Flu

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sneezing
 - sinus congestion and pressure
 - runny nose
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription

drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (***Nightshift Night Severe Cold & Flu only***)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (***Nightshift Night Severe Cold & Flu only***)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (***Nightshift Night Severe Cold & Flu only***)

When using this product

- **do not use more than directed**
- excitability may occur, especially in children (***Nightshift Night Severe Cold & Flu only***)
- marked drowsiness may occur (***Nightshift Night Severe Cold & Flu only***)
- alcohol, sedatives, and tranquilizers may increase drowsiness (***Nightshift Night Severe Cold & Flu only***)
- avoid alcoholic drinks (***Nightshift Night Severe Cold & Flu only***)
- use caution when driving a motor vehicle or operating machinery (***Nightshift Night Severe Cold & Flu only***)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Mucinex Fast-Max Day Cold & Flu

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Severe Cold & Flu

- **do not take more than directed (see Overdose warning)**
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients Mucinex Fast-Max Day Cold & Flu

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients Mucinex Nightshift Night Severe Cold & Flu

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - Kit Carton

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

NDC 72854-241-20

MAXIMUM STRENGTH

Mucinex®

FAST-MAX® NIGHTSHIFT

DAY
COLD & FLU
Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

HEADACHE
SORE THROAT
CHEST CONGESTION
BODY PAIN
FEVER
COUGH

ALL IN
ONE*

NASAL CONGESTION
SINUS CONGESTION
SINUS PRESSURE

ACTUAL SIZE
FOR AGES 12+
12 CAPLETS

FAST
RELEASE

NIGHT
SEVERE COLD & FLU

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal Decongestant
Triprolidine HCl - Antihistamine

HEADACHE
SORE THROAT
ITCHY THROAT
BODY PAIN
FEVER
COUGH

ALL IN
ONE*

NASAL CONGESTION
SNEEZING
RUNNY NOSE

ACTUAL SIZE
FOR AGES 12+

8 CAPLETS

TOTAL 20 CAPLETS

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

MAXIMUM STRENGTH

Mucinex
FAST-MAX NIGHTSHIFT

DAY
COLD & FLU

NIGHT
SEVERE COLD & FLU

FAST RELEASE. POWERFUL SYMPTOM RELIEF!

MAXIMUM STRENGTH

NDC 72854-241-20

Mucinex

FAST-MAX NIGHTSHIFT

DAY
COLD & FLU

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

NIGHT
SEVERE COLD & FLU

Acetaminophen - Pain Reliever/
Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Phenylephrine HCl - Nasal Decongestant
Triprolidine HCl - Antihistamine



ACTUAL SIZE
FOR AGES 12+



FAST
RELEASE



ACTUAL SIZE
FOR AGES 12+



12 CAPLETS

TOTAL 20 CAPLETS

8 CAPLETS

MAXIMUM STRENGTH
Mucinex
FAST-MAX NIGHTSHIFT

DAY **NIGHT**
COLD & FLU SEVERE COLD & FLU
Maximum Strength per 4-hour dose
Keep carton for full information.

Fast-Max Day Cold & Flu helps to relieve these symptoms day or night. Nightshift Night Severe Cold & Flu helps to relieve these symptoms at night. Do not take the Mucinex Fast-Max Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

PARENTS:
Learn about their medicine advice
www.StopMedicineAbuse.org

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.



HEALTH • HYGIENE • HOME

www.mucinex.com

Patents: www.rb.com/patents

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2021 RB Health

3176344 012721



6382499531 1

LOT:

EXP.:

MADE IN:

3176344



Do not take the Mucinex Fast-Max Day Cold & Flu and Mucinex Nightshift Night Severe Cold & Flu caplets at the same time. Always wait at least 4 hours before taking another dose of Mucinex caplets. TAKE ONLY AS DIRECTED.

3176344

Drug Facts

Active ingredients (in each caplet)

Purposes

010621

PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION

Mucinex Fast-Max Day Cold & Flu

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine HCl 5 mg.....	Nasal decongestant

Active ingredients (in each caplet)**Purposes****Mucinex Nightshift Night Severe Cold & Flu**

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant
Triprolidine HCl 1.25 mg.....	Antihistamine

Uses**Mucinex Fast-Max Day Cold & Flu**

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Mucinex Nightshift Night Severe Cold & Flu

- temporarily relieves these common cold and flu symptoms:
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - sneezing
 - sinus congestion and pressure
 - runny nose
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Reseal Area**Drug Facts (continued)**

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma (*Nightshift Night Severe Cold & Flu only*)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (*Nightshift Night Severe Cold & Flu only*)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Nightshift Night Severe Cold & Flu only*)

When using this product

- do not use more than directed
- excitability may occur, especially in children (*Nightshift Night Severe Cold & Flu only*)
- marked drowsiness may occur (*Nightshift Night Severe Cold & Flu only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nightshift Night Severe Cold & Flu only*)
- avoid alcoholic drinks (*Nightshift Night Severe Cold & Flu only*)
- use caution when driving a motor vehicle or operating machinery (*Nightshift Night Severe Cold & Flu only*)

Hinge

Hinge

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Mucinex Fast-Max Day Cold & Flu

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Mucinex Nightshift Night Severe Cold & Flu

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients Mucinex Fast-Max Day Cold & Flu

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Inactive ingredients Mucinex Nightshift Night Severe Cold & Flu

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

MUCINEX FAST-MAX DAY COLD AND FLU AND MUCINEX NIGHTSHIFT NIGHT SEVERE COLD AND FLU MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride, and triprolidine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-241
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-241-10	1 in 1 CARTON	07/01/2021	
1		1 in 1 KIT		
2	NDC:72854-241-20	2 in 1 CARTON	07/01/2021	

2		1 in 1 KIT	
3	NDC:72854-241-40	4 in 1 CARTON	07/01/2021
3		1 in 1 KIT	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	6
Part 2	1 BLISTER PACK	4

Part 1 of 2

MAXIMUM STRENGTH MUCINEX FAST-MAX COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, film coated

Product Information

Item Code (Source)	NDC:63824-575
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6130)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	MSC;VW
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2021	

Part 2 of 2

MUCINEX NIGHTSHIFT SEVERE COLD AND FLU MAXIMUM STRENGTH

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride, and triprolidine hydrochloride tablet, coated

Product Information

Item Code (Source)	NDC:72854-234
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	

CROSPROVIDONE (UNII: 2S7830E561)
FERRIC OXIDE RED (UNII: 1K09F3G675)
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)
MICA (UNII: V8A1AW0880)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
POLYVINYL ALCOHOL GRAFT POLYETHYLENE GLYCOL COPOLYMER (3:1; 45000 MW) (UNII: 23ZQ42JZZH)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
STEARIC ACID (UNII: 4ELV7Z65AP)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	yellow	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	VW;LOGOcrescentmoonplus
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2021	

Labeler - Reckitt Benckiser LLC (094405024)