

**QUALITY CHOICE MAXIMUM STRENGTH ORAL PAIN RELIEVING-
benzocaine gel
Chain Drug Marketing Association Inc**

QC Maximum Strength Oral Analgesic 0.42oz 95362

Active ingredient Purpose

Benzocaine 20%.....Oral pain reliever

Uses

- temporarily relieves pain due to:
- toothache
- canker sores
- gum pain
- denture pain
- cheek bites

Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

Do not use

- more than directed
- for more than 7 days unless directed by a doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

If pregnant or breast-feeding , ask a health professional before use.

Keep out of reach of children. In cases of overdose, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 2 years of age and older.	Remove cap and cut tip of tube. Apply a small amount of product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or doctor.
Children under 12 years of age.	Should be supervised in the use of this product.
Children under 2 years of age.	Do not use.

Other information

- store at room temperature 20° to 25°C (68° to 77°F)
- this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted
- do not use continuously

Inactive ingredients

glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

Distributed By C.D.M.A., Inc.

43157 W 9 Mile Rd

Novi, MI 48375

www.qualitychoice.com

Made in South Korea



SAFETY-SEALED TUBE TIP

Maximum Strength

Oral Pain Relieving Gel

Fast Pain Relief

Benzocaine 20% (Oral Pain Reliever)

Fast Relief from:

Toothache | Canker Sores | Gum and Denture Pains | Cheek Bites

- Toothache
- Canker Sores
- Gum and Denture Pains
- Cheek Bites

Maximum Strength

SAFETY-SEALED TUBE TIP

Oral Pain Relieving Gel

Fast Pain Relief

Benzocaine 20% (Oral Pain Reliever)

NET WT. 0.42 oz (11.9 g)



Maximum Strength
Oral Pain
Relieving Gel

VC-896



Distributed by C.D.M.A., Inc.®
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 800-935-2362

MADE IN CHINA

95362-ZDP-BX-R1
0C-MD-ZDP105-R3

LOT & EXP.

Drug Facts (continued)	
Active Ingredient Benzocaine 20% Oral pain reliever	Purpose Oral pain reliever
Uses ■ for the temporary relief of pain due to: ■ toothache ■ canker sores ■ gum pain ■ denture pain ■ cheek bites	
Warnings Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: ■ pale, gray, or blue colored skin (cyanosis) ■ headache ■ rapid heart rate ■ shortness of breath ■ dizziness or lightheadedness ■ fatigue or lack of energy Allergy Alert: do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. Do not use ■ more than directed ■ for more than 7 days unless directed by a doctor ■ for teething ■ in children under 2 years of age	
Stop use and ask a doctor if ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain or redness does not go away ■ swelling, rash or fever develops	
Other Information ■ store at room temperature 20-25°C (68-77°F) ■ this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted. Do not use continuously.	
Inactive ingredients glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol	

IS OPEN OR IF TUBE TIP IS CUT PRIOR TO OPENING. TAMPER EVIDENT: DO NOT USE IF OUTER CARTON

WARNINGS AND PRODUCT INFORMATION. KEEP OUTER CARTON FOR COMPLETE

QUALITY CHOICE MAXIMUM STRENGTH ORAL PAIN RELIEVING benzocaine gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-953
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	20 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
PEPPERMINT OIL (UNII: AV092KU4JH)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-953-11	1 in 1 CARTON	05/31/2013	
1		11.76 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	05/31/2013	

Labeler - Chain Drug Marketing Association Inc (011920774)