

ST. JOSEPH LOW DOSE ASPIRIN- aspirin tablet, film coated
Foundation Consumer Healthcare

St. Joseph Low Dose Aspirin

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID) ¹

¹ nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pain
- ask your doctor about other uses for St. Joseph Safety Coated 81 mg Aspirin

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms occur. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years of age and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years of age: do not use unless directed by a doctor

Other information

- store at 20-25°C (68-77°F)
- close cap tightly after use

Inactive ingredients

colloidal anhydrous silica, FD&C red #40, FD&C yellow #6, hypromellose, methacrylic

acid- ethyl acrylate copolymer, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

Call **1-855-785-6381** weekdays 9 AM to 5 PM EST

Dist. by: Foundation Consumer Healthcare, LLC.
Pittsburgh, PA 15205

PRINCIPAL DISPLAY PANEL - 36 Tablet Bottle Box

•ADULT ASPIRIN REGIMEN †•

ST. JOSEPH ®

LOW DOSE ASPIRIN
SAFETY COATED
PAIN RELIEVER (NSAID)

81mg

AMERICA'S
ASPIRIN

100
YEARS

DOCTOR RECOMMENDED

Enteric
Coated

36
Coated
Tablets

No Print/No Varnish Lot & Expiry Area



The Product Does Not contain Gluten. Easy to swallow coated tablets.
 Coating Helps Protect Against Stomach Upset.
 Talk to your doctor before starting an aspirin regimen.
 Aspirin is not right for everyone.
 Dist. by: Foundation Consumer Healthcare, LLC,
 Pittsburgh, PA 15212 ©2022
 stjosephaspirin.com

B-0603-750-07R
 REV-A

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts
Active ingredient (in each tablet) Aspirin 81 mg (NSAID)
 *nonsteroidal anti-inflammatory drug

Purpose Pain reliever

Uses Temporarily relieves minor aches and pain
 ask your doctor about other uses for St. Joseph Safety Coated 81 mg Aspirin

Warnings
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
 Allergy alert: Aspirin may cause a severe allergic reaction, which may include: hives; facial swelling; asthma (wheezing); shock
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinning (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed.
 Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.

Drug Facts (continued)
 Ask a doctor before use if:
 stomach bleeding warning applies to you
 you have a history of stomach problems, such as heartburn
 you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 you are taking a diuretic
 you have asthma
 Ask a doctor or pharmacist before use if you are taking a prescription drug for gout, diabetes or arthritis.
 Stop use and ask a doctor if:
 an allergic reaction occurs. Seek medical help right away.
 you experience any of the following signs or stomach bleeding:
 red faint vomit blood
 have bloody or black stools
 have stomach pain that does not get better
 pain gets worse or lasts more than 10 days
 ringing in the ears or loss of hearing occurs
 redness or swelling is present
 new symptoms occur.
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)
Directions drink a full glass of water with each dose
 adults and children 12 years of age and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 children under 12 years of age: do not use unless directed by a doctor

Other information
 store at 20-25°C (68-77°F) close cap tightly after use

Inactive ingredients colloidal anhydrous silica, FD&C red #40, FD&C yellow #6, hydroxypropyl methylcellulose, methacrylic acid-ethyl acrylate copolymer, microcrystalline cellulose, polydextrose, polyethylene glycol, pregelatinized starch, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?
 Call 1-855-785-6381 weekdays 9 AM to 5 PM EST

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ADULT ASPIRIN REGIMEN.
ST. JOSEPH
 LOW DOSE ASPIRIN
 SAFETY COATED PAIN RELIEVER (NSAID)
 81mg
 DOCTOR RECOMMENDED
 100 YEARS
 36 Coated Tablets
 Enteric Coated

ADULT ASPIRIN REGIMEN.
ST. JOSEPH
 LOW DOSE ASPIRIN
 SAFETY COATED PAIN RELIEVER (NSAID)
 81mg
 DOCTOR RECOMMENDED
 100 YEARS
 36 Coated Tablets
 Enteric Coated

LEBGG775S2

ST. JOSEPH LOW DOSE ASPIRIN

aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69536-181
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TRIACETIN (UNII: XHX3C3X673)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
SHELLAC (UNII: 46N107B71O)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

Product Characteristics

Color	white (PEACH)	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	SJ
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69536-181-15	1 in 1 BOX	02/17/2017	
1		150 in 1 BOTTLE; Type 0: Not a Combination Product		

2	NDC:69536-181-36	1 in 1 BOX	02/17/2017	
2		36 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:69536-181-12	1 in 1 BOX	02/17/2017	
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:69536-181-20	1 in 1 BOX	02/17/2017	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:69536-181-54	1 in 1 BOX	02/17/2017	05/01/2018
5		54 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:69536-181-65	1 in 1 BOX	02/17/2017	
6		365 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/17/2017	

Labeler - Foundation Consumer Healthcare (079675882)

Revised: 4/2024

Foundation Consumer Healthcare