

**IBUPROFEN MINIS- ibuprofen capsule, liquid filled  
McKesson (Health Mart)**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)**

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - slurred speech
  - trouble breathing
  - leg swelling
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**

- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### **Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.
- swallow whole; do not crush, chew, or dissolve

### **Inactive ingredients**

FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol solution, white ink

### **Questions or comments ?**

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the active ingredient in Advil® Liqui-Gels® minis\*\*

Ibuprofen Capsules 200 mg

Pain Reliever/ Fever Reducer (NSAID)

Smaller Capsule • Same Strength

Softgels† (†Liquid-filled capsules)

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels® mini.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Distributed by McKesson

One Post Street, San Francisco, CA 94104

healthmart.com

### **Product Label**

Exp. Date:  
Lot No.:

**McKESSON**  
Distributed by McKesson  
One East Wacker Drive  
Murray Park Center  
Irvine, CA 94104  
healthmart.com

KEEP OUTER CARTON FOR COMPLETE  
WARNINGS AND PRODUCT INFORMATION. FDC05655  
PLD-A560A

TAPE HERE TO OPEN  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



**Drug Facts**

**Active ingredient**  
(in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)  
\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/  
fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
headache  
muscular aches  
minor pain of arthritis  
toothache  
backache  
the common cold  
menstrual cramps  
temporarily reduces fever

**Warnings**

**Allergy alert:** ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
hives  
facial swelling  
asthma (wheezing)  
shock  
skin reddening  
rash  
blisters. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older  
have had stomach ulcers or bleeding problems  
take a blood thinning (anticoagulant) or steroid drug  
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
have 3 or more alcoholic drinks every day while using this

**Drug Facts** (continued)

product take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer  
right before or after heart surgery

**Ask a doctor before use if**  
the stomach bleeding warning applies to you  
you have problems or serious side effects from taking pain relievers or fever reducers  
you have a history of stomach problems, such as heartburn  
you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke  
you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**  
under a doctor's care for any serious condition  
taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
taking any other drug

**When using this product,** take with food or milk if stomach upset occurs.

**Stop use and ask a doctor if**  
you experience any of the following signs of stomach bleeding:  
feel faint  
vomit blood  
have bloody or black stools  
have stomach pain that does not get better

**Drug Facts** (continued)

you have symptoms of heart, problems or stroke:  
chest pain  
trouble breathing  
weakness in one part or side of body  
slurred speech  
leg swelling  
pain gets worse or lasts more than 10 days  
fever gets worse or lasts more than 3 days  
redness or swelling is present in the painful area  
any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
do not take more than directed  
the smallest effective dose should be used  
adults and children 12 years and over:  
take 1 capsule every 4 to 6 hours while symptoms persist  
if pain or fever does not respond to 1 capsule, 2 capsules may be used  
do not exceed 6 capsules in 24 hours, unless directed by a doctor  
children under 12 years: ask a doctor

**Other information**  
each capsule contains: potassium 20 mg  
read all warnings and directions before use. Keep carton.

**Drug Facts** (continued)

store at 20°-25°C (68°-77°F)  
avoid excessive heat above 40°C (104°F)  
swallow whole; do not crush, chew, or dissolve

**Inactive ingredients**  
FD&C green #3, gelatin, medium-chain triglycerides, polyethylene glycol, potassium hydroxide, purified water, sorbitol sorbitan solution, white ink

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels® minis.

NDC 62011-0399-1

**HealthMart.** Compare to the active ingredient in Advil® Liqui-Gels® minis\*\*

**MINIS**

**Ibuprofen**  
Capsules 200 mg  
Pain Reliever/Fever Reducer (NSAID)  
Smaller Capsule • Same Strength

**20 Softgels**<sup>†</sup> (Liquid-filled capsules)

ACTUAL SIZE

**HEALTH MART Ibuprofen 200 mg MINIS Softgels**

<b>IBUPROFEN MINIS</b> ibuprofen capsule, liquid filled			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:62011-0399
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

**Inactive Ingredients**

Ingredient Name	Strength
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	

**Product Characteristics**

Color	BLUE	Score	no score
Shape	CAPSULE	Size	13mm
Flavor		Imprint Code	P33
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62011-0399-1	1 in 1 BOX	02/28/2018	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206568	02/28/2018	

**Labeler** - McKesson (Health Mart) (177667227)

Revised: 3/2019

McKesson (Health Mart)