

PLEO MUC- mucor racemosus ointment
Sanum Kehlbeck GmbH & Co. KG

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Pleo™ Muc
OINTMENT 3X

for topical use only
Homeopathic External Analgesic Medicine

1.05 oz (30g)

Indications

For temporary relief of pain of bruises, sprains, swelling of joints and muscles.

DOSAGE

Apply thinly on the affected area 1–3 times daily.

WARNING

If symptoms persist more than a few days, contact a licensed practitioner. As with any other drug, if you are pregnant or nursing a baby, seek the advice of a health care professional before using this product.

Keep this and all other medications out of the reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Protect from light and heat.

Tamper Evident

Do not use this product if metal seal under the tube cap is missing or broken or if imprinted security strip on carton is torn.

INGREDIENTS

Active Ingredients

1 g ointment contains; 0.1 g Mucor racemosus 3x.

Inactive Ingredients

lactic acid, lanolin alcohol, glycerylmonostearate, propylene glycol, polyglyceryl-3, cocoate magnesium sulfate, water.

Made in Germany

Distributed by:
SANUM USA Corp.

1465 Slater Road
Ferndale, WA 98248

Manufactured By:
Sanum-Kehlbeck
GmbH & Co. KG

Rev. 12/2006

PRINCIPAL DISPLAY PANEL - 30 g Ointment Carton

**Pleo™ Muc
OINTMENT 3X**

**for topical
use only
Homeopathic
External Analgesic
Medicine**

Indications: For
temporary relief of
pain of bruises,
sprains, swelling of
joints and muscles.

ORIGINAL ENDERLEIN

1.05 oz (30g)



1806-1

Pleo™ Muc OINTMENT 3X

for topical
use only
Homeopathic
External Analgesic
Medicine

Indications: For temporary relief of pain of bruises, sprains, swelling of joints and muscles.



1.05 oz (30g)

DOSEAGE: Apply thinly on the affected area 1-3 times daily.

WARNING: If symptoms persist more than a few days, contact a licensed practitioner. As with any other drug, if you are pregnant or nursing a baby, seek the advice of a health care professional before using this product. **Keep this and all other medications out of the reach of children.** In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Protect from light and heat.

Lot #:

Exp. Date:

Tamper Evident: Do not use this product if metal seal under the tube cap is missing or broken or if imprinted security strip on carton is torn.

INGREDIENTS: Active Ingredients: 1 g ointment contains; 0,1 g Mucor racemosus 3x. Inactive Ingredients: lactic acid, lanolin alcohol, glycerylmonostearate, propylene glycol, polyglyceryl-3, cocoate magnesium sulfate, water.

Made in Germany

Distributed by:
SANUM USA Corp.
1465 Slater Road
Ferndale, WA 98248

Manufactured By:
Sanum-Kehlbeck
GmbH & Co. KG

Rev. 12/2006

1806-1

4005A3XFS30US612

PLEO MUC

mucor racemosus ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:60681-1806	
Route of Administration	TOPICAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
mucor racemosus (UNII: 17RH99LQ7G) (mucor racemosus - UNII:17RH99LQ7G)		mucor racemosus	3 [hp_X] in 30 g	
Inactive Ingredients				
Ingredient Name			Strength	
glyceryl monostearate (UNII: 230OU9XXE4)				
lactic acid (UNII: 33X04XA5AT)				
propylene glycol (UNII: 6DC9Q167V3)				
water (UNII: 059QF0K00R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60681-1806-1	1 in 1 CARTON		
1		30 g in 1 TUBE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
Unapproved homeopathic		03/19/1996		

Labeler - Sanum Kehlbeck GmbH & Co. KG (318386133)

Revised: 11/2009

Sanum Kehlbeck GmbH & Co. KG