

BETAMETHASONE DIPROPIONATE- betamethasone dipropionate spray

Sun Pharmaceutical Industries, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use **BETAMETHASONE DIPROPIONATE SPRAY** safely and effectively. See full prescribing information for **BETAMETHASONE DIPROPIONATE SPRAY**.

BETAMETHASONE DIPROPIONATE spray, for topical use
Initial U.S. Approval: 1975

INDICATIONS AND USAGE

Betamethasone dipropionate spray is a corticosteroid indicated for the treatment of mild to moderate plaque psoriasis in patients 18 years of age or older. (1)

DOSAGE AND ADMINISTRATION

- Apply to the affected skin areas twice daily. Rub in gently. (2)
- Use betamethasone dipropionate spray for up to 4 weeks and not beyond. (2)
- Discontinue treatment when control is achieved. (2)
- Do not use if atrophy is present at the treatment site. (2)
- Do not use with occlusive dressings unless directed by a physician. (2)
- Avoid use on the face, scalp, axilla, groin, or other intertriginous areas. (2)
- Not for oral, ophthalmic, or intravaginal use. (2)

DOSAGE FORMS AND STRENGTHS

Spray: 0.05% (equivalent to 0.5 mg betamethasone/g) (3)

CONTRAINDICATIONS

- None (4)

WARNINGS AND PRECAUTIONS

- Betamethasone dipropionate spray can produce reversible HPA axis suppression with the potential for glucocorticosteroid insufficiency during or after treatment. High Potency corticosteroids, large treatment surface areas, prolonged use, use of occlusive dressings, altered skin barrier, liver failure and use in pediatric patients may predispose to HPA axis suppression. Use of topical corticosteroids may require periodic evaluation for HPA axis suppression. Modify use if HPA axis suppression develops. (5.1)
- Topical corticosteroids may increase the risk of cataracts and glaucoma. If visual symptoms occur, consider referral to an ophthalmologist. (5.2)

ADVERSE REACTIONS

The most common adverse reactions ($\geq 1\%$) are application site reactions, including pruritus, burning and/or stinging, pain, and atrophy. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Taro Pharmaceuticals U.S.A., Inc. at 1-866-923-4914 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 6/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Effects on Endocrine System
- 5.2 Ophthalmic Adverse Reactions
- 5.3 Allergic Contact Dermatitis

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied/Storage
- 16.2 Handling/Instructions for the Pharmacist

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Betamethasone dipropionate spray is indicated for the treatment of mild to moderate plaque psoriasis in patients 18 years of age or older.

2 DOSAGE AND ADMINISTRATION

Shake well before use.

Apply betamethasone dipropionate spray to the affected skin areas twice daily and rub in gently.

Use betamethasone dipropionate spray for up to 4 weeks of treatment. Treatment beyond 4 weeks is not recommended.

Discontinue betamethasone dipropionate spray when control is achieved.

Do not use if atrophy is present at the treatment site.

Do not bandage, cover, or wrap the treated skin area unless directed by a physician.

Avoid use on the face, scalp, axilla, groin, or other intertriginous areas.

Betamethasone dipropionate spray is for topical use only. It is not for oral, ophthalmic, or intravaginal use.

3 DOSAGE FORMS AND STRENGTHS

Spray, 0.05% for topical use. Each gram of betamethasone dipropionate spray contains 0.643 mg betamethasone dipropionate USP (equivalent to 0.5 mg betamethasone) in a slightly thickened, white to off-white oil-in-water emulsion.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Effects on Endocrine System

Betamethasone dipropionate spray can produce reversible hypothalamic-pituitary-adrenal (HPA) axis suppression with the potential for glucocorticosteroid insufficiency. This may occur during or after withdrawal of treatment. Factors that predispose to HPA axis suppression include the use of high-potency corticosteroids, large treatment surface areas, prolonged use, use of occlusive dressings, altered skin barrier, liver failure, and young age.

Evaluation for HPA axis suppression may be done by using the adrenocorticotrophic hormone (ACTH) stimulation test.

In a study including 48 evaluable subjects 18 years of age or older with moderate to severe plaque psoriasis, abnormal ACTH stimulation test results suggestive of adrenal suppression were identified in 5 out of 24 (20.8%) subjects after treatment with betamethasone dipropionate spray twice daily for 15 days. No subject (0 out of 24) had abnormal ACTH stimulation test results after treatment with betamethasone dipropionate spray twice daily for 29 days [*see Clinical Pharmacology (12.2)*].

If HPA axis suppression is documented, gradually withdraw the drug, reduce the frequency of application, or substitute with a less potent corticosteroid. If signs and symptoms of steroid withdrawal occur, supplemental systemic corticosteroids may be required.

Systemic effects of topical corticosteroids may also manifest as Cushing's syndrome, hyperglycemia, and glucosuria. These events are rare and generally occur after prolonged exposure to larger than recommended doses, particularly with high-potency topical corticosteroids.

Minimize the unwanted risks from endocrine effects by mitigating the risk factors favoring increased systemic bioavailability and by using the product as recommended [*see Dosage and Administration (2)*].

Pediatric patients may be more susceptible to systemic toxicity due to their larger skin surface to body mass ratios. Use of betamethasone dipropionate spray is not recommended in pediatric patients [*see Use in Specific Populations (8.4)*].

5.2 Ophthalmic Adverse Reactions

Use of topical corticosteroids, including betamethasone dipropionate spray, may increase the risk of posterior subcapsular cataracts and glaucoma. Cataracts and glaucoma have been reported postmarketing with the use of topical corticosteroid products, including betamethasone dipropionate [see Adverse Reactions (6.2)] .

Avoid contact of betamethasone dipropionate spray with eyes. Advise patients to report any visual symptoms and consider referral to an ophthalmologist for evaluation.

5.3 Allergic Contact Dermatitis

Allergic contact dermatitis with corticosteroids is usually diagnosed by observing failure to heal rather than noting a clinical exacerbation. Corroborate such an observation with appropriate diagnostic patch testing. If irritation develops, discontinue the topical corticosteroid and institute appropriate therapy.

6 ADVERSE REACTIONS

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

In two randomized, multicenter, prospective vehicle-controlled clinical trials, subjects with moderate plaque psoriasis of the body applied betamethasone dipropionate spray or vehicle spray twice daily for 4 weeks. A total of 352 subjects applied betamethasone dipropionate spray and 180 subjects applied vehicle spray.

Adverse reactions that occurred in at least 1% of subjects treated with betamethasone dipropionate spray for up to 28 days are presented in Table 1.

Table 1: Adverse Reactions Occurring in $\geq 1\%$ of Subjects Treated with Betamethasone Dipropionate Spray for up to Four Weeks

	Betamethasone Dipropionate Spray <i>b.i.d.</i> (N=352)	Vehicle Spray <i>b.i.d.</i> (N=180)
Application site pruritus	6.0%	9.4%
Application site burning and/or stinging	4.5%	10.0%
Application site pain	2.3%	3.9%
Application site atrophy	1.1%	1.7%

Less common adverse reactions (with occurrence lower than 1% but higher than 0.1%) in subjects treated with betamethasone dipropionate spray were application site reactions including telangiectasia, dermatitis, discoloration, folliculitis and skin rash, in addition to dysgeusia and hyperglycemia. These adverse reactions were not observed in

subjects treated with vehicle.

6.2 Postmarketing Experience

Because adverse reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Postmarketing reports for local adverse reactions to topical corticosteroids have also included striae, irritation, dryness, acneiform eruptions, hypopigmentation, perioral dermatitis, allergic contact dermatitis, secondary infection, hypertrichosis, and miliaria.

Hypersensitivity reactions, consisting of predominantly skin signs and symptoms, e.g., contact dermatitis, pruritus, bullous dermatitis, and erythematous rash have been reported.

Ophthalmic adverse reactions of cataracts, glaucoma, and increased intraocular pressure have been reported with the use of topical corticosteroids, including topical betamethasone products.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no available data on betamethasone dipropionate spray use in pregnant women to evaluate a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes.

Observational studies suggest an increased risk of low birthweight infants with the use of greater than 300 grams of potent or very potent topical corticosteroid during a pregnancy. Advise pregnant women that betamethasone dipropionate spray may increase the risk of having a low birthweight infant and to use betamethasone dipropionate spray on the smallest area of skin and for the shortest duration possible.

In animal reproduction studies, increased malformations, including umbilical hernias, cephalocele, and cleft palate, were observed after intramuscular administration of betamethasone dipropionate to pregnant rabbits during the period of organogenesis (see *Data*). The available data do not allow the calculation of relevant comparisons between the systemic exposure of betamethasone dipropionate observed in animal studies to the systemic exposure that would be expected in humans after topical use of betamethasone dipropionate spray.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Administration of 0.05 mg/kg betamethasone dipropionate intramuscularly to pregnant

rabbits during the period of organogenesis caused malformations. The abnormalities observed included umbilical hernias, cephalocele, and cleft palate.

8.2 Lactation

Risk Summary

There are no data regarding the presence of betamethasone dipropionate in human milk, the effects on the breastfed infant, or the effects on milk production after topical application of betamethasone dipropionate spray to women who are breastfeeding.

It is possible that topical administration of betamethasone dipropionate could result in sufficient systemic absorption to produce detectable quantities in human milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for betamethasone dipropionate spray and any potential adverse effects on the breastfed infant from betamethasone dipropionate spray or from the underlying maternal condition.

Clinical Considerations

To minimize potential exposure to the breastfed infant via breast milk, use betamethasone dipropionate spray on the smallest area of skin and for the shortest duration possible while breastfeeding. Advise breastfeeding women not to apply betamethasone dipropionate spray directly to the nipple and areola to avoid direct infant exposure [*see Use in Specific Populations (8.4)*] .

8.4 Pediatric Use

Safety and effectiveness of betamethasone dipropionate spray in patients younger than 18 years of age have not been studied; therefore use in pediatric patients is not recommended. Because of a higher ratio of skin surface area to body mass, pediatric patients are at greater risk of systemic toxicity, including HPA axis suppression and adrenal insufficiency, when treated with topical drugs. [*see Warnings and Precautions (5.1)*]

Rare systemic effects such as Cushing's syndrome, linear growth retardation, delayed weight gain, and intracranial hypertension have been reported in pediatric patients, especially those with prolonged exposure to large doses of high potency topical corticosteroids.

Local adverse reactions including skin atrophy have also been reported with use of topical corticosteroids in pediatric patients.


8.5 Geriatric Use

Clinical studies of betamethasone dipropionate spray did not include sufficient numbers of subjects who were 65 years of age or older to determine whether they respond differently from younger subjects.

11 DESCRIPTION

Betamethasone dipropionate spray contains 0.0643% betamethasone dipropionate (equivalent to 0.05% betamethasone), a synthetic, fluorinated corticosteroid for topical use.

The chemical name for betamethasone dipropionate is 9-fluoro-11(β), 17, 21-trihydroxy-16(β)-methylpregna-1,4-diene-3,20-dione-17,21-dipropionate. The empirical formula is $C_{28}H_{37}FO_7$ and the molecular weight is 504.6. The structural formula is shown below.



Each gram of betamethasone dipropionate spray contains 0.643 mg of betamethasone dipropionate USP (equivalent to 0.5 mg betamethasone) in a slightly thickened, white to off-white, oil-in-water, non-sterile emulsion with the following inactive ingredients: butylated hydroxytoluene, cetostearyl alcohol, diazolidinyl urea, hydroxyethyl cellulose, methylparaben, mineral oil, oleyl alcohol, polyoxyl 20 cetostearyl ether, propylparaben, purified water, sorbitan monostearate. Betamethasone dipropionate spray is co-packaged with a manual spray pump for installation by the pharmacist prior to dispensing to patients.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Corticosteroids play a role in cellular signaling, immune function, inflammation, and protein regulation; however, the precise mechanism of action of betamethasone dipropionate spray in psoriasis is unknown.

12.2 Pharmacodynamics

Vasoconstrictor studies performed with betamethasone dipropionate spray in healthy subjects indicate that it is in the mid-range of potency as compared with other topical corticosteroids; however, similar blanching scores do not necessarily imply therapeutic equivalence.

The potential for HPA axis suppression by betamethasone dipropionate spray was evaluated in a study randomizing 52 adult subjects with moderate to severe plaque psoriasis. Betamethasone dipropionate spray was applied twice daily for 15 or 29 days, in subjects with psoriasis involving a mean of 29.0% and 26.5% body surface area at baseline across the 2 treatment duration arms, respectively. Forty-eight (48) subjects were evaluated for HPA axis suppression at the end of treatment. The proportion of subjects demonstrating HPA axis suppression was 20.8% (5 out of 24) in subjects treated with betamethasone dipropionate spray for 15 days. No subjects (0 out of 24) treated with betamethasone dipropionate spray for 29 days had HPA axis suppression.

In this study HPA axis suppression was defined as serum cortisol level ≤ 18 mcg/dL 30-minutes post-cosyntropin stimulation. In the 4 subjects with available follow-up values, all subjects had normal ACTH stimulation tests at follow-up.

12.3 Pharmacokinetics

The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids are absorbed through normal intact skin. Inflammation and/or other disease processes in the skin may increase percutaneous absorption.

Plasma concentrations of betamethasone dipropionate, betamethasone-17-propionate, and betamethasone were measured at baseline, and before and after the last dose (1, 3, and 6 hours) in the HPA axis suppression trial in subjects with psoriasis [see *Clinical Pharmacology* (12.2)]. The majority of subjects had no measurable plasma concentration (<5.00 pg/mL) of betamethasone dipropionate, while the metabolites, betamethasone-17-propionate and betamethasone, were detected in the majority of subjects (Table 2). There was high variability in the data but there was a trend toward higher systemic exposure at Day 15 and lower systemic exposure at Day 29.

Table 2: Mean (\pm SD) Maximum Plasma Concentrations (pg/mL) of Betamethasone Dipropionate Metabolites after 15 or 29 Days of Treatment with Betamethasone Dipropionate Spray

Analyte (pg/mL)	Betamethasone Dipropionate Spray <i>b.i.d.</i> (15 days)	Betamethasone Dipropionate Spray <i>b.i.d.</i> (29 days)
Betamethasone-17-propionate	120 \pm 127	63.9 \pm 52.6
Betamethasone	119 \pm 176	57.6 \pm 55.9

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic potential of betamethasone dipropionate.

In a 90-day repeat-dose toxicity study in rats, topical administration of betamethasone dipropionate spray formulation at dose concentrations of 0.05% and 0.1% (providing dose levels up to 0.5 mg/kg/day in males and 0.25 mg/kg/day in females) resulted in a toxicity profile consistent with long-term exposure to corticosteroids including reduced body weight gain, adrenal atrophy, and histological changes in bone marrow, thymus and spleen indicative of severe immune suppression. A no observable adverse effect level (NOAEL) could not be determined in this study. Although the clinical relevance of the findings in animals to humans is not clear, sustained glucocorticoid-related immune suppression may increase the risk of infection and possibly the risk of carcinogenesis.

Betamethasone was negative in the bacterial mutagenicity assay (*Salmonella typhimurium* and *Escherichia coli*), and in the mammalian cell mutagenicity assay (CHO/HGPRT). It was positive in the *in vitro* human lymphocyte chromosome aberration assay, and equivocal in the *in vivo* mouse bone marrow micronucleus assay.

Studies in rabbits, mice, and rats using intramuscular doses up to 1 mg/kg, 33 mg/kg, and 2 mg/kg, respectively, resulted in dose-related increases in fetal resorptions in rabbits and mice.

14 CLINICAL STUDIES

Two multi-center, randomized, double-blind, vehicle-controlled clinical trials were conducted in subjects aged 18 years and older with moderate plaque psoriasis. In both trials, randomized subjects applied betamethasone dipropionate spray or vehicle spray to the affected areas twice daily for 28 days. Enrolled subjects had body surface area of involvement between 10% to 20%, and an Investigator Global Assessment (IGA) score of 3 (moderate).

Efficacy was assessed as the proportion of subjects who were considered a treatment success (defined as having an IGA score of 0 or 1 [clear or almost clear] and at least a 2-grade reduction from baseline). Table 3 presents the efficacy results at Day 15 and Day 29.

Table 3: Proportion of Subjects with Plaque Psoriasis with Treatment Success *after 14 Days and 28 Days of Treatment

	Study 1		Study 2	
	Betamethasone Dipropionate Spray <i>b.i.d.</i> (N=182)	Vehicle Spray <i>b.i.d.</i> (N=95)	Betamethasone Dipropionate Spray <i>b.i.d.</i> (N=174)	Vehicle Spray <i>b.i.d.</i> (N=87)
Treatment Success at Day 15	21.5%	7.4%	19.0%	2.3%
Treatment Success at Day 29	42.7%	11.7%	34.5%	13.6%

* Treatment success is defined as an IGA of 0 or 1 (clear or almost clear) and at least a 2-grade reduction from baseline.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied/Storage

Betamethasone dipropionate spray, 0.05% is a slightly thickened, white to off-white, non-sterile emulsion supplied in high density polyethylene bottles with a heat induction seal lined polypropylene cap. The drug is supplied in the following volumes:

- 60 mL
- 120 mL
- NDC 51672-1390-4
- NDC 51672-1390-8

Store at 20° to 25°C (68° to 77°F)[see USP Controlled Room Temperature].

Each unit is co-packaged with a manual spray pump for installation by the pharmacist prior to dispensing.

16.2 Handling/Instructions for the Pharmacist

1. Remove the spray pump from the wrapper.
2. Remove and discard the cap from the bottle.
3. Keeping the bottle upright, insert the spray pump into the bottle and turn clockwise until it is closed tightly.
4. Dispense the bottle with the spray pump inserted.
5. Include the date dispensed in the space provided on the carton.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Inform patients of the following:

- Discontinue therapy when control is achieved, unless directed otherwise by the physician.
- Do not use for longer than 4 consecutive weeks.
- Avoid contact with the eyes.
- Avoid use of betamethasone dipropionate spray on the face, scalp, underarms, groin or other intertriginous areas, unless directed by the physician.
- Do not occlude the treatment area with bandage or other covering, unless directed by the physician.
- Local reactions and skin atrophy are more likely to occur with occlusive use, prolonged use, or use of higher potency corticosteroids.
- Advise a woman to use betamethasone dipropionate spray on the smallest area of skin and for the shortest duration possible while pregnant or breastfeeding. Advise breastfeeding women not to apply betamethasone dipropionate spray directly to the nipple and areola to avoid direct infant exposure.

Mfd. by: Taro Pharmaceuticals Inc., Brampton, Ontario, Canada L6T 1C1

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**, Hawthorne, NY 10532

Revised: June 2020

PPK-8555-0 21

PATIENT INFORMATION **Betamethasone Dipropionate**

(bay'' ta meth' a sone dye proe' pee oh nate) Spray, 0.05%

Important: Betamethasone dipropionate spray is for use on the skin only. Do not get betamethasone dipropionate spray near or in your eyes, mouth, or vagina.

What is betamethasone dipropionate spray?

- Betamethasone dipropionate spray is a prescription corticosteroid medicine used to treat mild to moderate plaque psoriasis in people 18 years of age and older.

It is not known if betamethasone dipropionate spray is safe and effective in children under 18 years of age. Betamethasone dipropionate spray is not recommended for use in patients under 18 years of age.

Before you use betamethasone dipropionate spray, tell your doctor about all of your medical conditions, including if you:

- are allergic to any of the ingredients in betamethasone dipropionate spray. See the end of this leaflet for a list of the ingredients in betamethasone dipropionate spray.
- have thinning of the skin (atrophy) at the treatment site.
- are pregnant or plan to become pregnant. It is not known if betamethasone dipropionate spray will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if betamethasone dipropionate spray passes into breast milk.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your doctor if you take other corticosteroid medicines by mouth or use other products on your skin that contain corticosteroids.

How should I use betamethasone dipropionate spray?

See the " Instructions for Use" for detailed information about the right way to apply betamethasone dipropionate spray.

- Use betamethasone dipropionate spray exactly as your doctor tells you to use it.
- Your doctor should tell you how much betamethasone dipropionate spray to use and where to apply it.
- Apply betamethasone dipropionate spray 2 times a day.
- Use betamethasone dipropionate spray for the shortest amount of time needed to treat your plaque psoriasis. Tell your doctor if your skin condition is not getting better after 4 weeks of using betamethasone dipropionate spray. Do not use betamethasone dipropionate spray for longer than 4 weeks.
- Wash your hands after applying betamethasone dipropionate spray.
- Do not use betamethasone dipropionate spray on your face, scalp, underarms (armpits), groin, or areas where your skin may touch or rub together.
- Do not bandage, cover, or wrap the treated skin area, unless your doctor tells you to.

What are the possible side effects of betamethasone dipropionate spray?

- **Betamethasone dipropionate spray can pass through your skin.** Too much betamethasone dipropionate spray passing through your skin can cause your adrenal glands to stop working. Your doctor may do blood tests to check for adrenal gland problems.

The most common side effects of betamethasone dipropionate spray include itching, burning, stinging, pain, and thinning of skin (atrophy) at the treated site. These are not all the possible side effects of betamethasone dipropionate spray. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store betamethasone dipropionate spray?

- Store betamethasone dipropionate spray at room temperature between 68°F to 77°F (20°C to 25°C).
- Throw away (discard) any unused betamethasone dipropionate spray after 4 weeks.

Keep betamethasone dipropionate spray and all medicines out of the reach of children.

General information about the safe and effective use of betamethasone dipropionate spray.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use betamethasone dipropionate spray for a condition for which it was not prescribed. Do not give betamethasone dipropionate spray to other people even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or doctor for information about betamethasone dipropionate spray that is written for health professionals.

What are the ingredients in betamethasone dipropionate spray?

Active ingredient: betamethasone dipropionate

Inactive ingredients: butylated hydroxytoluene, cetostearyl alcohol, diazolidinyl urea, hydroxyethyl cellulose, methylparaben, mineral oil, oleyl alcohol, polyoxyl 20 cetostearyl ether, propylparaben, purified water, sorbitan monostearate.

Mfd. by: Taro Pharmaceuticals Inc., Brampton, Ontario, Canada L6T 1C1

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**, Hawthorne, NY 10532

This Patient Information has been approved by the U.S. Food and Drug Administration
Revised: June 2020

Instructions for Use

Betamethasone Dipropionate

(bay'' ta meth' a sone dye proe' pee oh nate) Spray, 0.05%

Important: Betamethasone dipropionate spray is for use on the skin only. Do not get betamethasone dipropionate spray near or in your eyes, mouth, or vagina.

Read this "Instructions for Use" before you start using betamethasone dipropionate spray and each time you get a refill. There may be new information. This information does not take the place of talking with your doctor about your medical condition or treatment.

Parts of the betamethasone dipropionate spray bottle. (See Figure A)

Figure A



How to apply betamethasone dipropionate spray:

Step 1: Shake the betamethasone dipropionate spray bottle well. Remove the cap from the pump top.

Step 2: Hold the bottle in an upright position while pointing the opening of the pump top in the direction of the affected area. To spray, push down on the pump top. Apply betamethasone dipropionate spray to the affected area as instructed by your doctor. (**See Figure B**)

Figure B



Step 3: Spray only enough betamethasone dipropionate spray to cover the affected area, for example, the elbow (**See Figure C**). Rub in betamethasone dipropionate spray gently.

Figure C



Repeat Steps 2 and 3 to apply betamethasone dipropionate spray to other affected areas as instructed by your doctor.

Step 4: After applying betamethasone dipropionate spray, place the cap back onto the pump top. (**See Figure D**)

Figure D



How should I store betamethasone dipropionate spray?

- Store betamethasone dipropionate spray at room temperature between 68°F to 77°F (20°C to 25°C).
- Throw away (discard) any unused betamethasone dipropionate spray after 28 days.

Keep betamethasone dipropionate spray and all medicines out of the reach of children.

This "Instructions for Use" has been approved by the U.S. Food and Drug Administration.

Mfd. by: Taro Pharmaceuticals Inc., Brampton, Ontario, Canada L6T 1C1

Dist. by: **Taro Pharmaceuticals U.S.A., Inc.**, Hawthorne, NY 10532

Revised: June 2020

PPK-8555-0 21

PRINCIPAL DISPLAY PANEL - 60 mL Bottle Carton**60 mL****NDC 51672-1390-4****Betamethasone****Dipropionate****Spray****0.05%****Potency expressed
as betamethasone**

FOR TOPICAL USE ONLY
NOT FOR ORAL, OPHTHALMIC
OR INTRAVAGINAL USE

**Keep this and all
medications out of the
reach of children.****Rx only****TARO**



BETAMETHASONE DIPROPIONATE

betamethasone dipropionate spray

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:51672-1390
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of	Strength
-----------------	----------	----------

Ingredient Name		Strength	Strength	
BETAMETHASONE DIPROPIONATE (UNII: 826Y60901U) (BETAMETHASONE - UNII:9842X06Q6M)		BETAMETHASONE	0.5 mg in 1 g	
Inactive Ingredients				
Ingredient Name			Strength	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)				
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)				
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)				
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)				
METHYLPARABEN (UNII: A2I8C7HI9T)				
MINERAL OIL (UNII: T5L8T28FGP)				
OLEYL ALCOHOL (UNII: 172F2WN8DV)				
POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)				
PROPYLPARABEN (UNII: Z8IX2SC1OH)				
WATER (UNII: 059QF0KO0R)				
SORBITAN MONOSTEARATE (UNII: NVZ4I0H58X)				
Product Characteristics				
Color	white (white to off-white)	Score		
Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51672-1390-4	1 in 1 CARTON	06/17/2020	
1		60 g in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
2	NDC:51672-1390-8	1 in 1 CARTON	06/17/2020	
2		120 g in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA211722	06/17/2020	

Labeler - Sun Pharmaceutical Industries, Inc. (146974886)

Establishment

Name	Address	ID/FEI	Business Operations
Sun Pharma Canada Inc.		243339023	manufacture(51672-1390)

Revised: 7/2025

Sun Pharmaceutical Industries, Inc.