RIGHT REMEDIES PAIN RELIEF EXTRA STRENGTH CAPLETacetaminophen tablet, coated Strive Pharmaceuticals Inc.

RIGHT REMEDIES Pain Relief EXTRA STRENGTH Caplet

Drug Facts

Active Ingredient (in each caplet)

Acetaminophen 500mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold headache
 - backache
 minor pain of arthritis
 - toothache muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

• skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever

gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed. See overdose warning

adult &	 take 2 caplets every 6 hours while
children	symptoms last
12 years	 do not take more than 6 caplets in 24
of age	hours unless directed by a doctor
and over	 do not take more than 10 days, unless
	directed by a doctor
children	 ask a doctor
under	
12 years	
of age	

Other information

- each caplet contains: sodium 1 mg (very low sodium)
- store at room temperature between 20 25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

Inactive ingredients

croscarmellose sodium, lactose monohydrate, magnesium stearate, polyethylene glycol, polyvinyl alcohol, povidone k30, purified water, sodium starch glycolate, starch corn, talc, titanium dioxide

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

*compare to the active ingredient of Extra Strength **Tylenol**®

TAMPER EVIDENT. DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING FROM BOTTLE

Distributed by: Strive Pharmaceuticals inc. East Brunswick, NJ 08816

Product of India

Packaged & Quality Assured in USA

READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION


*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division., owner of the registered trademark Tylenol[®] Extra Strength Caplets.

REV.00- 102022

Packaging



PACKAGE - 200 CAPLETS



Purpose Pain reliever/fever reducer

Drug Facts

Active ingredient (in each caplet)

Acetamino

Uses

- temporarily relieves minor aches and pains due to: the common cold teadache to backache minor pain of arthritis toothache
- Inscular aches premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 more than 4,000 mg of acetaminophen in 24 hours
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 skin reddening blisters rash
 fa skin reaction occurs, stop use and seek medical help right away.
 Do not use withth any other drug containing acetaminophen, ask a doctor or pharmacist.
 if you have ever had an allergic reaction to this product or any of its ingredients.
 Aks a doctor before use if you have liver disease

Stop use and ask a doctor if

pain gets worse or lasts more than 10 days
 new symptoms occur
 These could be signs of a serious condition.

lays fever gets worse or lasts more than 3 days redness or swelling is present

Drug Facts (continued) Warnings (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms. Directions do not take more than directed (see overdose warning) adults & children 12 take 2 caplets every 6 hours while symptoms last vears of age and over do not take more than 6 caplets in 24 hours, unless directed by a doctor do not take for more than 10 days unless directed by a doctor children under 12 ask a doctor years of age Other information each caplet contains: sodium 1 mg (very low sodium) avoid excessive heat, cold and humidity. Close cap tightly after use. store between 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, lactose monohydrate, magnesium stearate, polethylene glycol, polyvinyl alcohol, povidone K30, purified water, sodium starch glycolate, starch corn, talc, titanium dioxide

Questions or comments? 1-888-577-8033 Monday - Friday Bam - 4pm EST

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816 Product of India

Packaged and Quality Assured in the USA

ADHESIVE

PACKAGE - 40 CAPLETS



PACKAGE - 300 CAPLETS

Compare to the active ingredient in Extra Strength Tylenol". NDC 70692-150-31 EXTRA STRENGTH Pain Relief Acetaminophen 500 mg	Purpose Pain releventever reducer	us to: minor pain of athrits	Serere her damage may occur if you take Serere her damage may occur if you take product aactors. Symphons may include: e skin redening aastors and sek medical help right ang.	er wurd wir per foung vandenning accentinghenen (prescupion of merkeschperon in zweiter war aus auf der gentrals seterlinichen, sis auchter of pfarmeistig = / you are allengt, fo acterninghen or any of eingredients in his product.	nex userus e premierar avoire ere r year eu cump de concernant y ang en nume. Stop use and ask dender if a poil pais verse or bists more than 10 days . First pais verse or bists more Than 3 days are may anytoms cour in chriss or sveling is present. These outdide spins da serious condition	I pregnant or heast-beeling, ask a health professional before use. Geep out or reeator ideator. Devendos artificants in cases of evendose, pat medical help or contract a Poisson Control Center right away. Ouck medical alterntion is critical for adults as well as for children even if you do not notice any signs or symptoms.	0	take 2 caplets every 6 hours while symptoms last to ond take more than 6 caplets in 24 hours unless directed by a doord take for more than 10 days, unless directed by a doord	ask a doctor	 store at room temperature between 20-25°C (68-77°F) ptightly after use 	l <i>inactive i ingredientis</i> croscameloos sodum, laciose morotyviciais, magnesium stearate, polyethylene glycol, polyvinyi aicoriot, povidone K30, puntiled waler, sodium starch gyrobite, starch com, tac, ittanium dioxide	Monday-Friday 8am-4pm EST	y Johnson & Johnson Consumer Inc., McNeil istered trademark Tylenole Extra Strength Diserburded hos	Shine Pharmosultish Inc. East Brunswick, NJ 08816 Product of India Packaged and quality assured in the USA	LB7069215031 LOT: No Vamish REV.00-072023 EXP: No Vamish	·
Pain reliever Fever reducer	Drug Facts Active ingredient (in each caplet) Actimiconen 500 mg	Uses = Temporarily relieves minor aches and pains due to: the common cod' = headache = bookache = minor pain of athrhis = toothache = muscular aches = premenstrual and menstrual camps	Warnings Warnings Uver wardings: This product contains acetaminophen. Severe liker damage may occur if a more than 4,000 mg datarimophen in Nulle using this pooluct a or more alcoholic dimis servi val wille using this pooluct Allergy alter A coleminophen may cause severe skin reactions. Symphone may a billeris a rash. It a bin ackdon occurs, stop use and sever metical help right aver- and the severe and the occurs. Stop use and sever metical help right aver-	We must be must be used containing examinationer presentation or induces the method of contains actaininghen ask dock or planmacks = 1 you are allergic to acids the inactive ingredients in this product. Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before use 1 you have liver (Sease Ask a obtain before the liver (Sease Ask a obt	Stop use and ask a doctor if a pain gets worse or tast than 3 days a new symptoms occur a redness or sw	If pregnant or heast-leading, ask a health professional before use Reep out of reach of childing, ask a neglical help or contact Overdoes warming: In case of overdoes, gat medical help or contact medical attention is oritical for adults as well as for children even if	Directions Do not take more than directed. See overdose warning	e and over	children under 12 years of age	Other information = each captet contains: sodium 1 mg (very low sodium) = avoid eccessive heat, cold and humdity = close captign	Inactive ingredients croscarmellose sodum, lacto polyvinyl alcohol, povidone K30, purified water, sodium s	Questions or comments? 1-888-577-8033 Monday-Friday 8am-4pm EST	This product is not manufactured or distributed by Consumer Healthcare Division, owner of the regis		3 70692 00082 5	i

RIGHT REMEDIES PAIN RELIEF EXTRA STRENGTH CAPLET

acetaminophen tablet, coated

Product Information						
Product Type	duct Type HUMAN OTC DRUG Item Code (Source) ND					
Route of Administration	ORAL					
Active Ingredient/Active	Moioty					
	-					
Ingr	redient Name			Basis of St	rength	Strength
ACETAMINOPHEN (UNII: 36209IT	L9D) (ACETAMINOF	PHEN - UNI	l:362O9ITL9D)	ACETAMINOPH	EN	500 mg
Inactive Ingredients						
	Ingredient	Name			S	trength
CROSCARMELLOSE SODIUM (UN	NII: M28OL1HH48)					
LACTOSE MONOHYDRATE (UNII:	EWQ57Q8I5X)					
MAGNESIUM STEARATE (UNII: 70	0097M6I30)					
POLYETHYLENE GLYCOL, UNSP	ECIFIED (UNII: 3V	VJQOS DW1A	()			
POLYVINYL ALCOHOL, UNSPEC	IFIED (UNII: 532B5	59J990)				
POVIDONE K30 (UNII: U725QWY3	2X)					
WATER (UNII: 059QF0KO0R)						
SODIUM STARCH GLYCOLATE T	YPE A (UNII: H8A)	/0SQX4D)				
STARCH, CORN (UNII: 08232NY35	5J)					
TALC (UNII: 7SEV7J4R1U)						
TITANIUM DIOXIDE (UNII: 15FIX9)	V2JP)					
Product Characteristics						
Color	to	Score		n	o score	

Color	white	Score	no score
Shape	capsule	Size	18mm
Flavor		Imprint Code	S150
Contains			

Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692- 150-05	50 in 1 PACKAGE; Type 0: Not a Combination Product	07/20/2022	
2	NDC:70692- 150-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	07/20/2022	
3	NDC:70692- 150-44	1 in 1 CARTON	11/21/2023	
3		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:70692- 150-31	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/21/2023	

Marketing Information

Marketing

Application Number or Monograph Marketing Start Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M013	07/20/2022	

Labeler - Strive Pharmaceuticals Inc. (080028013)

Revised: 11/2023

Strive Pharmaceuticals Inc.