

**CLINIMIX- leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection**  
**CLINIMIX- leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection**  
**CLINIMIX- leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection**  
**Baxter Healthcare Corporation**

-----

#### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use CLINIMIX safely and effectively. See full prescribing information for CLINIMIX.**

#### **CLINIMIX (amino acids in dextrose) injection, for intravenous use**

**Initial U.S. Approval: 1997**

#### ----- **INDICATIONS AND USAGE** -----

CLINIMIX is indicated as a source of calories and protein for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. CLINIMIX may be used to treat negative nitrogen balance in patients. (1)

#### ----- **DOSAGE AND ADMINISTRATION** -----

See full prescribing information for information on preparation, administration, instructions for use, dosing considerations, including the recommended dosage in adults and pediatrics, and dosage modifications in patients with renal impairment. (2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8)

#### ----- **DOSAGE FORMS AND STRENGTHS** -----

CLINIMIX is available in multiple strengths. See full prescribing information for detailed description of each formulation. (3, 11)

#### ----- **CONTRAINDICATIONS** -----

- Known hypersensitivity to one or more amino acids or dextrose. (4)
- Inborn errors of amino acid metabolism. (4)
- Patients with pulmonary edema or acidosis due to low cardiac output. (4)

#### ----- **WARNINGS AND PRECAUTIONS** -----

- Pulmonary Embolism due to Pulmonary Vascular Precipitates: if signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. (5.1)
- Hypersensitivity Reactions: monitor for signs and symptoms and discontinue infusion if reactions occur. (5.2)
- Risk of Infections, Refeeding Complications, and Hyperglycemia or Hyperosmolar Hyperglycemic State: monitor for signs and symptoms; monitor laboratory parameters. (5.3, 5.4, 5.5)
- Vein Damage and Thrombosis: solutions with osmolarity of  $\geq 900$  mOsm/L must be infused through a central catheter. (2.2, 5.6)
- Hepatobiliary Disorders: monitor liver function parameters and ammonia levels. (5.7)
- Aluminum Toxicity: increased risk in patients with renal impairment, including preterm infants. (5.8, 8.4)
- Parenteral Nutrition Associated Liver Disease: increased risk in patients who receive parenteral nutrition for extended periods of time, especially preterm infants; monitor liver function tests, if abnormalities occur consider discontinuation or dosage reduction. (5.9, 8.4)
- Electrolyte Imbalance and Fluid Overload: patients with cardiac insufficiency or renal impairment may require adjustment of fluid, protein and electrolyte content. (5.10, 8.4)

#### ----- **ADVERSE REACTIONS** -----

Adverse reactions include diuresis, extravasation, glycosuria, hyperglycemia, and hyperosmolar coma. (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare at 1-866-888-2472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

#### ----- **USE IN SPECIFIC POPULATIONS** -----

Pediatric Use: increased risk of hypoglycemia/hyperglycemia: monitor serum glucose concentrations. (8.4)

**See 17 for PATIENT COUNSELING INFORMATION.**

**Revised: 10/2018**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Preparation Prior to Administration
- 2.2 Important Administration Instructions
- 2.3 Instructions for Use
- 2.4 Preparation and Addition of Lipid Emulsion
- 2.5 Dosing Considerations
- 2.6 Recommended Dosage in Adults
- 2.7 Dosage Modifications in Patients with Renal Impairment
- 2.8 Recommended Dosage in Pediatric Patients
- 2.9 Discontinuation of CLINIMIX

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates
- 5.2 Hypersensitivity Reactions
- 5.3 Risk of Infections
- 5.4 Refeeding Syndrome
- 5.5 Hyperglycemia or Hyperosmolar Hyperglycemic State
- 5.6 Vein Damage and Thrombosis
- 5.7 Hepatobiliary Disorders
- 5.8 Aluminum Toxicity
- 5.9 Risk of Parenteral Nutrition Associated Liver Disease
- 5.10 Electrolyte Imbalance and Fluid Overload
- 5.11 Monitoring/Laboratory Tests

### **6 ADVERSE REACTIONS**

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

### **15 REFERENCES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

CLINIMIX is indicated as a source of calories and protein for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. CLINIMIX may be used to treat negative nitrogen balance in patients.

## 2 DOSAGE AND ADMINISTRATION

### 2.1 Preparation Prior to Administration

- Tear protective clear overwrap across top at slit and remove solution container. Small amounts of moisture may be found on the solution container from water permeating from inside the container. The amount of permeated water is insufficient to affect the solution significantly. If larger amounts of water are found, the container should be checked for tears or leaks.
- Inspect the container prior to activation. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Evaluate the following:
  - If the outlet or additive port protectors are damaged, detached, or not present, discard container as solution path sterility may be impaired.
  - Check to ensure seal between chambers is intact, solutions are contained in separate chambers, and the content of the individual chambers is clear, colorless or slightly yellow. Discard if the seal is broken or if the solution is bright yellow or yellowish brown.
  - Check for minute leaks by separately squeezing each chamber. If external leaks or leakage between the chambers are found, discard solution as sterility or stability may be impaired.
- Lipids and/or additives can be introduced to the container after opening seal between chambers. Because additives may be incompatible, evaluate all additions to the plastic container for compatibility. Activate chambers of container prior to introduction of additives. Mix thoroughly when additives have been introduced. Supplemental medication may be added with a 19 to 22 gauge needle through the medication port.
- Calcium and phosphate ratios must be considered. Excess addition of calcium and phosphate, especially in the form of mineral salts, may result in the formation of calcium phosphate precipitates [see *Warnings and Precautions (5.1)*].
- Inspect the container to ensure precipitates have not formed during the mixing or addition of additives. A slight yellow color does not alter the quality and efficacy of this product. If lipid has been added, ensure the emulsion has not separated. Separation of the emulsion can be visibly identified by a yellowish streaking or the accumulation of yellowish droplets in the mixed emulsion. Discard the admixture if any of the above are observed

### 2.2 Important Administration Instructions

- Set the vent to the closed position on a vented intravenous administration set to prevent air embolism.
- Use a dedicated line without any connections to avoid air embolism.
- CLINIMIX is for intravenous infusion only into a central or peripheral vein. The choice of a central or peripheral venous route should depend on the osmolarity of the final infusate. Solutions with osmolarity of 900 mOsm/L or greater must be infused through a central catheter [see *Warnings and Precautions (5.6)*].
  - For central vein infusion only: CLINIMIX 4.25/10, 5/15, 5/20
  - For central or peripheral vein infusion: CLINIMIX 4.25/5
- The solution should be inspected for precipitates before admixing, after admixing, and again before administration.
- Use a 0.22 micron filter for administration of CLINIMIX. If a lipid is also administered, use a 1.2 micron filter.
- If lipid emulsion is added, do not use administration sets and lines that contain di-2-ethylhexyl

phthalate (DEHP). Administration sets that contain polyvinyl chloride (PVC) components have DEHP as a plasticizer.

### 2.3 Instructions for Use


1. Open by tearing protective clear overwrap, which includes an oxygen-absorbing sachet, across top at slit and remove solution container. Discard the oxygen-absorbing sachet after removal from the clear overwrap.
2. To proceed with activation, the container should be at room temperature. Lay the room temperature container onto a flat surface. Grasp the container firmly on each side of the top of the container (**Figure 1**).
3. Starting from the top, using some pressure, slowly roll the container to open seal between chambers as shown in **Figure 2**. Do not pull or rip the seal apart. The seal must be completely opened towards the port side of the container. The upper section of the seal towards the hanger side can remain unbroken.
4. Mix the contents thoroughly by inverting the container upside down to ensure a homogenous admixture (**Figure 3**).
5. Once the container is mixed, check for leaks.
6. Make additions (if prescribed).

Because additives may be incompatible, evaluate all additions to the container for compatibility and stability of the resulting preparation. Consult with pharmacist, if available. Questions about compatibility may be directed to Baxter. If it is deemed advisable to introduce additives, use aseptic technique. For information on adding lipid emulsions *see Dosage and Administration (2.4)*.

- a. Prepare medication port.
  - b. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
  - c. Mix solution and medication thoroughly (**Figure 3**). For high density medication (high specific gravity), such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.
7. Inspect final solution for discoloration and particulate matter. Check for leaks.
  8. Spike and hang container.
    - d. Suspend container from eyelet support.
    - e. Twist off protector from outlet port at bottom of container (**Figure 4**).
    - f. Attach administration set. Refer to complete directions accompanying set.

For single dose only. Discard unused portion.

**Figures 1–4:**



### Instructions on Storage

#### *Storage After Removal of Overwrap:*

Once removed from the protective clear overwrap, mixed (peel seal activated) or unmixed (peel seal intact) CLINIMIX solutions may be stored under refrigeration for up to 9 days.

#### *Storage Once any Additive is Added:*

Use promptly after mixing. Any storage with additives should be under refrigeration and limited to a brief period of time, less than 24 hours. After removal from refrigeration, use promptly and complete the infusion within 24 hours. Any remaining mixture must be discarded.

### **2.4 Preparation and Addition of Lipid Emulsion**

1. Prior to adding lipid emulsion, mix amino acid and dextrose injection as shown in **Figures 1-3**.
2. Prepare lipid emulsion transfer set following instructions provided.
3. Attach transfer set to lipid emulsion container using aseptic technique.
4. Twist off protector on the additive port of the container.
5. Attach the transfer set to the exposed additive port.
6. Open clamp on transfer set.
7. After completing transfer, use appropriate plastic clamp or metal ferrule to seal off additive port tube.
8. Remove transfer set.
9. Mix contents of container thoroughly. Inspect final solution for discoloration and particulate matter. Check for leaks.

#### *Storage Once Lipids are Added:*

Use promptly after mixing. Any storage with additives should be under refrigeration and limited to a brief period of time, no longer than 24 hours. After removal from refrigeration, use promptly and complete the infusion within 24 hours. Any mixture remaining must be discarded.

## 2.5 Dosing Considerations

- The dosage of CLINIMIX should be individualized based on the patient’s clinical condition (ability to adequately metabolize amino acids and dextrose), body weight and nutritional/fluid requirements, as well as additional energy given orally/enterally to the patient. Prior to initiating CLINIMIX the following patient information should be reviewed: all concomitant medications, gastrointestinal function and laboratory data such as electrolytes (including magnesium, calcium, and phosphorus), glucose, urea/creatinine, liver panel, complete blood count and triglyceride level (if adding lipid emulsion). Refer to the complete prescribing information of lipid emulsion for dosing information.
- CLINIMIX formulations have varying concentrations of protein and carbohydrate; thus infusion rates to achieve requirements will vary. Protein, caloric, fluid and electrolyte requirements all need to be taken into consideration when determining individual patient dosage needs.
- The dosage selection is based only on the recommended protein requirements. The maximum dextrose infusion rates and calorie and fluid requirements must also be considered when determining the clinically appropriate infusion rate for patients.
- CLINIMIX meets the total nutritional requirements for protein and dextrose in stable patients, and can be individualized to meet specific needs with the addition of nutrients.
- Total daily fluid requirements can be met beyond the volume of amino acids solution by supplementing with non-carbohydrate or carbohydrate-containing electrolyte solutions. In many patients, provision of adequate calories in the form of hypertonic dextrose may require the administration of exogenous insulin to prevent hyperglycemia and glycosuria.
- Prior to administration of CLINIMIX correct severe fluid, electrolyte and acid-base disorders.
- Monitor levels of serum potassium during therapy. It may be necessary to add potassium to the CLINIMIX admixture.
- Lipid emulsion administration should be considered with prolonged use (more than 5 days) of CLINIMIX in order to prevent essential fatty acid deficiency (EFAD). Serum lipids should be monitored for evidence of EFAD in patients maintained on fat-free parenteral nutrition. See prescribing information of lipid emulsion.
- The flow rate should be increased gradually. The flow rate must be adjusted taking into account the dose being administered, the daily volume intake, and the duration of the infusion.

## 2.6 Recommended Dosage in Adults

The recommended daily nutritional requirements for protein and dextrose compared to the amount of nutrition provided by CLINIMIX are shown in **Table 1**.

As indicated on an individual basis, maintenance vitamins, electrolytes, trace elements and other components (including lipids) should be administered as required to prevent deficiencies and complications from developing.

The maximum infusion rates in adult patients are show in Table 2.

In addition to meeting protein needs, the administration rate should be governed, especially during the first few day of therapy, by the patient’s tolerance to dextrose. Daily intake of amino acids and dextrose should be increased gradually to the maximum required dose as indicated by frequent determinations of blood glucose levels.

**Table 1: Nutritional Comparison –Adult Patients**

	<b>Recommended Nutritional</b>	<b>Recommended CLINIMIX Adult Dosage</b>
--	--------------------------------	--

	Requirements		CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
	Stable Patients	Critically Ill Patients *				
<b>Fluid</b> (mL/kg/day)	30 to 40	Minimum needed to deliver adequate nutrition	19 to 40	19 to 40	16 to 40	16 to 40
<b>Protein</b> <sup>†</sup> (g/kg/day) (Nitrogen g/kg/day)	0.8 to 1 (0.13 to 0.16)	1.5 to 2 (0.24 to 0.32)	0.8 to 1.7 (0.13 to 0.27)	0.8 to 1.7 (0.13 to 0.27)	0.8 to 2 (0.13 to 0.32)	0.8 to 2 (0.13 to 0.32)
<b>Dextrose</b> (g/kg/day)	≤10	≤5.8	0.95 to 2	1.9 to 4	2.4 to 6	3.2 to 8

\* Do not use in patients with conditions that are contraindicated [see *Contraindications (4)*].

† Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

**Table 2: Maximum Infusion Rate in Adult Patients :**

		Maximum Infusion Rates in Adults Patients			
		CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
<b>Maximum Infusion Rate (mL/kg/hour)</b>		2.4	2.4	1.67	1.25
<b>Corresponding infusion rate</b>	<b>Amino Acid (g/kg/hour)</b>	0.1*	0.1*	0.08	0.06
	<b>Dextrose (g/kg/hour)</b>	0.12	0.24	0.25*	0.25*

\* Rate limiting factor

## 2.7 Dosage Modifications in Patients with Renal Impairment

Prior to administration, correct severe fluid or electrolyte imbalances. Closely monitor serum electrolyte levels and adjust the volume of CLINIMIX administered as required [see *Warnings and Precautions (5.10)*].

Patients with renal impairment not needing dialysis require 0.6 to 0.8 g of protein/kg/day. Serum electrolyte levels should be closely monitored. Patients on hemodialysis or continuous renal replacement therapy should receive 1.2 to 1.8 g of protein/kg/day up to a maximum of 2.5 g of protein/kg/day based on nutritional status and estimated protein losses.<sup>2</sup> The CLINIMIX dosage can be adjusted based on the severity of renal impairment, supplementing protein as indicated. If required, additional amino acids may be added to the CLINIMIX container or infused separately. Compatibility of additions should be evaluated by a pharmacist and questions may be directed to Baxter.

## 2.8 Recommended Dosage in Pediatric Patients

The dosage and constant infusion rate of intravenous dextrose must be selected with caution in pediatric patients, particularly neonates and low weight infants, because of the increased risk of hyperglycemia/hypoglycemia [see *Use in Specific Populations (8.4)*]. Frequent monitoring of serum glucose concentrations is required when dextrose is prescribed to pediatric patients, particularly neonates and low birth weight infants. The infusion rate and volume should be determined by the consulting physician experienced in pediatric intravenous fluid therapy.

In pediatric patients, CLINIMIX is dosed on the basis of protein provided as amino acids. The

recommended dosage, by age group is provided in **Tables 3 - 6**. Infusion rates are based on protein and do not take carbohydrates, fluid or electrolytes into consideration.

This product does not contain the amino acids cysteine and taurine, considered conditionally essential for neonates and infants. If possible, these amino acids should be added to this product if used in this pediatric population.

**Table 3: Preterm and Term Infants Less than 1 Month of Age**

	Recommended Nutritional Requirements <sup>1</sup>	Recommended CLINIMIX Dosage in Preterm and Term Infants Less than 1 Month of Age			
		CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
<b>Infusion Rate Range</b> (mL/kg/hr)		2.9 to 3.9	2.9 to 3.9	2.5 to 3.3	2.5 to 3.3
<b>Fluid</b> (mL/kg/day)	100 to 150	70 to 94	70 to 94	60 to 79	60 to 79
<b>Protein*</b> (g/kg/day) (Nitrogen g/kg/day)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)
<b>Dextrose</b> (g/kg/day)	7 to 20	3.5 to 4.7	7 to 9.4	9 to 11.9	12 to 15.8

\* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

**Table 4: Pediatric Patients 1 Month to Less than 1 Year of Age**

	Recommended Nutritional Requirements <sup>1</sup>	Recommended CLINIMIX Dosage in Pediatric Patients 1 Month to Less than 1 Year of Age			
		CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
<b>Infusion Rate Range</b> (mL/kg/hr)		2 to 2.9	2 to 2.9	1.7 to 2.5	1.7 to 2.5
<b>Fluid</b> (mL/kg/day)	100 mL/kg for the first 10 kg + 50 mL/kg for the second 10 kg.	48 to 70	48 to 70	41 to 60	41 to 60
<b>Protein*</b> (g/kg/day) (Nitrogen g/kg/day)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)
<b>Dextrose</b> (g/kg/day)	7 to 20	2.4 to 3.5	4.8 to 7	6.1 to 9	8.2 to 12

\* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

**Table 5: Pediatric Patients 1 Year to Less than 11 Years of Age**

	Recommended Nutritional	Recommend CLINIMIX Dosage in Pediatric Patients 1 Year to Less than 11 Years of Age
--	-------------------------	---



	Requirements <sup>1</sup>	CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
<b>Infusion Rate Range</b> (mL/kg/hr)		1 to 2	1 to 2	0.8 to 1.7	0.8 to 1.7
<b>Fluid</b> (mL/kg/day)	100 mL/kg for the first 10 kg + 50 mL/kg for the second 10 kg + 20 mL/kg for weight > 20 kg	24 to 48	24 to 48	19 to 41	19 to 41
<b>Protein*</b> (g/kg/day) (Nitrogen g/kg/day)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)
<b>Dextrose</b> (g/kg/day)	7 to 14	1.2 to 2.4	2.4 to 4.8	2.9 to 6.1	3.8 to 8.2

\* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

**Table 6: Pediatric Patients 11 Years to 17 Years of Age**

	Recommended Nutritional Requirements <sup>1</sup>	Recommended CLINIMIX Dosage in Pediatric Patients 11 Years to 17 Years of Age			
		CLINIMIX 4.25/5	CLINIMIX 4.25/10	CLINIMIX 5/15	CLINIMIX 5/20
<b>Infusion Rate Range</b> (mL/kg/hr)		0.8 to 1.5	0.8 to 1.5	0.7 to 1.3	0.7 to 1.3
<b>Fluid</b> (mL/kg/day)	100 mL/kg for the first 10 kg + 50 mL/kg for the second 10 kg + 20 mL/kg for weight > 20 kg	19 to 36	19 to 36	17 to 31	17 to 31
<b>Protein*</b> (g/kg/day) (Nitrogen g/kg/day)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)
<b>Dextrose</b> (g/kg/day)	5 to 9	1 to 1.8	1.9 to 3.6	2.5 to 4.7	3.4 to 6.2

\* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

## 2.9 Discontinuation of CLINIMIX

To reduce the risk of hypoglycemia after discontinuation, a gradual decrease in flow rate in the last hour of infusion should be considered.

## 3 DOSAGE FORMS AND STRENGTHS

CLINIMIX is available in 1000 mL and 2000 mL dual chamber containers. The individual chambers contain essential and nonessential amino acids and dextrose. Table 7 describes the individual components of CLINIMIX.

**Table 7 INGREDIENTS PER 100mL OF CLINIMIX**

<b>Strength of CLINIMIX</b>		CLINIMIX 4.25/5 sulfite-free (4.25% Amino Acid in 5% Dextrose) Injection	CLINIMIX 4.25/10 sulfite-free (4.25% Amino Acid in 10% Dextrose) Injection	CLINIMIX 5/15 sulfite-free (5% Amino Acid in 15% Dextrose) Injection	CLINIMIX 5/20 sulfite-free (5% Amino Acid in 20% Dextrose) Injection
	Dextrose Hydrous, USP (g/100 mL)	5	10	15	20
	Amino Acids (g/100 mL)	4.25	4.25	5	5
	Total Nitrogen (mg/100 mL)	702	702	826	826
<b>Essential Amino Acids (mg/100 mL)</b>	Leucine	311	311	365	365
	Isoleucine	255	255	300	300
	Valine	247	247	290	290
	Lysine (added as the hydrochloride salt)	247	247	290	290
	Phenylalanine	238	238	280	280
	Histidine	204	204	240	240
	Threonine	179	179	210	210
	Methionine	170	170	200	200
	Tryptophan	77	77	90	90
<b>Nonessential Amino Acids (mg/100 mL)</b>	Alanine	880	880	1035	1035
	Arginine	489	489	575	575
	Glycine	438	438	515	515
	Proline	289	289	340	340
	Serine	213	213	250	250
	Tyrosine	17	17	20	20
<b>Anion Profile (mEq/L)*</b>	Acetate <sup>†</sup>	37	37	42	42
	Chloride <sup>‡</sup>	17	17	20	20
	pH <sup>§</sup> (Range)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)
	Osmolarity (mOsmol/L) (calc)	675	930	1255	1505
<b>Caloric Content (kcal/L)</b>	From Dextrose	170	340	510	680
	From Amino Acids	170	170	200	200
	TOTAL (Dextrose and Amino Acids)	340	510	710	880

\* Balanced by ions from amino acids.

† Derived from glacial acetic acid (for pH adjustment) and sodium acetate.

‡ Contributed by calcium chloride, lysine hydrochloride, magnesium chloride, and sodium chloride.

§ pH of sulfite-free amino acid injection in the outlet port chamber was adjusted with glacial acetic acid.

## 4 CONTRAINDICATIONS

The use of CLINIMIX is contraindicated in:

- Patients with known hypersensitivity to one or more amino acids or dextrose [*see Warnings and Precautions (5.2)*].
- Patients with inborn errors of amino acid metabolism due to risk of severe metabolic and neurologic complications.
- Patients with pulmonary edema or acidosis due to low cardiac output.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates

Pulmonary vascular precipitates causing pulmonary vascular emboli and pulmonary distress have been reported in patients receiving parenteral nutrition. In some cases, fatal outcomes due to pulmonary embolism have occurred. CLINIMIX contains no added phosphorus. Patients, especially those with hypophosphatemia, may require the addition of phosphate. To prevent hypocalcemia, calcium supplementation should always accompany phosphate administration. Excessive addition of calcium and phosphate increases the risk of the formation of calcium phosphate precipitates. Precipitates have been reported even in the absence of phosphate salt in the solution. Precipitation following passage through an in-line filter and suspected in vivo precipitate formation has also been reported. If signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. In addition to inspection of the solution [*see Dosage and Administration (2.1, 2.2, 2.3, 2.4)*], the infusion set and catheter should also periodically be checked for precipitates.

### 5.2 Hypersensitivity Reactions

Hypersensitivity/infusion reactions including anaphylaxis have been reported with CLINIMIX. Stop infusion immediately and treat patient accordingly if any signs or symptoms of a hypersensitivity reaction develop. Signs or symptoms may include: hypotension, hypertension, peripheral cyanosis, tachycardia, dyspnea, vomiting, nausea, urticaria, rash, pruritus, erythema, hyperhidrosis, pyrexia, and chills.

### 5.3 Risk of Infections

Patients who require parenteral nutrition are at high risk of infections because the nutritional components of these solutions can support microbial growth. Infection and sepsis may also occur as a result of the use of intravenous catheters to administer parenteral nutrition.

The risk of infection is increased in patients with malnutrition-associated immunosuppression, hyperglycemia exacerbated by dextrose infusion, long-term use and poor maintenance of intravenous catheters, or immunosuppressive effects of other concomitant conditions, drugs, or other components of the parenteral formulation (e.g., lipid emulsion).

To decrease the risk of infection, ensure aseptic technique in catheter placement and maintenance, as well as aseptic technique in the preparation and administration of the nutritional formula.

Monitor for signs and symptoms (including fever and chills) of early infections, including laboratory test results (including leukocytosis and hyperglycemia) and frequent checks of the parenteral access device and insertion site for edema, redness and discharge.

### 5.4 Refeeding Syndrome

Refeeding severely undernourished patients may result in refeeding syndrome, characterized by the intracellular shift of potassium, phosphorus, and magnesium as the patient becomes anabolic. Thiamine deficiency and fluid retention may also develop. To prevent these complications, monitor severely undernourished patients and slowly increase nutrient intakes.

### **5.5 Hyperglycemia or Hyperosmolar Hyperglycemic State**

When using CLINIMIX in patients with diabetes mellitus, impaired glucose tolerance may worsen hyperglycemia. Administration of dextrose at a rate exceeding the patient's utilization rate may lead to hyperglycemia, coma, and death. Patients with underlying confusion and renal impairment who receive dextrose infusions, may be at greater risk of developing hyperosmolar hyperglycemic state. Monitor blood glucose levels and treat hyperglycemia to maintain optimum levels while administering CLINIMIX. Insulin may be administered or adjusted to maintain optimal blood glucose levels during CLINIMIX administration.

### **5.6 Vein Damage and Thrombosis**

Solutions with osmolarity of 900 mOsm/L or greater must be infused through a central catheter. CLINIMIX solutions containing more than 5% dextrose have an osmolarity greater than or equal to 900 mOsm/L. CLINIMIX 4.25/10, 5/15 and 5/20 are indicated for administration into a central vein only, such as the superior vena cava [see *Dosage and Administration (2.2)*]. The infusion of hypertonic nutrient injections into a peripheral vein may result in vein irritation, vein damage, and/or thrombosis.

CLINIMIX 4.25/5 is indicated for peripheral administration, or may be infused into a central vein [see *Dosage and Administration (2.2)*]. The primary complication of peripheral access is venous thrombophlebitis, which manifests as pain, erythema, tenderness or a palpable cord. Remove the catheter as soon as possible, if thrombophlebitis develops.

### **5.7 Hepatobiliary Disorders**

Hepatobiliary disorders are known to develop in some patients without preexisting liver disease who receive parenteral nutrition, including cholecystitis, cholelithiasis, cholestasis, hepatic steatosis, fibrosis and cirrhosis, possibly leading to hepatic failure. The etiology of these disorders is thought to be multifactorial and may differ between patients.

Increase in blood ammonia levels and hyperammonemia may occur in patients receiving amino acid solutions. In some patients this may indicate hepatic insufficiency or the presence of an inborn error of amino acid metabolism [see *Contraindications (4)*.]

Monitor liver function parameters and ammonia levels. Patients developing signs of hepatobiliary disorders should be assessed early by a clinician knowledgeable in liver diseases in order to identify possible causative and contributory factors, and possible therapeutic and prophylactic interventions.

### **5.8 Aluminum Toxicity**

CLINIMIX contains no more than 25 mcg/L of aluminum. However, with prolonged parenteral administration in patients with renal impairment, the aluminum contained in CLINIMIX may reach toxic levels. Preterm infants are at a greater risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Patients with renal impairment, including preterm infants, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day, accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

### **5.9 Risk of Parenteral Nutrition Associated Liver Disease**

Parenteral Nutrition Associated Liver Disease (PNALD) has been reported in patients who receive parenteral nutrition for extended periods of time, especially preterm infants, and can present as cholestasis or steatohepatitis. The exact etiology is unknown and is likely multifactorial. If CLINIMIX treated patients develop liver test abnormalities consider discontinuation or dosage reduction.

## 5.10 Electrolyte Imbalance and Fluid Overload

Patients with renal impairment, such as pre-renal azotemia, renal obstruction, and protein-losing nephropathy may be at increased risk of electrolyte and fluid volume imbalance. Patients with cardiac insufficiency due to left ventricular systolic dysfunction are susceptible to excess fluid accumulation. Use CLINIMIX with caution in patients with cardiac insufficiency or renal impairment. CLINIMIX dosage may require adjustment with specific attention to fluid, protein, and electrolyte content in these patients.

Monitor renal function parameters. Patients developing signs of renal impairment should be assessed early by a clinician knowledgeable in renal disease in order to determine the appropriate CLINIMIX dosage and other treatment options.

## 5.11 Monitoring/Laboratory Tests

Monitor fluid and electrolyte status, serum osmolarity, blood glucose, liver and kidney function, blood count and coagulation parameters throughout treatment.

Patients receiving CLINIMIX should be monitored frequently and their electrolyte requirements individualized.

## 6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the prescribing information.

- Pulmonary embolism due to pulmonary vascular precipitates [*see Warnings and Precautions (5.1)*]
- Hypersensitivity reactions [*see Warnings and Precautions (5.2)*]
- Risk of Infections [*see Warnings and Precautions (5.3)*]
- Refeeding syndrome [*see Warnings and Precautions (5.4)*]
- Hyperglycemia or hyperosmolar hyperglycemic state [*see Warnings and Precautions (5.5)*]
- Vein damage and thrombosis [*see Warnings and Precautions (5.6)*]
- Hepatobiliary disorders [*see Warnings and Precautions (5.7)*]
- Parenteral Nutrition Associated Liver Disease [*see Warnings and Precautions (5.9)*]
- Electrolyte imbalance and fluid overload [*see Warnings and Precautions (5.10)*]

The following adverse reactions from voluntary reports or clinical studies have been reported with CLINIMIX. Because many of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Diuresis
- Extravasation
- Glycosuria
- Hyperglycemia
- Hyperosmolar coma

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### *Risk Summary*

There are no adequate or well-controlled studies in pregnant women with CLINIMIX. Additionally, animal reproduction studies have not been conducted with amino acids and electrolytes and dextrose. It

is not known whether CLINIMIX can cause fetal harm when administered to a pregnant woman.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. However, the estimated background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies.

### *Clinical Considerations*

#### Disease-Associated Maternal and/or Embryo-Fetal Risk

Based on clinical practice guidelines, parenteral nutrition should be considered in cases of severe maternal malnutrition where nutritional requirements cannot be fulfilled by the enteral route because of the risks to the fetus associated with severe malnutrition, such as preterm delivery, low birth weight, intrauterine growth restriction, congenital malformations and perinatal mortality.

## **8.2 Lactation**

### *Risk Summary*

It is not known whether CLINIMIX is present in human milk. There are no data on the effects of CLINIMIX on the breastfed infant or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for CLINIMIX and any potential adverse effects on the breastfed child from CLINIMIX or from the underlying maternal condition.

## **8.4 Pediatric Use**

Safety and effectiveness of CLINIMIX in pediatric patients have not been established by adequate and well-controlled studies. Use of dextrose, amino acid infusions and electrolytes in pediatric patients is based on clinical practice [see *Dosage and Administration (2.8)*].

Newborns, especially those born premature and with low birth weight, are at increased risk of developing hypo – or hyperglycemia and therefore need close monitoring during treatment with intravenous glucose solutions to ensure adequate glycemic control in order to avoid potential long term adverse effects. Hypoglycemia in the newborn can cause prolonged seizures, coma and brain damage. Hyperglycemia has been associated with intraventricular hemorrhage, late onset bacterial and fungal infection, retinopathy of prematurity, necrotizing enterocolitis, bronchopulmonary dysplasia, prolonged length of hospital stay, and death. Plasma electrolyte concentrations should be closely monitored in the pediatric population as this population may have impaired ability to regulate fluids and electrolytes.

Because of immature renal function, preterm infants receiving prolonged treatment with CLINIMIX may be at risk of aluminum toxicity [see *Warnings and Precautions (5.8)*].

Patients, including pediatric patients, may be at risk for Parenteral Nutrition Associated Liver Disease (PNALD) [see *Warnings and Precautions (5.9)*].

Hyperammonemia is of special significance in infants (birth to two years). This reaction appears to be related to a deficiency of the urea cycle amino acids of genetic or product origin. It is essential that blood ammonia be measured frequently in infants [See *Warnings and Precautions (5.7)*].

## **8.5 Geriatric Use**

Clinical studies of CLINIMIX did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from other younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

## **10 OVERDOSAGE**

An increased infusion rate of CLINIMIX cause hyperglycemia, hyperosmolality, and adverse effects on water and electrolyte balance [see Warnings and Precautions (5.5, 5.10)].

Severe hyperglycemia and severe dilutional hyponatremia, and their complications, can be fatal.

Discontinue infusion and institute appropriate corrective measures in the event of overhydration or solute overload during therapy, with particular attention to respiratory and cardiovascular systems.

For current information on the management of poisoning or overdosage, contact the National Poison Control Center at 1-800-222-1222 or [www.poisson.org](http://www.poisson.org).

## 11 DESCRIPTION


CLINIMIX sulfite-free (amino acids in dextrose) injection for intravenous use consists of sterile, nonpyrogenic, hypertonic solutions in a dual chamber container.

The outlet port chamber contains essential and nonessential amino acids. The formulas for the individual amino acids found in CLINIMIX sulfite-free (amino acids in dextrose) injections are provided in Table 8.

**Table 8: Formulas for Amino Acids**

<b>Essential Amino Acids</b>	
Leucine	$(\text{CH}_3)_2 \text{CHCH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Isoleucine	$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3) \text{CH}(\text{NH}_2) \text{COOH}$
Valine	$(\text{CH}_3)_2 \text{CHCH}(\text{NH}_2) \text{COOH}$
Lysine (added as the hydrochloride salt)	$\text{H}_2\text{N}(\text{CH}_2)_4 \text{CH}(\text{NH}_2) \text{COOH}$
Phenylalanine	$(\text{C}_6\text{H}_5) \text{CH}_2 \text{CH}(\text{NH}_2) \text{COOH}$
Histadine	$(\text{C}_3\text{H}_3\text{N}_2) \text{CH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Threonine	$\text{CH}_3\text{CH}(\text{OH}) \text{CH}(\text{NH}_2) \text{COO}$
Methionine	$\text{CH}_3\text{S}(\text{CH}_2)_2 \text{CH}(\text{NH}_2) \text{COOH}$
Tryptophan	$(\text{C}_8\text{H}_6\text{N}) \text{CH}_2 \text{CH}(\text{NH}_2) \text{COOH}$
<b>Nonessential Amino Acids</b>	
Alanine	$\text{CH}_3\text{CH}(\text{NH}_2) \text{COOH}$
Arginine	$\text{H}_2\text{NC}(\text{NH}) \text{NH}(\text{CH}_2)_3 \text{CH}(\text{NH}_2) \text{COOH}$
Glycine	$\text{H}_2\text{NCH}_2\text{COOH}$
Proline	$[(\text{CH}_2)_3 \text{NH CH}] \text{COOH}$
Serine	$\text{HOCH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Tyrosine	$[\text{C}_6\text{H}_4(\text{OH})] \text{CH}_2\text{CH}(\text{NH}_2) \text{COOH}$

The injection port chamber contains dextrose. Dextrose, USP, is chemically designated D-glucose, monohydrate ( $\text{C}_6\text{H}_{12}\text{O}_6 \cdot \text{H}_2\text{O}$ ) and has the following structure:



## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

CLINIMIX is used as a supplement of nutrition in patients, providing macronutrients (amino acids and dextrose) parenterally.

The amino acids provide the structural units that make up proteins and are used to synthesize proteins and other biomolecules or are oxidized to urea and carbon dioxide as a source of energy.

The administered dextrose is oxidized to carbon dioxide and water, yielding energy.

### 12.3 Pharmacokinetics

The disposition of infused amino acids and dextrose, are essentially the same as those absorbed from ordinary food.

## 15 REFERENCES

1. Ayers Phil, et al. *A.S.P.E.N. Parenteral Nutrition Handbook*, 2<sup>nd</sup> ed. 2014, pg. 123.
2. Mueller CM ed. *The A.S.P.E.N. Nutrition Support Core Curriculum*, 2<sup>nd</sup> ed. 2012. Chapter 29. Wolk R, Foulks C. Renal Disease, pg. 500.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

CLINIMIX (amino acids in dextrose) injection (sulfite-free) is available in 1000 mL and 2000mL volumes (See Table 9).

**Table 9: CLINIMIX Formulations**

After mixing, the product represents	1000 mL Code and NDC Number	2000 mL Code and NDC Number
CLINIMIX 4.25/5 sulfite-free (4.25% Amino Acid in 5% Dextrose) Injection	Code 2B7726 NDC 0338-1133-03	Code 2B7704 NDC 0338-1089-04



CLINIMIX 4.25/10 sulfite-free (4.25% Amino Acid in 10% Dextrose) Injection	Code 2B7727 NDC 0338-1134-03	Code 2B7705 NDC 0338-1091-04
CLINIMIX 5/15 sulfite-free (5% Amino Acid in 15% Dextrose) Injection	Code 2B7730 NDC 0338-1137-03	Code 2B7709 NDC 0338-1099-04
CLINIMIX 5/20 sulfite-free (5% Amino Acid in 20% Dextrose) Injection	Code 2B7731 NDC 0338-1138-03	Code 2B7710 NDC 0338-1101-04

Minimize exposure of CLINIMIX to heat and avoid excessive heat.

Protect from freezing.

Store CLINIMIX at room temperature (25°C/77°F) (may briefly store at up to 40°C/104°F).

Refrigerated storage is limited to 9 days once the protective clear overwrap has been opened.

Do not use if the protective clear overwrap has been previously opened or damaged.

For storage of admixed solutions see *Dosage and Administration* (2.3, 2.4).

## 17 PATIENT COUNSELING INFORMATION

Inform patients, caregivers, or home healthcare providers of the following risks of CLINIMIX:

- Pulmonary embolism due to pulmonary vascular precipitates [see *Warnings and Precautions* (5.1)]
- Hypersensitivity reactions [see *Warnings and Precautions* (5.2)]
- Risk of Infections [see *Warnings and Precautions* (5.3)]
- Refeeding syndrome [see *Warnings and Precautions* (5.4)]
- Hyperglycemia or hyperosmolar hyperglycemic state [see *Warnings and Precautions* (5.5)]
- Vein damage and thrombosis [see *Warnings and Precautions* (5.6)]
- Hepatobiliary disorders [see *Warnings and Precautions* (5.7)]
- Aluminum toxicity [see *Warnings and Precautions* (5.8)]
- Parenteral Nutrition Associated Liver Disease (PNALD) [see *Warnings and Precautions* (5.9)]
- Electrolyte imbalance and fluid overload [see *Warnings and Precautions* (5.10)]

### **Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

Printed in USA

Baxter and Clinimix are registered trademarks of Baxter International Inc.

07-19-00-0372

## PACKAGE LABEL - PRINCIPAL DISPLAY PANEL

2B7725 NDC 0338-1132-03

# CLINIMIX 2.75/5

**SULFITE-FREE**  
**(2.75% Amino Acids  
in 5% Dextrose)**  
**Injection**

**500 mL INJECTION PORT CHAMBER**  
**10% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER**  
**5.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 5 g  
**ESSENTIAL AMINO ACIDS**  
LEUCINE 201 mg  
ISOLEUCINE 165 mg  
VALINE 160 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 159 mg  
PHENYLALANINE 154 mg  
HISTIDINE 132 mg  
THREONINE 116 mg  
METHIONINE 110 mg  
TRYPTOPHAN 50 mg

**NONESSENTIAL AMINO ACIDS**  
ALANINE 570 mg  
ARGININE 316 mg  
GLYCINE 283 mg  
PROLINE 187 mg  
SERINE 138 mg  
TYROSINE 11 mg

**mg/L**  
ACETATE 24  
CHLORIDE 11

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7728 NDC 0338-1132-03

**CLINIMIX**  
**2.75/5**  
**SULFITE-FREE**

**(2.75% Amino Acid  
in 5% Dextrose)**  
**Injection**

**500 mL INJECTION PORT CHAMBER**  
**10% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER**  
**5.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS  
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 201 mg

ISOLEUCINE 165 mg

VALINE 160 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 159 mg

PHENYLALANINE 154 mg

HISTIDINE 132 mg

THREONINE 116 mg

METHIONINE 110 mg

TRYPTOPHAN 50 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 570 mg

ARGININE 316 mg

GLYCINE 283 mg

PROLINE 187 mg

SERINE 138 mg

TYROSINE 11 mg

mEq/L

ACETATE 24

CHLORIDE 11

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

***Baxter***

**BAXTER HEALTHCARE CORPORATION**

DEERFIELD IL 60015 USA

MADE IN USA

2B7727 NDC 0338-1134-03

# CLINIMIX 4.25/10

**SULFITE-FREE**

**(4.25% Amino Acids  
in 10% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**500 mL INJECTION PORT CHAMBER  
20% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 10 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NOESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

**mEq/L**

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7727 NDC 0338-1134-03

**CLINIMIX**

**4.25/10**

**SULFITE-FREE**

**(4.25% Amino Acids  
in 10% Dextrose)  
Injection**

**500 mL INJECTION PORT CHAMBER  
20% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 10 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

***Baxter***

**BAXTER HEALTHCARE CORPORATION**

DEERFIELD IL 60015 USA

MADE IN USA

2B7701 NDC 0338-1083-04

**2.75/5**

**CLINIMIX 2.75/5**  
sulfite-free  
**(2.75% Amino Acid  
in 5% Dextrose)**  
**Injection**

**1000 mL INJECTION PORT CHAMBER**  
**10% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**5.5% Amino Acid Injection**

**Rx Only**

CHECK FOR MINUTE LEAKS BY SQUEEZING  
EACH CHAMBER OF THE BAG

BEFORE USE GRASP EACH SIDE OF  
THE TOP OF THE BAG AND ROLL BAG  
TO OPEN SEAL BETWEEN CHAMBERS

**MIX THOROUGHLY**

AFTER MIXING THE PRODUCT  
REPRESENTS 2000 mL

ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH  
PHARMACIST IF AVAILABLE WHEN INTRODUCING  
ADDITIVES USE ASEPTIC TECHNIQUE  
MIX THOROUGHLY DO NOT STORE

DOSAGE INTRAVENOUSLY AS DIRECTED BY A  
PHYSICIAN DO NOT USE UNLESS SOLUTION IS  
CLEAR REFRIGERATED STORAGE AFTER MIXING  
LIMITED TO 24 HOURS DISCARD UNUSED PORTION  
SEE ACCOMPANYING DIRECTIONS FOR USE

CAUTIONS MUST NOT BE USED IN SERIES  
CONNECTIONS DISCONTINUE INFUSION  
IF ADVERSE REACTION OCCURS

CLARITY DUAL CHAMBER CONTAINER PL 2401 PLASTIC

LOT

EXP

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION  
DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE	201 mg
ISOLEUCINE	185 mg
VALINE	180 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	159 mg
PHENYLALANINE	154 mg
HISTIDINE	132 mg
THREONINE	116 mg
METHIONINE	110 mg
TRYPTOPHAN	50 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE	570 mg
ARGININE	316 mg
GLYCINE	283 mg
PROLINE	187 mg
SERINE	138 mg
TYROSINE	11 mg

**mEq/L**

ACETATE	24
CHLORIDE	11

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

pH 8.0 (4.5 TO 7.0) HYPERTONIC  
OSMOLARITY 525 mOsmol/L (CALC)

STERILE NONPYROGENIC  
SINGLE DOSE CONTAINER

**Baxter**

BAXTER HEALTHCARE CORPORATION  
CLINTEC NUTRITION DIVISION  
DEERFIELD, IL 60015 USA  
MADE IN USA  
PATENT PENDING  
BAXTER, CLINIMIX AND CLARITY ARE TRADEMARKS  
OF BAXTER INTERNATIONAL INC

**Container Label**

LOT EXP

2B7701 NDC 0338-1083-04

**CLINIMIX 2.75/5**  
sulfite-free  
**(2.75% Amino Acid  
in 5% Dextrose)**  
**Injection**

**1000 mL INJECTION PORT CHAMBER**  
**10% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**5.5% Amino Acid Injection**

**Rx Only**

CHECK FOR MINUTE LEAKS BY SQUEEZING  
EACH CHAMBER OF THE BAG  
BEFORE USE GRASP EACH SIDE OF

THE TOP OF THE BAG AND ROLL BAG

TO OPEN SEAL BETWEEN CHAMBERS

MIX THOROUGHLY

AFTER MIXING THE PRODUCT

REPRESENTS 2000 mL

ADDITIVES MAY BE INCOMPATIBLE CONSULT WITH

PHARMACIST IF AVAILABLE WHEN INTRODUCING

ADDITIVES USE ASEPTIC TECHNIQUE

MIX THOROUGHLY DO NOT STORE

DOSAGE INTRAVENOUSLY AS DIRECTED BY A

PHYSICIAN DO NOT USE UNLESS SOLUTION IS

CLEAR REFRIGERATED STORAGE AFTER MIXING

LIMITED TO 24 HOURS DISCARD UNUSED PORTION

SEE ACCOMPANYING DIRECTIONS FOR USE

CAUTIONS MUST NOT BE USED IN SERIES

CONNECTIONS DISCONTINUE INFUSION

IF ADVERSE REACTION OCCURS

CLARITY DUAL CHAMBER CONTAINER PL 2401 PLASTIC

CONTENTS OF EACH 100 mL OF THE

ADMIXED INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 201 mg

ISOLEUCINE 165 mg

VALINE 160 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 159 mg

PHENYLALANINE 154 mg

HISTIDINE 132 mg

THREONINE 116 mg

METHIONINE 110 mg

TRYPTOPHAN 50 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 570 mg

ARGININE 316 mg

GLYCINE 283 mg

PROLINE 187 mg

SERINE 138 mg

TYROSINE 11 mg

mEq/L

ACETATE 24

CHLORIDE 11

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

pH 6.0 (4.5 TO 7.0) HYPERTONIC  
OSMOLARITY 525 mOsmol/L (CALC)  
STERILE NONPYROGENIC  
SINGLE DOSE CONTAINER

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**CLINTEC NUTRITION DIVISION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

**PATENT PENDING**

**BAXTER CLINIMIX AND CLARITY ARE TRADEMARKS  
OF BAXTER INTERNATIONAL INC**

2B7726 NDC 0338-1133-03

**CLINIMIX**

**4.25/5**

**SULFITE-FREE**

**(4.25% Amino Acids  
in 5% Dextrose)  
Injection**

**500 mL INJECTION PORT CHAMBER  
10% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

**LOT**

**EXP**

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 980 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

**mEq/L**

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7726 NDC 0338-1133-03

CLINIMIX

4.25/5



SULFITE-FREE

(4.25% Amino Acids  
in 5% Dextrose)  
Injection

500 mL INJECTION PORT CHAMBER  
10% Dextrose Injection USP

500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection

ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37  
 CHLORIDE 17  
 BALANCED BY IONS FROM AMINO ACIDS  
 pH ADJUSTED WITH GLACIAL ACETIC ACID  
 STERILE  
 SINGLE DOSE CONTAINER  
 ROOM TEMPERATURE (25°C/77°F)  
 AVOID EXCESSIVE HEAT  
 PROTECT FROM FREEZING  
 SEE PRESCRIBING INFORMATION

**Baxter**

BAXTER HEALTHCARE CORPORATION  
 DEERFIELD IL 60015 USA  
 MADE IN USA

2B7704 NDC 0338-1089-04

**CLINIMIX**  
**4.25/5**

**SULFITE-FREE**

**(4.25% Amino Acids  
 in 5% Dextrose)  
 Injection**

**1000 mL INJECTION PORT CHAMBER**  
**10% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
 MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
 ACTIVATION  
 AFTER MIXING THE PRODUCT REPRESENTS 2000 mL  
 REFRIGERATED STORAGE IS LIMITED TO 9 DAYS  
 A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
 AND EFFICACY OF THIS PRODUCT  
 ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

**LOT EXP**

CONTENTS OF EACH 100 mL OF THE  
 ADMIXED INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE	311 mg
ISOLEUCINE	255 mg
VALINE	247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	247 mg
PHENYLALANINE	238 mg
HISTIDINE	204 mg
THREONINE	179 mg
METHIONINE	170 mg
TRYPTOPHAN	77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE	880 mg
ARGININE	489 mg
GLYCINE	438 mg
PROLINE	289 mg
SERINE	213 mg
TYROSINE	17 mg

mEq/L

ACETATE	37
CHLORIDE	17

BALANCED BY IONS FROM AMINO ACIDS  
 pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
 SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)  
 AVOID EXCESSIVE HEAT  
 PROTECT FROM FREEZING  
 SEE PRESCRIBING INFORMATION

**Baxter**  
 BAXTER HEALTHCARE CORPORATION  
 DEERFIELD IL 60015 USA  
 MADE IN USA

Container Label

LOT EXP

**2B7704 NDC 0338-1089-04**

**CLINIMIX 4.25/5  
SULFITE-FREE  
(4.25% Amino Acid  
in 5% Dextrose)  
Injection**

1000 mL INJECTION PORT CHAMBER  
10% Dextrose Injection USP

1000 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 5 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER  
ROOM TEMPERATURE (25°C/77°F)  
AVOID EXCESSIVE HEAT  
PROTECT FROM FREEZING  
SEE PRESCRIBING INFORMATION

**Baxter**  
**BAXTER HEALTHCARE CORPORATION**  
**DEERFIELD IL 60015 USA**  
**MADE IN USA**

2B7705 NDC 0338-1091-04

**CLINIMIX**

**4.25/10**

**SULFITE-FREE**

**(4.25% Amino Acids  
in 10% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**1000 mL INJECTION PORT CHAMBER**  
**20% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

**LOT**                      **EXP**

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP                      10 g

**ESSENTIAL AMINO ACIDS**

LEUCINE    311 mg

ISOLEUCINE                                        255 mg

VALINE    247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT)                      247 mg

PHENYLALANINE                                 238 mg

HISTIDINE                                         204 mg

THREONINE                                        179 mg

METHIONINE                                      170 mg

TRYPTOPHAN                                     77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE    880 mg

ARGININE                                         489 mg

GLYCINE    438 mg

PROLINE     289 mg

SERINE    213 mg

TYROSINE                                         17 mg

**mEq/L**

ACETATE    37

CHLORIDE                                         17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

**LOT EXP**

**2B7705 NDC 0338-1091-04**

**CLINIMIX 4.25/10**

**SULFITE-FREE**

**(4.25% Amino Acid**

**in 10% Dextrose)**

**Injection**

1000 mL INJECTION PORT CHAMBER

20% Dextrose Injection USP

1000 mL OUTLET PORT CHAMBER

8.5% Amino Acid Injection

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 10 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING  
SEE PRESCRIBING INFORMATION

**Baxter**

**BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA**

2B7728 NDC 0338-1135-03

**CLINIMIX**

**4.25/20**

**SULFITE-FREE**

**(4.25% Amino Acids  
in 20% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**500 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

**LOT**

**EXP**

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE

HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

**mEq/L**

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**

**BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA**

Container Label

**LOT EXP**

**2B7728 NDC 0338-1035-03**

**CLINIMIX**

**4.25/20**

**SULFITE-FREE**

**(4.25% Amino Acid  
in 20% Dextrose)**

**Injection**

**500 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

*Baxter*

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7706 NDC 0338-1093-04

# CLINIMIX 4.25/20

**SULFITE-FREE**  
**(4.25% Amino Acids  
in 20% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**1000 mL INJECTION PORT CHAMBER**  
**40% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g  
**ESSENTIAL AMINO ACIDS**  
LEUCINE 311 mg  
ISOLEUCINE 255 mg  
VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg  
PHENYLALANINE 238 mg  
HISTIDINE 204 mg  
THREONINE 179 mg  
METHIONINE 170 mg  
TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**  
ALANINE 890 mg  
ARGININE 489 mg  
GLYCINE 438 mg  
PROLINE 289 mg  
SERINE 213 mg  
TYROSINE 17 mg

**mEq/L**  
ACETATE 37  
CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID  
STERILE  
SINGLE DOSE CONTAINER

**Baxter**

BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7706 NDC 0338-1093-04

**CLINIMIX 4.25/20**  
**SULFITE-FREE**  
**(4.25% Amino Acid  
in 20% Dextrose)  
Injection**

**1000 mL INJECTION PORT CHAMBER**  
**40% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER**  
**8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL



REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY

AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

***Baxter***

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7729 NDC 0338-1136-03

# CLINIMIX

## 4.25/25

SULFITE-FREE

**(4.25% Amino Acids  
in 25% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**500 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

**mEq/L**

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

**LOT EXP**

**2B7729 NDC 0338-1136-03**

**CLINIMIX 4.25/25**

**SULFITE-FREE**

**(4.25% Amino Acid  
in 25% Dextrose)**

**Injection**

**500 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

***Baxter***

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7707 NDC 0338-1095-04

**CLINIMIX**  
**4.25/25**

**SULFITE-FREE**

**(4.25% Amino Acids  
in 25% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**1000 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT EXP

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7707 NDC 0338-1095-04

**CLINIMIX**

**4.25/25**

**SULFITE-FREE**

**(4.25% Amino Acid  
in 25% Dextrose)**

**Injection**

**1000 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
8.5% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 311 mg

ISOLEUCINE 255 mg

VALINE 247 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 247 mg

PHENYLALANINE 238 mg

HISTIDINE 204 mg

THREONINE 179 mg

METHIONINE 170 mg

TRYPTOPHAN 77 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 880 mg

ARGININE 489 mg

GLYCINE 438 mg

PROLINE 289 mg

SERINE 213 mg

TYROSINE 17 mg

mEq/L

ACETATE 37

CHLORIDE 17

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7730 NDC 0338-1137-03

# CLINIMIX 5/15

**SULFITE-FREE**

**(5% Amino Acids  
in 15% Dextrose)**

**Injection**

**CENTRAL LINE INFUSION ONLY**

**500 mL INJECTION PORT CHAMBER  
30% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 15 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 290 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

mEq/L

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

BAXTER HEALTHCARE CORPORATION

DEERFIELD IL 60015 USA

MADE IN USA

Container Label

LOT EXP

2B7730 NDC 0338-1137-03

**CLINIMIX**

**5/15**

**SULFITE-FREE**

**(5% Amino Acid  
in 15% Dextrose)**

**Injection**

**500 mL INJECTION PORT CHAMBER  
30% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 15 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

mEq/L

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7709 NDC 0338-1099-04

# CLINIMIX 5/15

**SULFITE-FREE**

**(5% Amino Acids  
in 15% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**1000 mL INJECTION PORT CHAMBER  
30% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION  
AFTER MIXING THE PRODUCT REPRESENTS 2000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS  
A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT  
ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

**LOT**

**EXP**

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP	15 g
<b>ESSENTIAL AMINO ACIDS</b>	
LEUCINE	365 mg
ISOLEUCINE	300 mg
VALINE	290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg
PHENYLALANINE	280 mg
HISTIDINE	240 mg
THREONINE	210 mg
METHIONINE	200 mg
TRYPTOPHAN	90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE	1035 mg
ARGININE	575 mg
GLYCINE	515 mg
PROLINE	340 mg
SERINE	250 mg
TYROSINE	20 mg

**mEq/L**

ACETATE	42
CHLORIDE	20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT  
PROTECT FROM FREEZING  
SEE PRESCRIBING INFORMATION

**Baxter**

BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

**LOT EXP**

**2B7709 NDC 0338-1099-04**

**CLINIMIX**

**5/15**

**SULFITE-FREE**

**(5% Amino Acid  
in 15% Dextrose)**

**Injection**

**1000 mL INJECTION PORT CHAMBER  
30% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON



ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY

AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 15 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

mEq/L

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7731 NDC 0338-1138-03

# CLINIMIX 5/20

**SULFITE-FREE**

**(5% Amino Acids  
in 20% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**500 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

**mEq/L**

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7731 NDC 0338-1138-03

**CLINIMIX**

**5/**

**SULFITE-FREE**

**(5% Amino Acid  
in 20% Dextrose)**

**Injection**

**500 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS  
A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT  
ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY  
CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg  
ISOLEUCINE 300 mg  
VALINE 290 mg  
LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg  
PHENYLALANINE 280 mg  
HISTIDINE 240 mg  
THREONINE 210 mg  
METHIONINE 200 mg  
TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg  
ARGININE 575 mg  
GLYCINE 515 mg  
PROLINE 340 mg  
SERINE 250 mg  
TYROSINE 20 mg

mEq/L

ACETATE 42  
CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)  
AVOID EXCESSIVE HEAT  
PROTECT FROM FREEZING  
SEE PRESCRIBING INFORMATION

**Baxter**  
**BAXTER HEALTHCARE CORPORATION**  
**DEERFIELD IL 60015 USA**  
**MADE IN USA**

2B7710 NDC 0338-1101-04

# CLINIMIX 5/20

**SULFITE-FREE**

**(5% Amino Acids  
in 20% Dextrose)  
Injection**

**CENTRAL LINE INFUSION ONLY**

**1000 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP	20 g
<b>ESSENTIAL AMINO ACIDS</b>	
LEUCINE	365 mg
ISOLEUCINE	300 mg
VALINE	290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg
PHENYLALANINE	280 mg
HISTIDINE	240 mg
THREONINE	210 mg
METHIONINE	200 mg
TRYPTOPHAN	90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE	1035 mg
ARGININE	575 mg
GLYCINE	515 mg
PROLINE	340 mg
SERINE	250 mg
TYROSINE	20 mg

mEq/L

ACETATE	42
CHLORIDE	20

BALANCED BY IONS FROM AMINO ACIDS  
PH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7710 NDC 0338-1101-04

CLINIMIX

5/20

**SULFITE-FREE**

**(5% Amino Acid  
in 20% Dextrose)**

**Injection**

**1000 mL INJECTION PORT CHAMBER  
40% Dextrose Injection USP**

**1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection**

**Rx Only**

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON

ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

mEq/L

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7732 NDC 0338-1139-03

# CLINIMIX 5/25

SULFITE-FREE

(5% Amino Acids  
in 25% Dextrose)  
Injection

**CENTRAL LINE INFUSION ONLY**

500 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP

500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection

Rx Only

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED  
INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 290 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

**mEq/L**

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7732 NDC 0338-1139-03

CLINIMIX

5/25

SULFITE-FREE

(5% Amino Acid  
in 25% Dextrose)

Injection

500 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP

500 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection

Rx Only

**ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE**

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

mEq/L

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE

SINGLE DOSE CONTAINER

**Baxter**

**BAXTER HEALTHCARE CORPORATION**

**DEERFIELD IL 60015 USA**

**MADE IN USA**

2B7711 NDC 0338-1103-04

# CLINIMIX 5/25

SULFITE-FREE

(5% Amino Acids  
in 25% Dextrose)  
Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP

1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

LOT

EXP

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg

ISOLEUCINE 300 mg

VALINE 290 mg

LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg

PHENYLALANINE 280 mg

HISTIDINE 240 mg

THREONINE 210 mg

METHIONINE 200 mg

TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg

ARGININE 575 mg

GLYCINE 515 mg

PROLINE 340 mg

SERINE 250 mg

TYROSINE 20 mg

**mEq/L**

ACETATE 42

CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

**Baxter**  
BAXTER HEALTHCARE CORPORATION  
DEERFIELD IL 60015 USA  
MADE IN USA

Container Label

LOT EXP

2B7711 NDC 0338-1103-04

CLINIMIX

5/25

SULFITE-FREE

(5% Amino Acid  
in 25% Dextrose)

Injection

1000 mL INJECTION PORT CHAMBER  
50% Dextrose Injection USP

1000 mL OUTLET PORT CHAMBER  
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND  
MIX THOROUGHLY BEFORE USE



SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON  
ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL  
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS  
A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY  
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE  
ADMIXED INJECTION

DEXTROSE HYDROUS USP 25 g

**ESSENTIAL AMINO ACIDS**

LEUCINE 365 mg  
ISOLEUCINE 300 mg  
VALINE 290 mg  
LYSINE (ADDED AS THE  
HYDROCHLORIDE SALT) 290 mg  
PHENYLALANINE 280 mg  
HISTIDINE 240 mg  
THREONINE 210 mg  
METHIONINE 200 mg  
TRYPTOPHAN 90 mg

**NONESSENTIAL AMINO ACIDS**

ALANINE 1035 mg  
ARGININE 575 mg  
GLYCINE 515 mg  
PROLINE 340 mg  
SERINE 250 mg  
TYROSINE 20 mg

mEq/L  
ACETATE 42  
CHLORIDE 20

BALANCED BY IONS FROM AMINO ACIDS  
pH ADJUSTED WITH GLACIAL ACETIC ACID

STERILE  
SINGLE DOSE CONTAINER

**Baxter**  
**BAXTER HEALTHCARE CORPORATION**  
**DEERFIELD IL 60015 USA**  
**MADE IN USA**

**CLINIMIX**

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1132
---------------------	-------------------------	---------------------------	---------------

Route of Administration INTRAVENOUS

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	201 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	154 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	159 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	110 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	165 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	160 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	132 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	116 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	50 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	570 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	283 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	316 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	187 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	138 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	11 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	5 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	
NITROGEN (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1132-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	05/14/2018

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1083
Route of Administration	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	201 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	154 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	159 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	110 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	165 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	160 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	132 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	116 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	50 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	570 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	283 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	316 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	187 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	138 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	11 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	5 g in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	
NITROGEN (UNII: N762921K75)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1083-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	05/14/2018

**CLINIMIX**

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

**Product Information**

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1133
Route of Administration	INTRAVENOUS		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	5 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	
NITROGEN (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1133-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1089
Route of Administration	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL

<b>PHENYLALANINE</b> (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	5 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1089-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1134
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL

<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	10 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1134-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/20/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	03/20/2012	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1091
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5017Y3R) (PHENYLALANINE - UNII:47E5017Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL

<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	10 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1091-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1135
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5017Y3R) (PHENYLALANINE - UNII:47E5017Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL

<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	20 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1135-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	05/14/2018

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1093
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5017Y3R) (PHENYLALANINE - UNII:47E5017Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL



<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	20 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1093-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	05/14/2018

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1136
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL

<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	25 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>NITROGEN</b> (UNII: N762921K75)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1136-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	02/29/2020

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	02/29/2020

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1095
<b>Route of Administration</b>	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LEUCINE</b> (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
<b>PHENYLALANINE</b> (UNII: 47E5017Y3R) (PHENYLALANINE - UNII:47E5017Y3R)	PHENYLALANINE	238 mg in 100 mL
<b>LYSINE</b> (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
<b>METHIONINE</b> (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
<b>ISOLEUCINE</b> (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
<b>VALINE</b> (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
<b>HISTIDINE</b> (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
<b>THREONINE</b> (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
<b>TRYPTOPHAN</b> (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
<b>ALANINE</b> (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
<b>GLYCINE</b> (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
<b>ARGININE</b> (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
<b>PROLINE</b> (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
<b>SERINE</b> (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL

<b>TYROSINE</b> (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL		
<b>DEXTROSE</b> (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	25 g in 100 mL		
<b>Inactive Ingredients</b>				
	<b>Ingredient Name</b>	<b>Strength</b>		
	ACETIC ACID (UNII: Q40Q9N063P)			
	WATER (UNII: 059QF0K00R)			
	NITROGEN (UNII: N762921K75)			
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0338-1095-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
NDA	NDA020734	09/29/1997	05/14/2018	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

<b>Product Information</b>			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0338-1137
<b>Route of Administration</b>	INTRAVENOUS		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
	PHENYLALANINE (UNII: 47E5017Y3R) (PHENYLALANINE - UNII:47E5017Y3R)	PHENYLALANINE	280 mg in 100 mL
	LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
	METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
	ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
	VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
	HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
	THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
	TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
	ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
	GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
	ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
	PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
	SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
	TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
	DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	15 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1137-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1099
Route of Administration	INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	15 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1099-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1138
Route of Administration	INTRAVENOUS		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	20 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

ACETIC ACID (UNII: Q40Q9N063P)				
WATER (UNII: 059QF0KO0R)				
NITROGEN (UNII: N762921K75)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1138-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA020734	09/29/1997		

<b>CLINIMIX</b>			
leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection			
<b>Product Information</b>			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1101
Route of Administration	INTRAVENOUS		
<b>Active Ingredient/Active Moiety</b>			
Ingredient Name	Basis of Strength	Strength	
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL	
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL	
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL	
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL	
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL	
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL	
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL	
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL	
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL	
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL	
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL	
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL	
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL	
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL	
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL	
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	20 g in 100 mL	
<b>Inactive Ingredients</b>			
Ingredient Name	Strength		
ACETIC ACID (UNII: Q40Q9N063P)			
WATER (UNII: 059QF0KO0R)			

NITROGEN (UNII: N762921K75)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1101-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1139
Route of Administration	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	25 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	
NITROGEN (UNII: N762921K75)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1139-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	02/29/2020

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020734	09/29/1997	02/29/2020

## CLINIMIX

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dextrose injection

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0338-1103
Route of Administration	INTRAVENOUS		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (DEXTROSE - UNII:IY9XDZ35W2)	DEXTROSE	25 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0K00R)	
NITROGEN (UNII: N762921K75)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------



1	NDC:0338-1103-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	09/29/1997	05/14/2018
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA020734		09/29/1997	05/14/2018

**Labeler** - Baxter Healthcare Corporation (005083209)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Baxter Healthcare Corporation		189326168	ANALYSIS(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , MANUFACTURE(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , LABEL(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , PACK(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , STERILIZE(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Baxter Healthcare Corporation		059140764	ANALYSIS(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , MANUFACTURE(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , LABEL(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , PACK(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103) , STERILIZE(0338-1132, 0338-1083, 0338-1133, 0338-1089, 0338-1134, 0338-1091, 0338-1135, 0338-1093, 0338-1136, 0338-1095, 0338-1137, 0338-1099, 0338-1138, 0338-1101, 0338-1139, 0338-1103)