

**BERKLEY AND JENSEN ASPIRIN- aspirin tablet, film coated  
BJWC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**BJWC Aspirin Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily relieves

- headache
- muscle pain
- toothache
- menstrual pain
- pain and fever of colds
- minor pain of arthritis

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- diabetes
- gout
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts more than 3 days
  - ringing in the ears or loss of hearing occurs
  - redness or swelling is present
  - new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose

- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)

### **Inactive ingredients**

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

### **Questions or comments?**

1-800-934-1204

### **Principal Display Panel**

Compare to the active ingredient in Bayer® Aspirin

ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

FOR ADULTS

SAFE PAIN RELIEF PLUS LIFESAVING BENEFITS

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART.

GLUTEN FREE

325 mg EACH

500 COATED TABLETS

325 mg EACH

ACTUAL SIZE

**Drug Facts** (continued)

**Directions**  
 Adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.  
 Children under 12 years: consult a doctor.

**Other Information**  
 ■ Store at 20°C-25°C (68°-77°F).  
 ■ Inactive ingredients contain calcium carbonate, calcium phosphate dibasic, hypromellose, talc, triacetin.

**Questions or comments?** 1-800-438-1204

With respect to the use of this product, please refer to the package insert of Bayer Aspirin.  
 Distributor of Bayer Aspirin:  
 100% Bayer Aspirin  
 100% Melted Bayer Aspirin  
 Distributed by:  
 BJS Wholesale Club  
 25 Pleasant Drive  
 Westborough, MA 01581

**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

Compare to the active ingredient in Bayer® Aspirin®



# ASPIRIN

PAIN RELIEVER/FEVER REDUCER (NSAID)

FOR ADULTS  
SAFE PAIN RELIEF PLUS LIFESAVING BENEFITS

TALK TO YOUR DOCTOR OR OTHER HEALTHCARE PROVIDER BEFORE USING THIS PRODUCT FOR YOUR HEART.

GLUTEN FREE

500

COATED TABLETS

325

mg EACH



ACTUAL SIZE

**Drug Facts** (in each tablet)

**Active ingredient:** Aspirin (25 mg Aspirin)  
**Purpose:** Pain reliever/fever reducer

**Uses:** temporarily relieve:  
 ■ headache  
 ■ muscle pain  
 ■ pain from fever of child  
 ■ minor pain of arthritis

**Warnings:** Children and teenagers who have an sore throat from chicken pox or the grippe/flu should not use this product. When using this product, if changes in behavior with continued vomiting occur, contact a doctor because these changes may be a sign of a serious condition.  
 ■ Allergic reactions may cause a severe allergic reaction with my blood.  
 ■ This is a hard swelling (edema) (swelling) which may cause stomach bleeding.  
 ■ This product contains an NSAID, which may cause:  
 ■ stroke  
 ■ heart attack  
 ■ kidney failure  
 ■ stomach bleeding  
 ■ take a blood thinner (anticoagulant) or steroid drug  
 ■ take aspirin with other pain relievers/fever reducers  
 ■ have 3 or more alcoholic drinks every day while using this product  
 ■ have a history of stomach ulcers or bleeding  
 ■ have a history of kidney disease  
 ■ have a history of liver disease  
 ■ have a history of heart disease  
 ■ have a history of high blood pressure  
 ■ have a history of asthma  
 ■ have a history of diabetes  
 ■ have a history of bleeding disorders  
 ■ have a history of low platelet counts  
 ■ have a history of low sodium levels  
 ■ have a history of low potassium levels  
 ■ have a history of low magnesium levels  
 ■ have a history of low calcium levels  
 ■ have a history of low iron levels  
 ■ have a history of low zinc levels  
 ■ have a history of low vitamin D levels  
 ■ have a history of low vitamin K levels  
 ■ have a history of low vitamin B12 levels  
 ■ have a history of low vitamin C levels  
 ■ have a history of low vitamin E levels  
 ■ have a history of low vitamin A levels  
 ■ have a history of low vitamin B6 levels  
 ■ have a history of low vitamin B9 levels  
 ■ have a history of low vitamin B1 levels  
 ■ have a history of low vitamin B2 levels  
 ■ have a history of low vitamin B3 levels  
 ■ have a history of low vitamin B5 levels  
 ■ have a history of low vitamin B7 levels  
 ■ have a history of low vitamin B8 levels  
 ■ have a history of low vitamin B10 levels  
 ■ have a history of low vitamin B11 levels  
 ■ have a history of low vitamin B12 levels  
 ■ have a history of low vitamin B13 levels  
 ■ have a history of low vitamin B14 levels  
 ■ have a history of low vitamin B15 levels  
 ■ have a history of low vitamin B16 levels  
 ■ have a history of low vitamin B17 levels  
 ■ have a history of low vitamin B18 levels  
 ■ have a history of low vitamin B19 levels  
 ■ have a history of low vitamin B20 levels  
 ■ have a history of low vitamin B21 levels  
 ■ have a history of low vitamin B22 levels  
 ■ have a history of low vitamin B23 levels  
 ■ have a history of low vitamin B24 levels  
 ■ have a history of low vitamin B25 levels  
 ■ have a history of low vitamin B26 levels  
 ■ have a history of low vitamin B27 levels  
 ■ have a history of low vitamin B28 levels  
 ■ have a history of low vitamin B29 levels  
 ■ have a history of low vitamin B30 levels  
 ■ have a history of low vitamin B31 levels  
 ■ have a history of low vitamin B32 levels  
 ■ have a history of low vitamin B33 levels  
 ■ have a history of low vitamin B34 levels  
 ■ have a history of low vitamin B35 levels  
 ■ have a history of low vitamin B36 levels  
 ■ have a history of low vitamin B37 levels  
 ■ have a history of low vitamin B38 levels  
 ■ have a history of low vitamin B39 levels  
 ■ have a history of low vitamin B40 levels  
 ■ have a history of low vitamin B41 levels  
 ■ have a history of low vitamin B42 levels  
 ■ have a history of low vitamin B43 levels  
 ■ have a history of low vitamin B44 levels  
 ■ have a history of low vitamin B45 levels  
 ■ have a history of low vitamin B46 levels  
 ■ have a history of low vitamin B47 levels  
 ■ have a history of low vitamin B48 levels  
 ■ have a history of low vitamin B49 levels  
 ■ have a history of low vitamin B50 levels  
 ■ have a history of low vitamin B51 levels  
 ■ have a history of low vitamin B52 levels  
 ■ have a history of low vitamin B53 levels  
 ■ have a history of low vitamin B54 levels  
 ■ have a history of low vitamin B55 levels  
 ■ have a history of low vitamin B56 levels  
 ■ have a history of low vitamin B57 levels  
 ■ have a history of low vitamin B58 levels  
 ■ have a history of low vitamin B59 levels  
 ■ have a history of low vitamin B60 levels  
 ■ have a history of low vitamin B61 levels  
 ■ have a history of low vitamin B62 levels  
 ■ have a history of low vitamin B63 levels  
 ■ have a history of low vitamin B64 levels  
 ■ have a history of low vitamin B65 levels  
 ■ have a history of low vitamin B66 levels  
 ■ have a history of low vitamin B67 levels  
 ■ have a history of low vitamin B68 levels  
 ■ have a history of low vitamin B69 levels  
 ■ have a history of low vitamin B70 levels  
 ■ have a history of low vitamin B71 levels  
 ■ have a history of low vitamin B72 levels  
 ■ have a history of low vitamin B73 levels  
 ■ have a history of low vitamin B74 levels  
 ■ have a history of low vitamin B75 levels  
 ■ have a history of low vitamin B76 levels  
 ■ have a history of low vitamin B77 levels  
 ■ have a history of low vitamin B78 levels  
 ■ have a history of low vitamin B79 levels  
 ■ have a history of low vitamin B80 levels  
 ■ have a history of low vitamin B81 levels  
 ■ have a history of low vitamin B82 levels  
 ■ have a history of low vitamin B83 levels  
 ■ have a history of low vitamin B84 levels  
 ■ have a history of low vitamin B85 levels  
 ■ have a history of low vitamin B86 levels  
 ■ have a history of low vitamin B87 levels  
 ■ have a history of low vitamin B88 levels  
 ■ have a history of low vitamin B89 levels  
 ■ have a history of low vitamin B90 levels  
 ■ have a history of low vitamin B91 levels  
 ■ have a history of low vitamin B92 levels  
 ■ have a history of low vitamin B93 levels  
 ■ have a history of low vitamin B94 levels  
 ■ have a history of low vitamin B95 levels  
 ■ have a history of low vitamin B96 levels  
 ■ have a history of low vitamin B97 levels  
 ■ have a history of low vitamin B98 levels  
 ■ have a history of low vitamin B99 levels  
 ■ have a history of low vitamin B100 levels

**Drug Facts** (continued)

**Directions**  
 Adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.  
 Children under 12 years: consult a doctor.

**Other Information**  
 ■ Store at 20°C-25°C (68°-77°F).  
 ■ Inactive ingredients contain calcium carbonate, calcium phosphate dibasic, hypromellose, talc, triacetin.

**Questions or comments?** 1-800-438-1204

With respect to the use of this product, please refer to the package insert of Bayer Aspirin.  
 Distributor of Bayer Aspirin:  
 100% Bayer Aspirin  
 100% Melted Bayer Aspirin  
 Distributed by:  
 BJS Wholesale Club  
 25 Pleasant Drive  
 Westborough, MA 01581

**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

<b>BERKLEY AND JENSEN ASPIRIN</b>			
aspirin tablet, film coated			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68391-416
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
Inactive Ingredients			
<b>Ingredient Name</b>	<b>Strength</b>		
STARCH, CORN (UNII: O8232NY3SJ)			
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			
TALC (UNII: 7SEV7J4R1U)			
TRIACETIN (UNII: XHX3C3X673)			
Product Characteristics			
<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	14mm

<b>Flavor</b>		<b>Imprint Code</b>	Aspirin;L	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68391-416-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/23/2008	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
OTC monograph not final	part343	07/23/2008		

**Labeler - BJWC (159082692)**

Revised: 8/2018

BJWC