

ALLERGY RELIEF D- loratadine, pseudoephedrine sulfate tablet, film coated, extended release
Meijer Distribution Inc

Meijer Distribution, Inc. Allergy Relief-D Drug Facts

Active ingredients (in each tablet)

Loratadine 5 mg

Pseudoephedrine sulfate 120 mg

Purpose

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- sneezing
- itchy, watery eyes
- runny nose
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed.

Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- **each tablet contains:** calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Claritin-D® 12 hour active ingredients

NON-DROWSY*

allergy relief-D

Pseudoephedrine Sulfate 120 mg | Nasal Decongestant

Loratadine 5 mg | Antihistamine

Extended-Release Tablets

INDOOR & OUTDOOR ALLERGIES

12 HOUR RELIEF OF:

Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing; Runny Nose; Itchy, Watery Eyes; Itchy Throat or Nose Due to Allergies

10 Tablets

ALLERGY & CONGESTION

*When taken as directed. See Drug Facts Panel.

actual size

12 HOUR allergy relief-D

ALLERGY & CONGESTION

meijer

NDC 41250-954-46

Compare to
Claritin-D® 12 hour
active ingredients**

NON-DROWSY*

allergy relief-D

Pseudoephedrine Sulfate 120 mg | Nasal Decongestant
Loratadine 5 mg | Antihistamine
Extended-Release Tablets

INDOOR & OUTDOOR ALLERGIES

12 HOUR RELIEF OF:

Nasal and Sinus Congestion Due to Colds or Allergies
Sneezing; Runny Nose; Itchy, Watery Eyes; Itchy Throat or Nose Due to Allergies

ALLERGY & CONGESTION

*When taken as directed. See Drug Facts Panel.

**10
Tablets**

actual
size



Drug Facts

Active ingredients (in each tablet)

Loratadine 5 mg.....Antihistamine
Pseudoephedrine sulfate 120 mg.....Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
 - sneezing
 - itchy, watery eyes
 - runny nose
 - itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores free breathing through the nose

Warnings

- Do not use**
 - if you have ever had an allergic reaction to this product or any of its ingredients
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Drug Facts (continued)

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not divide, crush, chew or dissolve the tablet
- adults and children 12 years and over: 1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

Other information

- each tablet contains calcium 25 mg
- do not use if blister unit is broken or torn
- store between 20° to 25°C (68° to 77°F)
- keep in a dry place

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate, hypromellose, lactose monohydrate, magnesium stearate, pharmaceutical ink, povidone, titanium dioxide

Drug Facts (continued)

Questions or comments? 1-800-719-9260

PID 528 577



7 13733 87640 7

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Claritin-D®.

DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



WWW.MEIJER.COM/
SATISFACTION

7U046 6E C3

OPEN OTHER END

CONVENIENT RECLOSING TAB

ALLERGY RELIEF D

loratadine, pseudoephedrine sulfate tablet, film coated, extended release

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-954	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)		LORATADINE	5 mg	
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)		PSEUDOEPHEDRINE SULFATE	120 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)				
HYPROMELLOSES (UNII: 3NXW29V3WO)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
POVIDONE (UNII: FZ989GH94E)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	WHITE (to off-white)	Score	no score	
Shape	ROUND	Size	12mm	
Flavor		Imprint Code	7U0	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-954-46	10 in 1 CARTON	06/15/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-954-60	20 in 1 CARTON	06/12/2018	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA076050	06/12/2018		

