

**VICKS DAYQUIL COLD AND FLU MULTI-SYMPTOM RELIEF- acetaminophen,
dextromethorphan hydrobromide, and phenylephrine hydrochloride liquid**
The Procter & Gamble Manufacturing Company

VICKS®
DayQuil™
COLD & FLU

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:

If sore throat is severe persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms get worse or last more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:**sodium 46 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: Do not use if printed shrinkband is broken or missing.

MADE IN CANADA

DIST. BY

PROCTER & GAMBLE, CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL

VICKS ®

DayQuil ®

COLD & FLU

Acetaminophen - Pain Reliever/Fever Reducer

Phenylephrine HCl - Nasal Decongestant
 Dextromethorphan HBr - Cough Suppressant

Non-Drowsy

Alcohol Free

4 FL OZ (118 mL)



Drug Facts (continued)
Uses temporarily relieves common cold/flu symptoms:
 • nasal congestion
 • cough due to minor throat & bronchial irritation
 • sore throat • headache
 • minor aches & pains • fever
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if
 • adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
 • child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
 • taken with other drugs containing acetaminophen
 • adult has 3 or more alcoholic drinks every day

Drug Facts (continued)
 while using this product
Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin redness • hives • rash
 If a skin reaction occurs, stop use and seek medical help right away
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)
Ask a doctor before use if you have
 • liver disease • heart disease
 • high blood pressure • thyroid disease
 • diabetes
 • trouble urinating due to enlarged prostate gland
 • cough that occurs with too much phlegm (mucus)
 • persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 • a sodium-restricted diet
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product, do not use more than directed.
Stop use and ask a doctor if
 • you get nervous, dizzy or sleepless
 • pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 • fever gets worse or lasts more than 3 days
 • redness or swelling is present

Drug Facts (continued)
 • new symptoms occur
 • cough comes back or occurs with rash or headache that lasts
 These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 • take only as directed
 • only use the dose cup provided
 • do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 4 hrs
children 6 to under 12 yrs 15 mL every 4 hrs
children 4 to under 6 yrs ask a doctor
children under 4 yrs do not use
Other information
 • each 15 mL contains: sodium 46 mg

Drug Facts (continued)
 • store at no greater than 25°C and do not refrigerate
Inactive ingredients citric acid, FD&C yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum
Questions? 1-800-362-1883



VICKS DAYQUIL COLD AND FLU MULTI-SYMPTOM RELIEF
 acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-980
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	APRICOT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-980-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/12/2022	06/15/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/12/2022	06/15/2026

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 8/2024

The Procter & Gamble Manufacturing Company