

**TUSSIN DM- dextromethorphan hydrobromide, guaifenesin liquid**  
**Discount Drug Mart**

-----

**Drug Facts**

**Active ingredients (in each 5 mL tsp)**

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

**Purpose**

Cough Suppressant

Expectorant

**Keep out of reach of children.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING.**

Do not use

- in a child under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before taking if you have**

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, bronchitis or emphysema

**Stop use and ask a doctor if**

cough lasts more than 7 days, comes back or is accompanied by fever, rash or

persistent headache.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Directions**

- do not take more than 6 doses in any 24-hour period

---

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

---

**Other information**

- **Keep carton for full Direction for use.**
- store at 20°-25°C (68°-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

**Inactive ingredients**

Anhydrous citric acid, dextrose, FD and red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?**

Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

**DISCOUNT drug mart FOOD FAIR Tussin DM product label**

**\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH AND CHEST CONGESTION DM**

**DISCOUNT**

**drug mart**

**FOOD FAIR**

**Tussin DM**

**Dextromethorphan HBr/ Guaifenesin**

**NON-DROWSY**

**COUGH SUPPRESSANT**

## EXPECTORANT

Helps Loosen

Chest Congestion

Cough Formula

for ages 12 & over

8 FL OZ (237 mL)

\* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

### **BX-039**

#### **SATISFACTION GUARANTEED**

IF DISSASIFIED , RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

DO NOT USE IF PRINTED SEAL UNDER CAP IS  
TORN OR MISSING

## Drug Facts

### Active ingredients Purpose

(in each 5 mL tsp)  
Dextromethorphan HBr, USP 10 mg.....Cough Suppressant  
Guaifenesin, USP 100 mg.....Expectorant

### Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

### Warnings

- Do not use**
- in a child under 12 years of age
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache.  
These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

### Other information

- Keep carton for full Direction for use.
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

\* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-039



\*COMPARE TO THE  
ACTIVE INGREDIENTS IN  
ROBITUSSIN® PEAK  
COLD COUGH &  
CHEST CONGESTION DM



\*COMPARE TO THE  
ACTIVE INGREDIENTS IN  
ROBITUSSIN® PEAK  
COLD COUGH &  
CHEST CONGESTION DM

# Tussin DM

Dextromethorphan HBr /  
Guaifenesin

**NON-DROWSY**

**COUGH SUPPRESSANT  
EXPECTORANT**

Helps Loosen  
Chest Congestion

Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

# Tussin DM

Dextromethorphan HBr /  
Guaifenesin

**NON-DROWSY**

**COUGH SUPPRESSANT  
EXPECTORANT**

Helps Loosen  
Chest Congestion

Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE  
PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR  
DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG  
MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256





\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

# Tussin DM

Dextromethorphan HBr/  
Guafenesin

**COUGH SUPPRESSANT/  
EXPECTORANT**

**NON-DROWSY**  
Helps Loosen  
Chest Congestion  
Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

## Drug Facts

**Active ingredients (in each 5 mL tsp)**  
Dextromethorphan HBr, USP 10 mg. Cough Suppressant  
Guafenesin, USP 100 mg. Expectorant

**Uses** ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold  
■ helps loosen phlegm (mucous) and thin bronchial secretions to drain bronchial tubes

### Warnings

**Do not use** ■ in a child under 12 years of age  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

■ cough that occurs with too much phlegm (mucus)  
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonsful every 4 hours
children under 12 years	do not use

### Other information

■ store at 20-25 °C (68-77 °F) ■ do not refrigerate  
■ dosage cup provided ■ sodium 3 mg per teaspoonful  
■ See Carton for full labeling

**Inactive ingredients** anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LR-069

LOT:

EXP:

DO NOT USE IF PRINTED SEAL UNDER CAP IS  
TORN OR MISSING

## Drug Facts

### Active ingredients Purpose

(in each 5 mL tsp)  
Dextromethorphan HBr, USP 10 mg.....Cough Suppressant  
Guaifenesin, USP 100 mg.....Expectorant

### Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

### Warnings

- Do not use**
- in a child under 12 years of age
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

### Other information

- Keep carton for full Direction for use.
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

**Inactive ingredients** Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

\* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-039



\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM



\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

# Tussin DM

Dextromethorphan HBr /  
Guaifenesin

**NON-DROWSY**

**COUGH SUPPRESSANT  
EXPECTORANT**

Helps Loosen  
Chest Congestion

Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

# Tussin DM

Dextromethorphan HBr /  
Guaifenesin

**NON-DROWSY**

**COUGH SUPPRESSANT  
EXPECTORANT**

Helps Loosen  
Chest Congestion

Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

SATISFACTION GUARANTEED  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



**DISCOUNT**  
**drug mart**  
WALK-INS

\*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

# Tussin DM

Dextromethorphan HBr/  
Guaifenesin  
**COUGH SUPPRESSANT/  
EXPECTORANT**

NON-DROWSY  
Helps Loosen  
Chest Congestion  
Cough Formula  
for ages 12 & over

8 FL OZ (237 mL)

**DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING**

**Drug Facts**

**Purpose**  
Cough Suppressant  
Expectorant

**Active ingredients (in each 5 mL tsp)**  
Dextromethorphan HBr, USP 10 mg  
Guaifenesin, USP 100 mg

**Uses** ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold  
■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

**Warnings**  
**Do not use** ■ in a child under 12 years of age  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have**  
■ cough that occurs with too much phlegm (mucus)  
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

**Stop use and ask a doctor** if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoons every 4 hours
children under 12 years	do not use

**Other information**  
■ store at 20-25 °C (68-77 °F) ■ do not refrigerate  
■ dosage cup provided ■ sodium 3 mg per teaspoonful  
■ See Carton for full labeling

**Inactive ingredients** anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

**Questions?** Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

\*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

**SATISFACTION GUARANTEED**  
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_

LR-069

## TUSSIN DM

dextromethorphan hydrobromide, guaifenesin liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53943-503
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>DEXTROSE</b> (UNII: IY9XDZ35W2)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	

### Packaging

Marketing Start      Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-503-26	237 mL in 1 PACKAGE; Type 0: Not a Combination Product	11/19/2014	
<b>Marketing Information</b>				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	11/19/2014	

**Labeler** - Discount Drug Mart (047741335)

Revised: 12/2025

Discount Drug Mart