

TUSSIN DM- dextromethorphan hydrobromide, guaifenesin liquid
Discount Drug Mart

Drug Facts

Active ingredients (in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg

Guaifenesin, USP 100 mg

Purpose

Cough Suppressant

Expectorant

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING.

Do not use

- in a child under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before taking if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, bronchitis or emphysema

Stop use and ask a doctor if

cough lasts more than 7 days, comes back or is accompanied by fever, rash or

persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- **Keep carton for full Direction for use.**
- store at 20°-25°C (68°-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients

Anhydrous citric acid, dextrose, FD and red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at **1-877-798-5944**

DISCOUNT drug mart FOOD FAIR Tussin DM product label

***COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH AND CHEST CONGESTION DM**

DISCOUNT

drug mart

FOOD FAIR

Tussin DM

Dextromethorphan HBr/ Guaifenesin

NON-DROWSY

COUGH SUPPRESSANT

EXPECTORANT

Helps Loosen

Chest Congestion

Cough Formula

for ages 12 & over

8 FL OZ (237 mL)

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-039

SATISFACTION GUARANTEED

IF DISSASIFIED , RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING

Drug Facts

Active ingredients **Purpose**
(in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg.....Cough Suppressant
Guaifenesin, USP 100 mg.....Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

- Do not use** ■ in a child under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- **Keep carton for full Direction for use.**
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-039



*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM



*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

Tussin DM

Dextromethorphan HBr /
Guaifenesin

NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT**

Helps Loosen
Chest Congestion

Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

Tussin DM

Dextromethorphan HBr /
Guaifenesin

NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT**

Helps Loosen
Chest Congestion

Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256





*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

Tussin DM

Dextromethorphan HBr/
Guaifenesin

**COUGH SUPPRESSANT/
EXPECTORANT**

NON-DROWSY
Helps Loosen
Chest Congestion
Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)
Dextromethorphan HBr, USP 10 mg. Cough Suppressant
Guaifenesin, USP 100 mg. Expectorant

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
■ helps loosen phlegm (mucous) and thin bronchial secretions to drain bronchial tubes

Warnings
Do not use ■ in a child under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ cough that occurs with too much phlegm (mucus)
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonsful every 4 hours
children under 12 years	do not use

Other information
■ store at 20-25 °C (68-77°F) ■ do not refrigerate
■ dosage cup provided ■ sodium 3 mg per teaspoonful
■ See Carton for full labeling

Inactive ingredients anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LR-069

LOT:

EXP:

res

DO NOT USE IF PRINTED SEAL UNDER CAP IS
TORN OR MISSING

Drug Facts

Active ingredients **Purpose**
(in each 5 mL tsp)

Dextromethorphan HBr, USP 10 mg.....Cough Suppressant
Guaifenesin, USP 100 mg.....Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

- Do not use** ■ in a child under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoonfuls every 4 hours
children under 12 years	do not use

Other information

- **Keep carton for full Direction for use.**
- store at 20-25°C (68-77°F)
- do not refrigerate
- dosage cup provided
- sodium 3 mg per teaspoonful

Inactive ingredients Anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

* This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.

BX-039



*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM



*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

Tussin DM

Dextromethorphan HBr /
Guaifenesin

NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT**

Helps Loosen
Chest Congestion

Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

Tussin DM

Dextromethorphan HBr /
Guaifenesin

NON-DROWSY

**COUGH SUPPRESSANT
EXPECTORANT**

Helps Loosen
Chest Congestion

Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



DISCOUNT
drug mart
WALK-INS

*COMPARE TO THE ACTIVE INGREDIENTS IN ROBITUSSIN® PEAK COLD COUGH & CHEST CONGESTION DM

Tussin DM

Dextromethorphan HBr/
Guaifenesin
**COUGH SUPPRESSANT/
EXPECTORANT**

NON-DROWSY
Helps Loosen
Chest Congestion
Cough Formula
for ages 12 & over

8 FL OZ (237 mL)

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredients (in each 5 mL tsp)
Dextromethorphan HBr, USP 10 mg. Cough Suppressant
Guaifenesin, USP 100 mg. Expectorant

Uses ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings
Do not use ■ in a child under 12 years of age
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ cough that occurs with too much phlegm (mucus)
■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than 6 doses in any 24-hour period

Age	Dose
adults & children 12 years & over	2 teaspoons every 4 hours
children under 12 years	do not use

Other information
■ store at 20-25 °C (68-77 °F) ■ do not refrigerate
■ dosage cup provided ■ sodium 3 mg per teaspoonful
■ See Carton for full labeling

Inactive ingredients anhydrous citric acid, dextrose, FD&C red # 40, flavor, glycerin, high fructose corn syrup, menthol, purified water, saccharin sodium, sodium benzoate.

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944.

LR-069

*This product is not manufactured or distributed by Pfizer, owner of the registered trademark Robitussin® Peak Cold.
SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

LOT: EXP:

TUSSIN DM

dextromethorphan hydrobromide, guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-503
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DEXTROSE (UNII: IY9XDZ35W2)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MENTHOL (UNII: L7T10EIP3A)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

Packaging

Marketing Start Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-503-26	237 mL in 1 PACKAGE; Type 0: Not a Combination Product	11/19/2014	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	11/19/2014	

Labeler - Discount Drug Mart (047741335)

Revised: 12/2024

Discount Drug Mart