ALKA-SELTZER PLUS SEVERE COLD AND FLU POWERFAST FIZZchlorpheniramine maleate, acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide tablet, effervescent Bayer HealthCare LLC.

Alka-Seltzer Plus Severe Cold & Flu PowerFast Fizz UI 1614461 Drug Facts

Active ingredients (in each tablet) Purposes

Acetaminophen 250 mg.......Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg......Antihistamine
Dextromethorphan hydrobromide 10 mg.....Cough suppressant
Phenylephrine hydrochloride 5 mg.....Nasal decongestant

Uses

- · temporarily relieves these symptoms due to a cold or flu:
- \cdot minor aches and pains \cdot headache \cdot cough
- · sore throat · runny nose · sneezing
- · nasal and sinus congestion
- · temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- \cdot more than 4,000 mg of acetaminophen in 24 hours
- \cdot with other drugs containing acetaminophen
- \cdot 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe

allergic reactions. Symptoms may include:

- \cdot skin reddening \cdot blisters \cdot rash \cdot hives
- \cdot facial swelling \cdot asthma (wheezing) \cdot shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than

2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease heart disease high blood pressure
- ◆ thyroid disease ◆ diabetes ◆ glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium restricted diet

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in childre

Stop use and ask a doctor if

· pain, cough, or nasal congestion gets worse or lasts more than

7 days

- · fever gets worse or lasts more than 3 days
- · redness or swelling is present
- · new symptoms occur
- · cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

· nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions

Directions

- · do not take more than the recommended dose
- \cdot adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
- · children under 12 years: do not use

Other information

- each tablet contains: potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

Inactive ingredients anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavors, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

Questions or comments

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)



ALKA-SELTZER PLUS SEVERE COLD AND FLU POWERFAST FIZZ

chlorpheniramine maleate, acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide tablet, effervescent

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0022
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	2 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	250 mg	

Inactive Ingredients		
Ingredient Name	Strength	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
DIMETHICONE (UNII: 92RU3N3Y1O)		
CALCIUM SILICATE (UNII: S4255P4G5M)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		

MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MANNITOL (UNII: 3OWL53L36A)	
POVIDONE (UNII: FZ989GH94E)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Product Characteristics				
Color	white (Speckled)	Score	no score	
Shape	ROUND	Size	25mm	
Flavor		Imprint Code	ASP;FLU	
Contains				

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0022- 01	10 in 1 CARTON	04/01/2020	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:0280-0022- 02	36 in 1 CARTON	09/15/2021	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:0280-0022- 03	3200 in 1 BOX	02/01/2022	
3		1 in 1 BOX; Type 0: Not a Combination Product		
4	NDC:0280-0022- 04	24 in 1 CARTON	03/09/2022	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:0280-0022- 05	12 in 1 CARTON	03/06/2023	
5		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2020	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2025 Bayer HealthCare LLC.